

EFFECT OF SOYABEAN DIET ON THE SERUM IMMUNOGLOBULIN PROFILE AND CD₄⁺ COUNT OF MALE DIABETIC RATS CHALLENGED WITH K.PNEUMONIAE

***ANDREW C. NWAKA¹, PETER N. UZOEGBU², ONYECHI OBIDOA³, UGWU OKECHUKWU P.C.⁴, OSSAI EMMANUEL C.⁴ and CHIGOR VINCENT N.⁵**

¹Department Of Biochemistry, Anambra State University, Uli, Nigeria.

²Tropical Diseases Research Unit, Department of Biochemistry, University of Nigeria, Nsukka, Nigeria.

³ Department of Biochemistry, Kogi State University, Anyingba, Kogi State.

⁴Department of Biochemistry, University of Nigeria, Nsukka.

⁵Department of Microbiology, University of Nigeria, Nsukka.

*Corresponding Author Email: andynwaka@yahoo.com

ABSTRACT

In this study, the effect of soyabean supplemented diet on the immunoglobulin profile and CD₄⁺ count of male diabetic rats challenged with *Klebsiella pneumoniae* were evaluated in a total of sixty-four mature albino rats, weighing 150-250 grams. The rats were separated into eight groups of eight rats each. The groups were as follows: A- rats fed on rat feed, B-rats fed on rat feed and soyabean meal, C-diabetes-induced rats fed on rat feed, D-diabetes-induced rats fed on rat feed and soyabean meal, E-rats fed on rat feed but challenged with *Klebsiella pneumoniae*, F- rats fed on rat feed and soyabean meal challenged with *Klebsiella pneumoniae*, G-diabetes induced rats fed on normal rat feed but challenged with *Klebsiella pneumoniae* and H-diabetes-induced rats fed on rat feed and soyabean meal but challenged with *Klebsiella pneumoniae*. The study lasted for 28 days. At the end of 28 days, the rats were bled and blood collected, processed and the serum obtained used for assessment of immunoglobulin profile and CD₄⁺ T-cell count. Immunoglobulin profile was carried out using immunoturbidimetric assay with Randox^(R) kit, while CD₄⁺ T-cell count was done using CY-flow machine. Results showed that there were significant increase ($P < 0.05$) in serum IgA, IgM, IgG levels and CD₄⁺ T-cell of diabetic rats fed on normal feed and soyabean meal when compared with their counterparts fed with only normal feed. Results of the study suggest that soyabean diet could assist in boosting the serum immunoglobulin levels and CD₄⁺ cell of diabetic rats and hence protecting the rats from *K. pneumoniae* infection.

KEY WORDS

Diabetes mellitus, Immunoglobulins, CD₄⁺-T-cell, *K. pneumoniae* and Soyabeans.

INTRODUCTION

Diabetes mellitus, often referred simply as diabetes, is a syndrome characterized by disordered metabolism and abnormally high fasting blood sugar (hyperglycaemia) resulting from insufficient levels of the hormone, insulin or its insensitivity (Tierney et al., 2002). It is a chronic disease characterized by elevated blood glucose levels and disturbances in carbohydrate, fat and

protein metabolism (Sky, 2000, Rother, 2007). The World Health Organization, recognizes three main form of diabetes mellitus type I, type II and gestational diabetes (occurring during pregnancy) (WHO, 1999).

Type I diabetes mellitus formerly called juvenile onset or insulin dependent diabetes mellitus (IDDM) is characterized by loss of the insulin

producing beta cell of the islets of langerhans of the pancreas, leading to a deficiency of insulin. The main cause of this loss of β - cell is a T-cell mediated autoimmune attack (Rother, 2007). There is no known preventive measure which can be taken against type I diabetes. While type 2 diabetes mellitus, also known as non-insulin dependent diabetes mellitus (NIDDM), is due to insulin resistance or reduced insulin sensitivities combined with reduced insulin secretion (Rosenbloom and Silverstain, 2003). Furthermore, gestational diabetes mellitus (GDM) resembles type 2 diabetes in several respects, involving a combination of relatively inadequate insulin secretion and responsiveness (WHO, 1999). In Nigeria, diabetes mellitus has emerged as one of the biggest health problems and its prevalence is increasing at an alarming rate (Unadike, 20120).

Soyabean, is a plant of the family leguminosae. Legumes are noted for their relatively high protein content when compared with cereals, tubers, fruits and vegetables. Soyabean contains about 40% protein, 18% fat and has a high concentration of minerals such as calcium, iron etc (Anosike et al., 2007).

Infections, no matter how mild, have adverse effects on nutritional status. The significance of these effects depends on the nutritional status of the individual, the nature and duration of the infection and the diet during the recovery period (Scrimshaw and Sangiovanni, 1997). Furthermore, it is not surprising that protein deficiency is so consistently observed to interfere with resistance to infection because most immune mechanisms are dependent on cell replication or the production of active protein compounds. However, because protein cannot be synthesized without a balance of essential amino acids, experimental amino acid deficiencies have the same effect as protein. Essentially all forms of

immunity have been shown to be affected by protein-energy malnutrition in young children, depending on the severity of the protein deficiency relative to energy. The effects include impaired antibody formation (Pretorius and De Villiers, 1962, Reddy and Srikantia, 1964 and Reddy et al., 1977), decreased serum immunoglobulin (Aref et al., 1977 and Watson et al., 1987), decreased thymic function (Watts, 1969, Purtilo and Connor, 1975 and Chandra et al., 1982) and splenic lymphocytes, delayed cutaneous hypersensitivity (Simythe et al., 1971 and Chandra, 1974), decreased complement formation (Sythe et al., 1971, Suskind et al., 1976), decreased interferon and effects on nonspecific mechanisms that include anatomic barriers and secretory substances such as lysosomes and mucus (Scrimshaw and Sangiovanni, 1997).

In this study, the effect of soyabean diet on the serum immunoglobulin profile (i.e. IgM, IgG and IgA) and CD_4^+ count of male diabetic rats challenged with *Klebsiella pneumoniae* (a Gram-negative bacteria) was evaluated.

MATERIALS

Chemical used: All the chemical used in this research were of analytical grade.

Animal used: Mature apparently healthy male albino rats weighing 150-250 grams were procured from the Animal House of Faculty of Veterinary Medicine, University of Nigeria, Nsukka campus. The animals were allowed to acclimatize for fourteen days before the commencement of the study.

Soyabean meal: Rats on soyabean meal were given 9.6g/kg body weight of soyabean powder dispersed in water by intubation each morning before the animals were fed for the day.

Rat feed: The normal rat feed used in this study is pelleted growers mash of Vital^(R) feed. It was produced by Grand Cereals and Oil Mills, Jos, Nigeria.

Pathogenic Organism: *Klebsiella pneumoniae* was used as the pathogenic organism in this study. Its stock was procured from the Department of Veterinary Microbiology, Faculty

of Veterinary Medicine, University of Nigeria, Nsukka.

METHODS

Experimental Design

Sixty-four mature male albino rats, weighing 150-250grams were used in the study. The rats were separated into eight groups of eight rats each and kept in eight different standard cages as follows:

Groups	Treatment
A	Rats fed on rat feed
B	Rats fed on rat feed and soyabean meal
C	Diabetes-induced rats fed on rat feed
D	Diabetes-induced rats fed on rat feed and soyabean meal.
E	Rats fed on rat feed but challenged with <i>Klebsiella pneumoniae</i>
F	Rats fed on rat feed and soyabean meal challenged with <i>Klebsiella pneumoniae</i>
G	Diabetes-induced rats fed on normal rat feed but challenged with <i>Klebsiella pneumoniae</i>
H	Diabetes-induced rats fed on rat feed and soyabean meal but challenged with <i>Klebsiella pneumoniae</i>

All the protocols as approved by Institutional Animal Ethics Committee (IAEC) were observed in the study. At the end of the 28days, the rats were bled from the retro-bulbar plexus of the medical canthus of the eye. The blood samples were collected into EDTA tubes and plain sample bottles. The EDTA acted as anticoagulant. The blood samples in the EDTA bottles were used for CD₄⁺ T-cell counts, while the blood sample collected in the plain tubes without EDTA were allowed to stand for 30 minutes and centrifuged at the speed of 3,000 revolutions per minutes for 10minutes. The serum was extracted and used for IgA, IgG and IgM assays using standard biochemical methods.

Induction of Diabetes

Diabetes was induced in the rats by injecting 180mg/kg body weight of alloxan monohydrate intraperitoneally in 0.9% w/v NaCl (normal saline) after fasting the rats for 12hours. Seven days after

the alloxan administration, the rats were fasted for 12hours and their blood glucose level was measured. Rats having blood glucose level of 400mg/dl and above were selected and used as diabetic rats in this study. Rats that were described as fasted were deprived of food for at least 12hours but were allowed free access (*ad libitum*) to drinking water.

Pathogen Challenge

The rats were challenged with *Klebsiella pneumoniae* by intraperitoneal administration of 0.4ml of the bacterial suspension containing 10⁸ c.f.u/ml in 0.9% (w/v) normal saline on the 22nd day of the study.

Proximate Analysis

The proximate composition of soyabean and the normal animal feed used in the study were evaluated. Their protein content were determined by micro khfelldal method as was described by

AOAC (2000) and Changsam (2003), while their ash, moisture and fats contents were determined by gravimetric method as described by AOAC (2000). Crude fiber were determined by gravimetric method as described by James (1995). The total carbohydrate were then estimated using the formula as was given by Bemiller (2003).

Immunoglobulin Profile and CD_4^+ T-cell Determination

The serum Igm, IgA and IgG were determined using immunoturbidimetric assay (Whicher et al., 1983) with Randox^(R) kit, while CD_4^+ count was carried out using CY Flow SL Green Machine.

RESULTS

The result of proximate analysis for the soyabean powder used in this study (Fig.1) indicated that its moisture content was 7.40%, protein 48.22%,

fibre 3.80%, fat 15.90%, ash 4.60% and carbohydrate content 20.08%. Also, the result of proximate analysis of normal feed (pelleted growers mash of vital^(R) feed) used in this study (Fig. 2.) showed that the feed's moisture content was 16.75%, fat 5.80%, ash 7.25%, fibre 2.60% and protein 17.95% and crbohydrate content was 49.65%.

Furthermore, when the different groups of rats were challenged with *Klebsiella pneumoniae*, it was observed that only the group G with diabetic rats fed with normal feed came down with the infection. This was indicated by the sickly appearance of the rats in the group and mortality of half of the rats in the group G before the end of the study.




Fig. 1: Soyabean proximate composition




Fig. 2: Normal feed proximate composition




Fig. 3: Mean serum IgM concentration for different groups of rats used




Fig. 4: Mean serum IgA concentration for different groups of rats used




Fig. 5: Mean serum IgG concentration for different groups of rats used




Fig. 6: Mean serum CD₄⁺ concentration for different groups of rats used

The result of the mean serum immunoglobulin M (IgM) for different groups of rats used in the study (**Fig 3**) indicated that the mean serum IgM of groups E, F and H were significantly higher ($P < 0.05$) than that of the control (group A). The result (**Fig. 3**) also showed that there were significant reductions ($P < 0.05$) in the serum IgM concentrations of diabetic rats fed on normal feed only, group C (79.21 ± 20.73 mg/dl) and diabetic rats fed on normal feed but challenged with *Klebsiella pneumoniae*, group G (62.49 ± 22.96 mg/dl) when compared to control, group A (120.55 ± 9.29 mg/dl). **Fig.3** indicated that the mean serum IgM of diabetic rats fed on soyabean diet (group D) was significantly higher ($P < 0.05$) than that of diabetic rats fed on normal feed. Result (**Fig.3**) also showed that the mean serum IgM of diabetic rats fed on soyabean diet challenged with *klebsiella pneumoniae* (group H) was significantly higher ($P < 0.05$) than that of diabetic rats fed on normal feed but challenged with *Klebsiella pneumoniae* (group G). **Fig.3** also

indicated that the mean serum IgM of normal rats fed on soyabean diet (group B) and normal rats fed on soyabean diet but challenged with *Klebsiella pneumoniae* (group F) were higher than their counterparts fed on normal feed (groups A and E respectively). The increases were statistically not significant ($P > 0.05$).

The result of the mean serum IgA concentrations for the different groups of rats used in this study (**Fig. 4**) revealed that the IgA levels of normal rats fed on normal rat feed and soyabean meal but challenged with *klebsiella pneumoniae*, group E (188.71 ± 15.09 mg/dl) and diabetic rats fed on normal rat feed and soyabean meal but challenged with *Klebsiella pneumoniae*, group H (193.67 ± 19.43 mg/dl) were significantly higher ($P < 0.05$) than that of the control, group A (169.61 ± 17.53 mg/dl). **Fig.4** further indicated that the mean serum IgA concentration of diabetic rats fed on normal feed, group C (124.12 ± 25.21 mg/dl) was significantly lower ($P < 0.05$) than those of the

control, group A ($169.61 \pm 17.53\text{mg/dl}$) and those of the diabetic rats fed on normal feed and soyabean meal, group D ($167.04 \pm 18.61\text{mg/dl}$).

Fig.4 also indicated that the mean serum IgA of rats fed on soyabean but challenged with *Klebsiella pneumoniae* (group F) and diabetic rats fed on soyabean but challenged with *Klebsiella pneumoniae* (group H) were significantly higher ($P < 0.05$) than their counterparts fed on normal feed (group E and G respectively).

The result of the mean serum IgG concentrations for the different groups of rats used in this study (**Fig.5**) showed that the mean serum IgG concentrations of normal rats fed on normal feed but challenged with *Klebsiella pneumoniae*, group E ($757.55 \pm 63.22\text{mg/dl}$) and normal rats but also challenged with *Klebsiella pneumoniae* group F ($899.31 \pm 56.23\text{mg/dl}$) were significantly higher ($P < 0.05$) than that of the control, group A ($599.96 \pm 45.06\text{mg/dl}$). **Fig.5** also revealed that the serum mean IgG concentration of diabetic rats fed on normal feed, group C ($311.57 \pm 53.36\text{mg/dl}$) was significantly reduced ($P < 0.05$) when compared to the control, group A ($597.96 \pm 45.06\text{mg/dl}$) and to that of the diabetic rats fed on normal rat feed and soyabean meal, group D ($502.94 \pm 41.54\text{mg/dl}$). Figure 5 further indicated that the mean serum IgG of rats fed on soyabean diet challenged with *Klebsiella pneumoniae* (group F) and diabetic rats fed on soyabean diet challenged with *Klebsiella pneumoniae* (group H) were significantly higher ($P < 0.05$) than their counterparts fed on normal feed only (groups E and G respectively). The result also revealed that there was no significant difference ($P < 0.05$) between the mean serum IgG of rats fed on soyabean diet (group B) and that of rats fed on normal feed only (group A).

The result of CD_4^+ cell count for the different groups of albino rats used in this study as shown in **Fig.6** indicated that the mean CD_4^+ cell count of

normal rats fed on normal feed and soyabean meal, group B ($331.25 \pm 22.50 \text{ cells}/\mu\text{l}$) and normal rats fed on soyabean diet but challenged with *Klebsiella pneumoniae* (group F) were significantly not different ($P > 0.05$) from their counterparts fed on normal feed only (groups A and E respectively). However, with the exception of group B, the mean CD_4^+ count of all the other groups were significantly lower ($P < 0.05$) than that of the control, group A. Moreover, the results shown in **Fig.6** indicated that there were significant reduction ($P < 0.05$) in CD_4^+ count of all the groups challenged with *Klebsiella pneumoniae* when compared with their counterparts not challenged with the pathogen. The results also showed that the mean CD_4^+ counts of diabetic rats fed on normal rat feed and soyabean meal, group D ($226.50 \pm 21.41 \text{ cells}/\mu\text{l}$) was significantly higher than that of the diabetic rats fed on only normal feed, group C ($161.75 \pm 28.74 \text{ cell}/\mu\text{l}$).

DISCUSSION

Over the years, most investigations on diabetes have concentrated on the effect of various diets or herbal extracts on glucose levels without adequate consideration of their effect on the immune response. However, the effect of soyabean diet on some immune markers (IgG, IgA, IgM and CD_4^+ cells) of diabetic rats were evaluated in this study.

The results of the immunoglobulin profile indicated significant increase ($P < 0.05$) in the IgM concentrations (**Fig.3**) of normal rats fed on normal rat feed but challenged with *Klebsiella pneumoniae* (group E), normal rats fed on normal feed and soyabean meal challenged with *Klebsiella pneumoniae* (group F) and diabetic rats fed on normal feed and soyabean meal but challenged with *Klebsiella pneumoniae* when compared to the control group (group A). This

suggested that the presence of *Klebsiella pneumoniae* (antigen), the state of health of the rats and the nutritive value of the soyabean diet used in the study could have likely been the reason for the increased IgM antibodies. This observation corroborated Edelman (1973) observation that in active immunity, the effective contact with antigens causes the host to produce antibodies in response to the antigens. When the IgM concentration of diabetic rats fed on normal rat feed (group C) was compared to the control, a significant reduction ($P < 0.05$) was observed in group C, suggesting that diabetes also causes depletion of the IgM level, thereby reducing immune competence in the rats. However, when the IgM concentration of diabetic rats fed on normal feed (group C) was compared with those of their counterparts fed on normal feed and soyabean meal (Group D), it was observed that the reduction in IgM concentration in group C was restored to normalcy with the incorporation of soyabean diet in group D, an observation that could be attributed to soyabean's high protein composition with its high content of essential amino acids. These amino acids could have helped in boosting the plasma proteins which could have led to increase in the IgM concentration, since immunoglobulins are synthesized by plasma cells. This observation is in accord with McGee and McMurray (1977) report that soyabean (protein) could help in boosting immunoglobulin concentration in the intestine of mice. Immunoglobulins are proteins. So the increase in IgM concentration observed in the diabetic rats fed on soyabean diet (group D) when compared to that of the diabetic rats fed on normal feed (group C) could be as a result of high essential amino acids present in the soyabean used.

Also, the results of IgA and IgG concentrations for normal rats fed on normal feed challenged with *Klebsiella pneumoniae* (group E), normal rats fed

on normal feed and soyabean meal challenged with *Klebsiella pneumoniae* (group F) and diabetic rats fed on normal feed and soyabean meal but challenged with *Klebsiella pneumoniae* (group H) indicated significant increases ($P < 0.05$) in IgA and IgG concentrations when compared with control group A (**Fig. 4 and 5**). This also suggested that the production of more antibodies (IgA and IgG) in both diabetic and non-diabetic rats (except in group G) as soon as they were challenged with *Klebsiella pneumoniae* in order to protect the rats from invading antigen could have been affected by the presence of *Klebsiella pneumoniae*, the state of health of the rats (i.e. whether diabetic or non-diabetic) and the nutritive value of the diet (soyabean) used. This observation is also in consonance with the report that the presence of antigen could trigger off the production of antibodies as protective measure of the host against invading antigen (Edelman, 1973).

It was also observed that more IgM, IgA, and IgG antibodies were synthesized in the groups in which soyabean meal was incorporated into their diet when compared with their counterparts fed on only normal feed. This suggests that soyabean diets could have helped in boosting their immunoglobulin levels, due to the presence of essential amino acids in the soyabean. These essential amino acids could have helped in boosting the immunoglobulin level. This observation also corroborates McMurray (1977) which stated that protein (essential amino acids) helps in boosting immunoglobulin concentration in the intestine of mice. Furthermore, the IgM, IgA and IgG concentrations of diabetic rats fed on normal feed (group C) were significantly reduced ($P < 0.05$) when compared to the control, an indication that diabetes causes depletion of immunoglobulin levels. This could be due to the diabetics' inability to utilize blood glucose for energy generation. Muscle proteins are therefore

metabolized for energy generation, leading to loss of muscle amino acids which could result to the shrinking of the muscles. The depletion of the muscle's amino acids could have caused the decrease in the immunoglobulin levels, since immunoglobulins are synthesized from amino acids. However, it was observed that the reduced immunoglobulin concentrations in group C was restored almost to normal level in diabetic rats fed on Soyabean diet (group D). The suspicion is that the essential amino acids in soyabean could have been replenished the lost amino acids for energy generation. The replenished essential amino acids from Soyabean Diet could have assisted in boosting the immunoglobulin concentration in the rats fed with Soyabean supplemented diet, since immunoglobulins are proteins.

Moreover, **Fig.6** indicated that with the exception of normal rats fed on normal rat feed and soyabean meal (group B), that the mean CD_4^+ count of all other groups (groups C, D, E, F, G and H) were significantly lower ($P < 0.05$) than the control (group A). This suggests that diabetes and the presence of pathogen (*Klebsiella pneumoniae*) could have caused reduction in CD_4^+ cells. The utilization of muscle protein for energy generation could have brought about the depletion of essential amino acids in the diabetic rats. This in turn could have caused the decreased CD_4^+ cell count in diabetic rats as observed, since CD_4^+ cells are produced from essential amino acids. Also results in **Fig.6** showed that both diabetic and non-diabetic rats challenged with *Klebsiella pneumoniae* but fed on soyabean incorporated diet had higher CD_4^+ count than their counterparts fed on only normal rat feed. This is because soyabean is rich in essential amino acids. These essential amino acids could have helped in replenishing the lost muscles and tissues amino acids during energy generation in diabetic rats fed in soyabean diet. The replenished essential amino

acids in diabetic rats fed on Soyabean supplemented diet could have boosted their CD_4^+ cell count when compared to that of the diabetic rats fed on normal feed. This should also be expected considering the high nutritive value of Soyabean (**Fig.1**) with high percentage of protein, minerals and vitamins including antioxidants (Regel et al., 2000; Anosike et al., 2007; and Villegas and Gao, 2008). The above observation also corroborates the report that decreased dietary intake of antioxidants could lead to a concomitant decrease in CD_4^+ T-cells (Oguntibeju et al., 2005 and 2006). Since the CD_4^+ T-cells, perform a central and coordinating role in the immune response (Hughes et al., 1997; Vajpayee et al., 2009), and Soyabean diet helped in raising the CD_4^+ cells in diabetic rats as observed in this study, Soyabean diet therefore could help in the general boosting of the immune system of diabetics.

CONCLUSION

This study revealed significant increases ($P < 0.05$) in serum IgA, IgM and IgG levels and CD_4^+ T-cells concentration of diabetic rats fed with normal feed supplemented with soyabean meal (group D) and the rats infected with *K. pneumoniae* and fed with normal feed and supplemented with soyabean meal (group H) when compared with their counterparts fed with only normal feed (groups C and F). Moreover, it was observed that when various groups of rats were challenged intraperitoneally with 0.4ml of *Klebsiella pneumoniae* suspension containing 10^8 c.f.u/ml, establishment of infection and high mortality rate of the infected rats were observed in diabetic rats fed with normal feed only (group G), whereas no mortality was observed in diabetic rats fed with normal rats feed supplemented with soyabean meal (group H). These results suggest that soyabean diet could be useful in the general

boosting of the immune system and in the general management of diabetes mellitus.

SUGGESTIONS FOR FURTHER STUDIES

It is suggested that further studies be carried out as follows:

- ❖ Inducing diabetes that could last for more than 3 months in order to appreciate the effect of soya protein more, in chronic complications of diabetes.
- ❖ Feeding the animals with soyabean diet for more than 3 months (i.e. longer period), since the longer the animal is exposed to the diet, the more the effect of the diet could be discovered.
- ❖ Using more than two different pathogenic organisms that are infectious to albino rats in subsequent study, to confirm these results.

ACKNOWLEDGEMENT

We are grateful to the following for assistance rendered towards the success of this research work: Prof. L.S.U. Ezeanyika and Prof. I.N.E. Onwurah, both from Department of Biochemistry, University of Nigeria, Nsukka. Also to Prof. J. I. Ihedioha, Prof. S.V.O Shoyinka and Prof. K. F. Chah, all from the Faculty of Veterinary Medicine, University of Nigeria, Nsukka. We will also not forget the Head of Hematological Section of Nnewi Teaching Hospital, Mr. A. Ozoabia, who assisted us in carrying out CD_4^+ count with their machine.

REFERENCES

- Anosike, C.A., Ezeanyika, L.U.S. and Obidoa, O. (2007). Effect of roasted soyabean (glycine max) diet on the histology of selected rat tissues. *Bio-Research*, 5 (2): 237-240.
- AOAC (2000). Official method of analysis of Association of Official Analytical Chemists. Washington D.C., 16th Edition. Pp. 12-18.
- Aref, G.H., el-Din, M.k. and Hassan, A.J., (1970). Immunoglobulins in Kwashiorkor. *Journal of Tropical Medicine and Hygiene*, 73:186-191.

- Bemiller, N.J. (2003) *Carbohydrate Analysis*. Food Analysis. Third edition. Kluwer Academic publisher, New York Pp. 143-146.
- Chandra, R.K., Gupta, S. and Singh, H. (1982). Inducer and suppressor T-cell subsets in protein-energy malnutrition: analysis by monoclonal antibodies. *Nutrition Research*, 2:21-26.
- Changsam, K.C. (2003). *Protein Analysis*. Food Analysis. Third edition kluwer Academic/Plenum publisher, New York. Pp. 134-135.
- Hughes, M. D., Johnson, V. A. and Hirsch, M. S. (1997). Monitoring plasma HIV-I RNA levels in addition to CD4+ lymphocyte count improves assessment of antiretroviral therapeutic response. *Annals of Internal Medicine*, 126:929-938.
- James, O.K. (1995). Gravimetric method of analysis employed in twenty different cereals and legumes, Wayton, U.K.
- Oguntibeju, O. O., Van den Heever, W. M. J. and Van Schakwyk, F. E. (2005). An analysis of based line dietary intake of HIV-positive/AIDS patients. *South African Journal of Medical Technology*, 19 (2): 3-9.
- Oguntibeju, O. O., Van den Heever, W. M. J. and Van Schakwyk, F. E. (2006). The effect of a liquid nutritional supplement on viral load and haematological parameters of HIV-positive/AIDS patients. *British Journal of Biomedical Science*, 63 (3): 134-139.
- Pretorius, P.J. and Villiers, L.S. (1962) Antibody response in children with Protein Malnutrition. *American Journal of Clinical Nutrition*, 10: 379-383.
- Purtilo, D.T. and Connor, D.H. (1975). Fatal infection in protein-calorie malnourished children with thymolymphatic atrophy. *Archives of Disease in Childhood* 50: 149-152
- Reddy, V. and Srikantia, S.G. (1964). Antibody response in Kwashiorkor. *Indian Journal of Medical Research*, 52: 1154-1158.
- Reddy, V., Bhaskaram, C. and Raghuramulu, N. (1977). Immunological responses in malnourished children. *Indian Journal of Padiatrics*, 14: 255- 258
- Regel, F. J., Fraser, G. D., Weeks, E. C. and Greenberg, A. N. (2000). Dietary phytoestrogen have anti-inflammatory activity in guinea pig. *Proceedings of the Society for Experiment Biology and Medicine*. 223 (4): 372-378
- Rosenbloom, A. and Silverstain, J.H., (2003) Type 2 Diabetes in children and Adolescents: A Clinicians Guide to Diagnosis, Epidemiology, Pathogenesis, Prevention and Treatment. American Diabetes Association, U.S.I.,
- Rother, K.I. (2007). "Diabetes Treatment Bridging the Divide". *The New England Journal of Medicine*, 356 (150): 1499-1501.

- Scrimshaw, N.S. and san Giovanni, J.P. (1997). Synergism of Nutrition, Infection and Immunity. *American Journal of clinical Nutrition*, 66:464s-477s.
- Sky, G.M. (2000). Kinetics of insulin secretion underlying notabolism events in diabetes mellitus. *Diabetes mellitus: A fundamental and clinical text*. Lippicott willians and Wilkins, Plucadelphia Pp. 2-11.
- Smythe, P.M., Brereton-stiles, G.G. and Grace, H.J. (1971). Thymolymphatic deficiency and depression of cell mediated immunity in protein-calorie malnutrition. *Lancet*, 2: 939-943.
- Suskind, R., Edelman, R., KulapongsP., Pariyanonda, A. and Sirisinha, S. 91976). Complement activity in Children with Protein Calorie Malnutrition. *American Journal of Clinical Nutrition*, 29: 1089-1092.
- Tierney, I.M., Mcphee, S.J. and Papadakis, M.A. (2002). *Current Medical Diagnosis and Treatment of Diabetes*. Large Medical books. McGraw-Hill, New York.Pp. 1203-1215.
- Unadike, B.c. (2010). Awareness and knowledge about diabetes mellitus among nursing students in the Niger Delta region of Nigeria. *Middle East Journal of Nursing*, 4(1): 1-8.
- Vajpayee, M., Kaushik, S., Sreenivas, V., Mojumdar, K., Mendiratta, N. and Chauhan, K. (2009). Role of immune activation in CD4+ T-Cell depletion in HIV –I infected Indian patients. *European Journal of Clinical Microbiology and Infectious Diseases*, 28: 69-73.
- Villegas, R. and Gao, Y. T. (2008). Legume and soy food intake and incidence of
- type 2 diabetes in the Shanghai womens health study. *American Journal of*
- *Clinical Nutrition*. 87 (1): 162-167.
- Watson, B.B, Reyes, M.A. and Mc Murray, D.N., (1978) Influence of malnutrition on the concentration of IgA, lysozymes, amylase and aminopeptidase in children's tears. *Proceedings of the Society for Experiment Biology and Medicine*, 157: 215-219.
- Watts, T. (1969). Thymus weights in malnourished children. *Journal of Tropical Pediatrics*, 15:155-158.
- Whicher, J.J., Price, C.P. and Spencer, K. (1983). Immunoturbidimetric assay. *Critical Reviews in Clinical Laboratory Science*, 18: 213-216.
- WHO (World Health Organization) (1999). Department of Non Communicable Disease Surveillance (1999). *Definition, Diagnosis and Classification of Diabetes Mellitus and its complications*.



***Corresponding Author:**

Andrew C. Nwaka, Ph.D,
Department of Biochemistry,
Anambra State University,
Uli, Anambra State, Nigeria.
E-mail: andynwaka@yahoo.com;
Phone: +234 (0) 8037834171.