

CHRONIC EFFECT OF ETHIDIUM BROMIDE ON OVARIAN HISTOARCHITECHTURE OF ALBINO MICE

MANISHA MATHUR

Department of Zoology, G.N.Khalsa College, Matunga, Mumbai, India

*Corresponding Author Email: manishakmathurs@yahoo.co.in

ABSTRACT

Adult Mice (B.Wt. 30 gms & 40gms) were treated at a 5mg/kg B.Wt. and 10 mg/kg B.Wt. of Ethidium Bromide for 10 days in drinking water. Control animals were given equal dose of deionised water. Qualitative changes were studied in Ovary. Histopathological changes in Ovary, Primary Follicles, Secondary Follicles, Tertiary Follicles, Graffian follicles, etc. were evaluated. Significant changes were noticed. These changes were characterized by disorganization of follicles and degenerative changes were noticed. Slight change in ovarian weight was recorded. Body weight of the animal was slightly altered after challenge with Ethidium Bromide.

KEY WORDS

Toxicity, Ethidium Bromide, Ovary, Histoarchitectur, Albino Mice.

INTRODUCTION

Mammalian Ovary exhibits seasonal and Cyclic Changes. In non-primate species these are synchronised with estrous and in primates with the menstrual cycle. The mammalian ovary functions as a cytogenous gland, i.e., it has both endocrine and exocrine functions. Ethidium Bromide forms one of the sources of toxicants in the laboratories. Ethidium bromide is an intercalating agent commonly used as a fluorescent tag (nucleic acid stain) in molecular biology laboratories for techniques such as agarose gel electrophoresis. It is commonly abbreviated as "EtBr", which is also an abbreviation for bromoethane. When exposed to ultraviolet light, it will fluoresce with an orange colour, intensifying almost 20-fold after binding to DNA. Under the name homidium, it has been commonly used since the 1950s in veterinary to treat trypanosomosis in cattle, a disease caused by trypanosomes (1). The high incidence of antibiotic resistance makes this treatment impractical in some areas, where the related isometamidium chloride is used instead (2, 3, 5). Ethidium bromide may be a mutagen, carcinogen or teratogen although this depends on the organism and conditions (4, 6, and 8). The National Toxicology

Program states it is nonmutagenic in rats and mice. (7, 9, 10) Ethidium bromide (Homidium brand) use in animals to treat trypanosome infection suggests that toxicity and mutagenicity are not high. Studies have been conducted in animals to evaluate EtBr as a potential antitumorigenic chemotherapeutic agent.[11] Its chemotherapeutic use is due to its toxicity to mitochondria.[12] The above studies do not support the commonly held idea that ethidium bromide is a potent mutagen in humans, but they do indicate that it can be toxic at high concentrations. No study has been done on the histopathological changes in the ovary of Albino mice due to Ethidium Bromide toxicity. Hence the present study has been done.

MATERIAL AND METHODS

Adult males and females of swiss albino mice weighing 30 gms and 40 gms were used as a model in the present study. Two groups of mice were given Ethidium Bromide at a dose level of 5mg/kg body wt. and 10 mg/kg body wt. for 10 days. Experimental animals were given sodium fluoride orally through drinking water.

Animals of experimental and control group were sacrificed on tenth day of treatment by cervical

dislocation. The contralateral ovaries of experimental and control group of mice were fixed in formalin for 24 hrs. They were dehydrated, in graded EtOH series, cleared in xylene, in filtered with and embedded in pure filtered paraffin wax (M.P.58 degree centigrade.) Deparaffinised sections (5-7 microns) were stained by haematoxylin and eosin to monitor the extent of changes in the ovary histoarchitecture. Every alternate section of the ovary was microscopically examined and appropriate areas were microphotographed and enlarged.

RESULTS AND DISCUSSIONS

The ovary of control mice weighed 0.536 mgs (mean value). Necrosis, cirrhosis, and ischemia occurred but was not marked in control animals (FIG:1&2). The overall shape of the ovary was not altered but there was slight decrease in the weight of the organ of mice treated with 5mgs (FIG.3) and still more decrease in weight was observed in mice treated with 10 mgs/kg BWt, as compared with control (FIG.4). Haemorrhage in blood capillaries was prominent in mice treated with 5mgs and very prominent in mice

treated with 10mgs/kg Bwt. But most of the follicles reached the maturity. The total number of atretic follicles in mice treated with 5mgs/kg Bwt was hence found to be high as compared with control. The overall shape of the ovary was not altered nor there did any significant change in organ weight as compared with control.

Five days chronic ingestion of (10mgs/kg B.Wt.) Ethidium bromide revealed pronounced degenerative changes in ovarian histoarchitecture. The sequence of atretic changes involved nuclear degeneration, characterized by Chromatolysis, rupture and dissolution of nuclear membrane. Granulosa cells associated with degenerating follicle types (bilaminar, Multilaminar and graffian follicle) showed desquamation, cytolysis and nuclear dissolution. Extensive vacuolation also occurred in these follicles. Only few follicles reached the maturity. In most of the females Graffian follicle failed to rupture, which led to failure in ovulation. This resulted in Sterility of the females.



FIG: 1 Low power photograph showing primary PF) and secondary follicles (SF) in control Ovary.
Haematoxylin -Eosin staining (10X)

FIG: 2 High power photomicrograph showing Graffian (GF) and Atretic Follicles (AF) in a control ovary.
Haematoxiline –Eosin staining (40X)




FIG. 3: High power photomicrograph of Mice Ovary after 5mg/kg Bwt.dose of Ethidium bromide.
NOTE-Secondary follicle (SF) and minor changes in Granulosa cells(40X).Well developed follicle with fertilization cone.Slight degeneration seen in follicular cells of Graffian Follicle(GF).Atretic Follicles seen clearly.




FIG.4: High power photomicrograph of an ovary demonstrating various degenerative changes after 10mg/kg Bwt.dose of Ethidium Bromide.

NOTE- Cell necrosis,infiltration of interstitial cells and vacuolation of corpus leuteum

ACKNOWLEDGEMENT

I dedicate my work to my parents. Without their blessings my work was not possible. I acknowledge my heartiest feelings for the sacrifice of my husband Sanjay and my son Gaurang during my study.

REFERENCES

1. Stevenson P, Sones KR, Gicheru MM, Mwangi EK. (1995). "Comparison of isometamidium chloride and homidium bromide as prophylactic drugs for trypanosomiasis in cattle at Nguruman, Kenya.". *Acta Trop.* 59 (2): 257–258. PMID 7676909.
2. Diaz F, Bayona-Bafaluy MP, Rana M, Mora M, Hao H, Moraes CT. (2002). "Human mitochondrial DNA with large deletions repopulates organelles faster than full-length genomes under relaxed copy number control.". *Nucleic Acids Res.* 30 (21): 4626–33. Doi: 10.1515/CCLM.2005.141. PMC 135822. PMID 16201894.
3. Huang Q, Fu WL (2005). "Comparative analysis of the DNA staining efficiencies of different fluorescent dyes in preparative agarose gel electrophoresis". *Clin. Chem. Lab. Med.* 43 (8): 841–2. doi:10.1515/CCLM.2005.141. PMC 135822. PMID 16201894.
4. Dean Madden, Safer stains for DNA. accessed 2009-12-08. a b Singer VL, Lawlor TE, Yue S. (1999). "Comparison of SYBR Green I nucleic acid gel stain mutagenicity and ethidium bromide mutagenicity in the Salmonella/mammalian microsome reverse mutation assay (Ames test)". *Mutat Res.* 439 (1): 37–47. PMID 10029672.
5. Ohta T, Tokishita S, Yamagata H. (2001). "Ethidium bromide and SYBR Green I enhance the genotoxicity of UV-irradiation and chemical mutagens in *E. coli*". *Mutat Res.* 492 (1-2): 91–7. PMID 11377248.
6. Novel Juice testing report.
7. M.J. Waring (1965). "Complex formation between ethidium bromide and nucleic acids.". *Journal of Molecular Biology* 13 (1): 269–282. Doi: 10.1016/S0022-2836(65)80096-1. PMID 5859041.
8. J McCann and B N Ames (1975). "Detection of carcinogens as mutagens in the Salmonella/microsome test: assay of 300 chemicals". *PNAS* 72 (12): 5135–5139. doi:10.1073/pnas.72.12.5135. PMC 388891. PMID 1061098.
9. National Toxicology Program (August 15, 2005). "Ethidium Bromide: Genetic Toxicity". <http://ntp.niehs.nih.gov/index.cfm?objectid=BDAF3AE4-123F-7908-7BE09D1BEA25B435>. Retrieved September 30, 2009
10. M.J. Kramer, E. Grunberg. (1973). "Effect of Ethidium Bromide against Transplantable Tumors in Mice and Rats.". *Experimental Chemotherapy* 19 (4): 254–258. doi:10.1159/000221462.
11. von Wurmb-Schwark N, Cavelier L, Cortopassi GA. (2006). "A low dose of ethidium bromide leads to an increase of total mitochondrial DNA while higher concentrations induce the mtDNA 4997 deletion in a human neuronal cell line.". *Mutat Res.* 596 (1-2): 57–63. doi:10.1016/j.mrfmmm.2005.12.003. PMID 16488450.
12. National Toxicology Program (August 15, 2005). "Executive Summary Ethidium Bromide: Evidence for Possible Carcinogenic Activity". <http://ntp.niehs.nih.gov/?objectid=6F5F63F6-F1F6-975E-79965F7EE68AE7C0>. Retrieved September 30, 2009.



*Corresponding Author:

MANISHA MATHUR
Department of Zoology,
G.N.Khalsa College,
Matunga, Mumbai, India