



BASIC ANATOMICAL AND PHARMACOGNOSTICAL STUDY OF *CASSIA FISTULA* LINN (CAESALPINIACEAE)

B.Lavanya*, N.Narayanan, A.Maheshwaran, S.Suganya, Y.Surya sree, S.Vel Aravindan and
M.Vigneshwar

Jaya College of Paramedical Science, College of Pharmacy, Thiruninravur, Chennai,
Tamilnadu, India.

*Corresponding Author Email: blavanyavels@gmail.com

ABSTRACT

Aim: To perform the basic anatomy and pharmacognostical study of the plant *cassia fistula* Linn. **Materials and Methods:** The *cassia fistula* was collected, authenticated and performed the review of the plant, physicochemical, microscopical, Powder microscopical and Pharmacognostical evaluations. **Conclusion:** Standardisation of the plant using the basic anatomical study such as microscopical and powder microscopical study and physiochemical studies for further proceedings in future.

KEY WORDS

Cassia fistula, Pharmacognostical evaluation physical evaluation, chemical evaluation.

AIM OF WORK

Introduce the plant by using Ethanobotanical study, Microscopical study and physicochemical evaluation.

PLAN OF THE WORK:

- Introduction about plant and authentication.
- Ethanobotanical review
- Secondary and tertiary metabolites study
- Folklore study of drug for future studies
- Microscopical studies
- Powder microscopy
- Physicochemical evaluation

I. INTRODUCTION

Cassia fistula Linn. (*Cassia*) family *Caesalpiniaceae* known as *Amulthus/indian laburnum*¹. It is one of the drugs present in tropical regions of Asia. *Cassia* is

naturally distributed across India. The various parts of the tree like the bark, root, flowers, leaves, fruit pulp are used medicinally and have several health benefits. It has been extensively used in Ayurvedic system of medicine for various ailments². By studying the ethanobotanical review, secondary and tertiary metabolites, folklore literature shows that, the drug which exhibit more than 25 pharmacological uses.

II. GEOGRAPHICAL DISTRIBUTION

Cassia fistula was available in deciduous forests throughout the greater parts of India, ascending to 1300 m in outer Himalaya³. In Maharashtra, it scattered as a tree throughout the Deccan and Konkan. The plant is cultivated as an ornamental throughout India.

III. ETHANO BOTANICAL REVIEW:



- Flowers deep yellow in terminal, drooping⁴ racemose.
- 30-60 cm in length.
- Calyx is usually oblong, obtuse and pubescent.
- Corolla with 5 subequal, obovate, small clawed petals.
- Stamens present 10 in numbers, axial present 3 in numbers with erect filaments with basal fixed curved anthers⁵.
- Cassia fistula is a medium sized tropical deciduous tree, 10 m tall with a straight trunk to 5 m, 1 m diameter and spreading branches.
- Stem is grey, smooth and slender when young and dark brown and rough when old.
- Leaves are alternate, usually pinnate, 30-40 cm in length, with 4-8 sets of ovate leaflets, 7.5-15 cm in length, 2-5 cm thick, entire, and the petioles 2-6 mm in length⁵.

IV. CLASSIFICATION⁵

Kingdom:	Plantae
Subkingdom:	Tracheobinota
Super Division:	Spermatophyta
Division:	Mangoliophyta
Class:	Magnoliopsida
Sub Class:	Rosidae
Order:	Fabales
Family:	Fabaceae
Genus:	Cassia
Species:	Fistula

V.VERNACULAR NAMES⁵

Bengali	<i>Amultash, sondal, sonali</i>
English	<i>Golden shower, Indian laburnum</i>
Gujarati	<i>Girmala</i>
Hindi	<i>Bandarlathi, bharva, suvarnaka</i>
Malayalam	<i>Tengguli, rajah</i>
Sanskrit	<i>Saraphala, survanaka, argwadha, rajtaru</i>
Tamil	<i>kavani, konnai, tirukontai, sarakkonne, Raelachettu</i>
Telugu	<i>Kakkemara</i>
Marathi	<i>Bahava</i>
Punjabi	<i>Amaltaas, Kaniyaar, Girdnalee</i>
Oriya	<i>Sunaari</i>
Urdu	<i>Amaltaas</i>
Arab	<i>Khayarsambhar, chaiyaphruek, khuun</i>
Thai	<i>Canâfistula mansa, chácara, GuayabaCimarrona</i>
Spanish	<i>Bâton casse, casse doux, casse espagnol</i>
Trade name	<i>Indian laburnum</i>

VI.COLLECTION AND AUTHENTICATION

The barks of *cassia fistula* were collected from Ariyalur district, Tamil Nadu and it was Taxonomic identified and authenticated by Prof. Dr.P. Jayaraman, Ph.D, Director, Retd, Plant Anatomy Research Centre (PARC) West Tambaram, Chennai. An authentication certificate number PARC/2015/3025. The bark of *cassia*

fistula was separated and shade dried. The dried material was reduced to a coarse powder and was successively extracted in soxhlet apparatus using ethyl acetate for 24 hrs). The solvents were redistilled and ethyl acetate extract was concentrated under reduced pressure and air dried. The yield of ethyl acetate extract was found to be 36%w/w.

VII.LITERATURE REVIEW

VII. i) BRIEF DESCRIPTION OF THE SECONDARY AND TERTIARY METABOLITES PRESENT IN THE VARIOUS PARTS OF CASSIA FISTULA

Bark	• Flavonol glycosides • Oxyanthraquinone • Dihydroxyanthraquinone
Pods	• Arabinopyranoside • 5-Nonatetracontanone • 2-Hentriacontanone

Leaves	• Quinone glycosides • Sennosides A • Glucoside • Barbaloin • Aloin • Formic acid • Butyric acid • Oxalic acid • Pectin • Tannin
Flowers	• Anthraquinone glycosides⁶ • Hentriacontanoic acid • Triacontanoic acid • Nonacosanoic acid • Heptacosanoic acids • Ceryl alcohol • Kaempferol • Anthraquinone glycoside • Fistulin • Aurantiamide acetate • β-sitosterol • β-D glucoside • Triterpenes
Fruit	• Ceryl alcohol • Kaempferol • Rhein • Bianthraquinone glycoside • Fistulin • Rhein, • Sennosides A & B • Anthraquinone • Tannins • Oxyanthraquinone⁷
Pulp of the pod	• Anthraquinone glycosides • Sennosides A & B • Rhein and its glucoside, • Barbaloin • Formic acid • Oxalic acid
Pulp	• Sugar moiety • Tannic substance • Albuminous starch • Calcium oxalate • Gluten • Sucrose, 31.3% • Fructose, 26.2 • Glucose, 42.5% • High concentration of potassium • Gum • Gluten
Root	• 7-Methylphyscion • Betulinic acid

	• <i>β-sitosterol</i>
Root bark	• <i>Tannins</i> • <i>Phlobaphenes and</i> • <i>Oxyanthraquinone</i>
Seed oil	• <i>Cyclopropenoid fatty acids</i> • <i>Vernolicid</i> • <i>Malvalic acid</i> • <i>Stetculic acid</i>
Fruit pulp	• <i>Cyclopropenoid fatty acids</i> • <i>Vernolic acid</i> • <i>Malvalic acid</i> • <i>Stetculic acid</i>
Powdered stem bark	• <i>Tannins</i> • <i>Lupeol</i> • <i>Haxacosanol</i> • <i>B-sitosterol</i>
Seed	• <i>Sterculic acid</i> • <i>Malvalic acid</i> • <i>Glycerides</i> • <i>Oleic acid</i> • <i>Linoleic acid</i> • <i>Stearic acid</i> • <i>Palmitic acid</i> • <i>Cephalin</i> • <i>Lecithin</i> • <i>Phospholipids and</i> • <i>Carbohydrates (11.8%).</i>
Flower's pollen	• <i>Amino acids</i> • <i>Carbohydrate (11.75)</i>
Wood	• <i>Fistucacidin</i> <i>(3, 4, 7, 8, 4'-pentahydroxyflavan).</i>
Vegetative organs	• <i>Proanthocyanidins,</i> • <i>Flavonoids</i>
Entire plant	• <i>Tannin</i> • <i>Saccharose 53-66%;</i> • <i>Invert sugar</i> • <i>Citric acid</i> • <i>Coloring matter</i> • <i>Pectin</i> • <i>Anthraquinone</i>
Flower oil	• <i>(E)-Nerolidol (38%)</i> • <i>2-Hexadecanone (17%),</i>
Leaf oil	• <i>Phytol (16.1%).</i>

III .ii) BRIEF STUDY OF THE CASSIA FISTULA PARTS

S.NO	PART OF THE PLANT	USES
1.	Seed	• Skin diseases • Abdominal discomfort • Leprosy • Laxative • Carminative • Cooling agent • Anti-pyretic properties • Constipation
2.	Flower	• Skin diseases • GIT disorder • Leprosy • Astringent • Laxative • Febrifugal • Wound healing properties • Decoction of the flowers is given for stomach troubles
3.	Fruit	• Skin diseases • Fever • Abdominal pain leprosy
4.	Root	• Cardiac disorders • Biliousness • Rheumatic condition • Hemorrhages • Wounds boils • Ulcers
5.	Pulp	• Purgative • Disorders of liver • Biliousness • Tonic • Gout • Rheumatic condition. • Antipyretic • Remedy for malaria • Blood – poisoning • Anthrax • Diahorrea • Leprosy • Diabetes • Removal of abdominal obstructions
6.	Leaves	• laxative
7.	Ripe pod	• traditional medicines • laxative drug • Anthelmintic Activity

VII. ii) BASED ON THE LITERATURE REVIEW THE FOLLOWING CASSIA FISTULA HAS PHARMACOLOGICAL ACTIVITIES:

- Antibacterial activity
- Larvicidal activity⁸
- Antifertility activity
- Antifungal activity
- Anti-inflammatory
- Antioxidant activities
- Anti-diabetic activity⁹
- Anti-itching activity
- Anti-parasitic activity
- Antiulcer activity
- Anti-leishmaniac activity
- Antimicrobial activity
- Antiparasitic activity
- Antipyretic activity¹⁰
- Analgesic activity
- Antitumor activity
- Antitussive activity
- Clastogenic effect
- CNS activities
- Hepatoprotective activity
- Hypolipidemic activity
- Laxative activity
- Leukotriene inhibition activity
- Wound healing activity

VIII. PHARMACOGNOSTICAL EVALUATION:

The crude drug is studied pharmacognostically and its microscopical studies as follows

VIII. i) morphological study

- **Colour:** Deep brown
- **Odour:** characteristic agreeable
- **Taste:** Bitter
- **Size** : 20-40 cm

VIII. ii) MICROSCOPICAL STUDY

The microscopical studies¹¹ are performed using the standard procedure and results are given below

VIII. iii) POWDER MICROSCOPY

The powder preparation of the bark tissue shows the following inclusions.

FIBRE:

There are long, thick walled fibre with narrow tapering ends

The lumen of the fibre is narrow and the cell walls are lignified.

The fibres are up to 1.3 mm in length and 50 µm in thickness.

SCLEREIDS:

Scattered in the powder are small, squarish highly thick walled sclereids which are called brachy sclereids⁷.

The sclereids are 10 µm in length and breadth

PHLOEM PARENCHYMA CELLS:

The vertical row of rectangular parenchyma cells is seen in thick bundles.

Within the parenchyma cells are seen cuboidal calcium oxalate crystals.

The crystals are in single vertical row within each parenchyma strand.

VIII. iv) PHYSICOCHEMICAL EVALUATION¹²:

a. The physical evaluations are performed for the crude drug of *cassia fistula* linn.

Moisture content

Moisture content were determined by heating the crude drug at 105°C in an oven to a constant weight by this we can determine the active chemical constituents of the drug on air dry bases.

Ash content

Total ash value

The sample of crude drug were admixed with sand, soil, calcium oxalate, chalk powder then it is made to incinerate, then the inorganic salt content was determined.

Acid insoluble ash

A part of total ash was dissolved in dil. HCl, filtered and then weight of the residue was measured.

Extractive value

Water soluble extractive

The drug is exhausted with water and extract were obtained`

Alcohol soluble extractive

Alcohol being the ideal solvent for extraction, the drug is exhausted with alcohol and extractive value was found.

Ether soluble extractive

The drug is exhausted with non-volatile ether so that extractive value of drugs could be obtained

b. The chemical evaluations are performed for the crude drug of *cassia fistula* linn.

The chemical test for the crude drug is performed and the results are tabulated below.

IX.RESULT AND DISCUSSION

IX. i) Result of microscopical studies¹⁴

Figure 1.1: Surface view of the bark showing thick flakes




Figure 1.2: Bark cut transversely showing outer bark (periderm) and inner secondary phloem of collapsed and non-collapsed phloem zones.




(cph – collapsed phloem; ob- Outer bark ; Ncph – Non-collapsed phloem ;)

Figure 2.1: T.s of collapsed phloem zone showing thick dark tangential lines.



(cph – collapsed phloem; PhR – phloem ray; PhF – phloem fibre)


Figure 2.2: T.s of collapsed phloem showing tangential dark lines of crushed phloem elements.



(PhR – phloem ray; Dp – dilated parenchyma; Cph – collapsed phloem)

(PhR – phloem ray; PhP – phloem parenchyma; CC – Companion cell; SE – Sieve elements
PhF- phloem fibre)

Figure 3.2: Non- collapsed phloem elements enlarged




(PhR – phloem ray; PhF – phloem fibre; se- sieve elements; Cc- companion cell;)

Figure 4.1: T.L of phloem showing phloem rays




(Cr – crystal; Se- sieve elements; PhR- phloem ray)

Figure 4.2: Phloem rays and sieve elements enlarged




(pa – parenchyma; PhR – phloem ray; Se – sieve elements; Sp- sieve plate)

Figure 5.1: T.L of view of phloem showing ray seriation and ray arrangement




(pa – parenchyma; PhR – phloem ray)

Figure 5.2: Phloem, uniseriate and biseriate rays.




(Ur – Uniseriate; Br – biseriate ray;)

Figure 6.1: Phloem showing biseriate rays, multiseriate rays and uniseriate rays.




(Br- biseriate ray; Pc – procumbent; Uc- Uniseriate ray; Mr- multiseriate rays)

Figure 7.1: R.L.S of phloem showing horizontal oriented phloem rays.



(Pa- parenchyma; PhR – phloem ray;)

Figure 8.1: Bark powder showing wide fibre and branch sclereids.



(Bsc- Brachy sclereids; fi- Fibres;)


Figure 8.2: Long narrow fibre



(Fi- Fibre;)

IX. ii) Powder microscopy of crude drug¹³:

S.NO	NAME OF THE PARTS	DESCRIPTION	IMAGE
1	XYLEM	Xylem cells are long and narrow like pipes ROLE: They carry water	
2	PHLOEM	The main components of phloem are Sieve element Companion cells ROLE: They conduct sugar and amino acid.	
3	VESSEL	They are stacked one on top of another in long column. ROLE: Water conducting tissue of plants.	

4	CORTEX	Cortex is the outer most layer thin walled parenchyma cells in the ground tissues. ROLE: Food storage. Mechanical support.	
5	PERIDERM	Periderm is the soft growing tissue between bark and wood. ROLE: Outer corky protective of the woody stem and roots.	

IX.iii) Results of physical evaluation:

S.No	Evaluation test	Values obtained in % w/w
1.	Moisture content	0.973
2.	Ash content	
	Total ash	9.03
	Acid insoluble ash	1.6
	Water insoluble ash	4.70
3.	Extractive value	
	Water soluble extractive	4.66
	Alcohol soluble extractive	10.039
	Ether soluble extractive	101.55mcg/ml

CONCLUSION

- *Cassia fistula* linn belonging to family caesalpiniaceae, popularly known as Indian laburnum is highly valued medicinal plant.
- It is known as a rich source of tannins, triterpenes, flavanoids²⁰ and glycosides present in *cassia fistula*, might be medically important and/or nationally valuable.
- The plant undergone the standardisation procedure such as
 - I. Authentication
 - II. Physico chemical studies
 - III. Pharmacognostical studies.
- A further study has to be performed for the characterisation, Isolation, Spectral studies of compounds. This gives the ideas about wide range of compounds in *cassia fistula*. Hence *cassia fistula* usage in medicinal field to treat various ailments.

REFERENCES

1. Sushma Kainsa, Praveen Kumar, Poonam Rani. Pharmacological Potentials of *Cassia auriculata* and *Cassia Fistula* Plants: Review. Pakistan Journals of Biological Science 2012; 15(9): 408-417
2. Gamble. J.S, et al. Flora of the Presidency of Madras – Vol I, II, & III. Botanical Survey of India, Calcutta, India.
3. Henry. A.N, et al. Flora of TamilNadu, India- Vol.3. Botanical Survey of India, Southern Circle, Coimbatore, India. pp-258:1987.
4. Mathew K.M, et al. The Flora of Tamil Nadu Karnatic. Vol. Polypetalae,. Vol.3. Gamopetalae & Monochlamydae. The Ranipat Herbarium, St. John's College, Tiruchirappalli, India. Pp. 688, pp689-1540.
5. Metcalfe C.R, et al. Anatomy of the Dicotyledons- Vol. I & II. Clarendon Press, Oxford. 1950.
6. Aurapa sakulpanich, et al. Determination of anthraquinone glycoside content in *cassia fistula* leaf extract for alternative source of laxative drug; international journal of biomedical and pharmaceutical sciences, 3(1); 42-45.

7. Aurapa sakulpanich, et al. extraction method for high content of anthraquinone from cassia fistula pods. *J health res*, 2008; 22(4):167-172.
8. Markout M, Bekkouche K, Larhsini M. Bousaid M. Lazrek HB, Jana M, Evaluation of some Moroccan medicinal plant extracts for larvicidal activity. *J. Ethanopharmacol*, 2000; 73:293-297.
9. Alam. MM, et al. Treatment of diabetes through herbal drugs in rural India.
10. Patel D, et al. Antipyretic and analgesic Activities of *Aconitum spicatum* and *Cassia fistula*. *Pharm. Biol.* 1965; 157: 2227.
11. Satyavati GV, et al. *Medicinal plant in India: ICMR: New Delhi, India, 1989.*
12. Kokate.C.K. *Evaluation of crude drug. (ed). Practical Pharmacognosy, 5th ed.: M.K.Jain; 2014. pp. 125-127.*
13. O. Brien, et al. Polychromatic Staining of Plant Cell Walls by toluidine blue- O, *Protoplasma*:59:364-373:1964.
14. Kokate. C.K. *Analytical Pharmacognosy. (ed). Pharmacognosy, 33rd ed.: Nirali Prakashan; Nov: 2005. pp. 97-132.*
15. http://www.deserttropicals.com/plant/Fabaceae/cassia_fistula_a.html

***Corresponding Author:**

B Lavanya*

Email: blavanyavels@gmail.com