International Journal of Pharmacy and Biological Sciences
ISSN: 2321-3272 (Print), ISSN: 2230-7605 (Online)

IJPBS | Volume 7 | Issue 4 | OCT-DEC| 2017 | 106-115

Research Article | Biological Sciences | Open Access | UGC Approved | MCI Approved Journal

ETHNOBOTANICAL PLANTS USED IN VARIOUS FORMS TO CURE DIABETES- A BRIEF STUDY IN BHOWRAGURI VILLAGE OF KOKRAJHAR DISTRICT, BTC, ASSAM, INDIA

Tamal Das^{1*} and Tribeni Mandal²

^{1*}Student of BSc Semester V, Department of Zoology, Science College, Kokrajhar, BTC, Assam- 783336, India. ²Asst.Professor, Department of English, Science College, Kokrajhar, BTC, Assam-783336, India.

*Corresponding Author Email: creamice29@gmail.com

ABSTRACT

Aim: Diabetes mellitus (DM) commonly referred to as diabetes, is a group of metabolic disorder in which there is high blood sugar level for a prolonged period. The main objective of the study is to present the medicinal plants used in Bhowraguri village of kokrajhar district for anti-diabetic purpose. Methods: A thorough study was carried out in Bhowraguri, which is a small village in kokrajhar district. The collected plant samples were identified with the help of taxonomic literature. The voucer specimens were deposited in the same herbarium for record. The photos were collected from our own houses, some neighbours etc. Result: A total of 39 medicinal plant species belonging to 31 families were found to be used by the local people of the area surveyed under study for treatment of diabetes. The plant parts used for the treatment of diabetes include leaves, fruits, flowers, barks, stems, roots, seeds as well as whole plants. Conclusion: For the treatment of diabetes various anti-diabetic drugs are used which are not only expensive but also produce side effect. Therefore, there has been a growing interest in the ethnobotanical approach to examine the anti-diabetic properties of plant traditionally used by the ethnic groups in different parts of the world. During our study we recorded 39 species of antidiabetic medicinal plant from Bhowraguri, a small village under kokrajhar district. Hence our record on anti-diabetic medicinal plants from this region is very important for further studies.

KEY WORDS

Anti-diabetic medicinal plant, anti-diabetic drugs, diabetes mellitus, Ethnobotany.

Introduction:

Diabetes mellitus (DM) commonly referred to as diabetes, is a group of metabolic disorders in which there are high blood sugar levels over a prolonged period. Symptoms of high blood sugar include frequent urination, increased thirst and increased hunger.

Diabetes is due to either the pancreas not producing enough insulin or the cells of the body not responding properly to the insulin produced. There are three main types of diabetes mellitus which are-Type 1, Type 2 and Gestational diabetes.

According to WHO report globally, approximately 347 million people or 5-8% of the global population is estimated to be affected by this disease. Diabetes now

is becoming the third "killer" of mankind along with cancer, cardiovascular and cerebrovascular disease. IT has also been predicted that by the year 2025 more than 75% of people with diabetes will reside in developing countries, as compared with 62% in 1995.A good number of synthetic commercial anti-diabetic drugs/agent like sulfonylureas, biguanides, glucosidase inhibitors and thiazolindiones are well known today which are not only expensive but also produces side effect. Therefore, there has been a growing interest in the ethnobotanical approach to examine the anti-diabetic properties of plant traditionally used by the ethnic groups in different parts of the world. A large no. of plants and plant-parts has been investigated for their



beneficial role and anti-diabetic properties. In view of its medicinal importance the present study has been focused to know the traditional medicinal plant wealth that is being used by the tribal people of BTC area against diabetes (Swargiary et al., 2013).

Materials and Methods:

Geographical position of study area:

Kokrajhar district is an administrative district in the state of Assam in North-Eastern India. The district headquarter is located in Kokrajhar town. It is located at a latitude of 24° 24′ 0″ N and a longitude of 90° 16′ 0″ E. It covers an area of 3,169.22 sq.km. The district is hot in summer. In Kokrajhar the highest day temperature is in between 28 °c to 40 °c. The average temperature in Jan is 20 °c, Feb is 22 °c, March is 28 °c April is 29 °c, May is 28 °c. Bhowraguri is a small village in kokrajhar district and it is 26 kms away from kokrajhar town.

A thorough study was carried out in Bhowraguri which is a small village in kokrajhar district. During our survey the information was gathered from the local elderly people (kobiraj,ojha) and some diabetic patients from the survey areas. The collected plant samples were identified with the help of taxonomic literature. The voucer specimens were deposited in the same herbarium for record. The photos were collected from our own houses, some neighbours etc.

Result and Discussion:

A total of 39 medicinal plant species belonging to 31 families were found to be used by the local people of the area surveyed under study for treatment of diabetes (table 1). It was observed that the plant parts used for the treatment of diabetes include leaves, fruits, flower, barks, stems, roots, seeds as well as whole plants. The number of species along with their families are given below-

Rutaceae - 2 (species), Asteraceae- 2 (species), Celastraceae- 1 (species) , Meliaceae -1 (species), Apiaceae- 1 (species), Fabaceae -2 (species), Annonaceae- 1 (species), Phyllanthaceae- 1 (species), Crassulaceae- 1 (species), Zingiberaceae- 1 (species), Moraceae- 3 (species), Poaceae- 1 (species), Myrtaceae-1 (species), Lamiaceae- 4 (species), Cucurbitaceae- 1 (species), Anacardiaceae- 1 (species), Apocynaceae- 1 (species), Menispermaceae - 1 (species), Arecaceae- 1 (species), Oleaceae -1 (species), Caricaceae- 1 (species), Cannabaceae- 1 (species), Araceae- 1 (species), Lauraceae- 1 (species), Moringaceae- 1 (species), Malvaceae- 1 (species), Asphodeloideae- 1 (species), Oxalidaceae- 1 (species), Musaceae -1 (species), Combretaceae- 1 (species) and Acanthaceae - 1 (species).



Table -1: The scientific names, families, local names, parts used, and preparations done in order to collect the anti-diabetic medicinal plants are listed below-

| Sl.No | Scientific name & Family | Local name | Part used | Preparation to use |
|-------|--------------------------|---------------|--------------|---|
| 1 | Murraya koenigii (L.) | Narahingha | Leaves | The leaf of this plant is consumed raw, |
| | sprengel. (Rutaceae) | | | early in the morning, in empty stomach. |
| 2 | Chromolaena odorata (L.) | Motmoti | Leaves | The leaf of this plant is pasted, and the |
| | R.M.King & H.Rob. | | | juice is consumed raw, early in the |
| | (Asteraceae) | | | morning, in empty stomach. |
| 3 | Artocarpus heterophyllus | Kothal | Leaves | The leaf of this tree is deepened into a |
| | Lam. | | | glass of water overnight and the water is |
| | (Moraceae) | | | drunk in the next morning. |
| 4 | Tinospora cordifolia | Gulonccholota | Stem | The stem of this plant is soaked in water |
| | (Thunb.) Miers | | | overnight and the water is drunk the next |
| | (Menispermaceae) | | | morning. |
| 5 | Catharanthus roseus (L.) | Noyontora | Leaves | Fresh leaf extracts or fresh leaves may be |
| | G. Don,1837 | | | chewed in empty stomach (Swargiary et |
| | (Apocynaceae) | | | al.,2013) |
| 6 | Mikania micrantha Kunth | Kemlota | Leaves | The leaf of this plant is chewed raw. |
| | (Asteraceae) | | | |
| 7 | Salacia reticulata Wight | Xeora | Leaves | The leaf of this plant is pasted, and the |
| | (Celastraceae) | | | juice is taken. |
| 8 | Mangifera indica L. | Aam | Leaves | The new leaves of this tree are chewed |
| | (Anacardiaceae) | | | raw. |
| 9 | Ficus benghalensis L. | Bot | Leaves | Bark infusion or decoction is used (Mondal |
| | 1753 | | | et al.,2013) |
| | (Moraceae) | | | |
| 10 | Cucumis sativus | Telakucch | Leaves | The leaf of this plant is pasted, and the |
| | Manicgardener, March | | | juice is taken. |
| | 17,2011 | | | |
| | (Cucurbitaceae) | | | |
| 11 | Clerodendrum | Bhetita | Leaves | Pasted leaves is taken orally (Mondal et |
| | infortunatum L. | | | al,2013) |
| | (Lamiaceae) | | | |
| 12 | Ocimum tenuiflorum L. | Tuloxi | Leaves | Powdered leaves are taken with honey to |
| | (Lamiaceae) | | | treat diabetes (Mondal et al,2013) |
| 13 | Aegle marmelos (L.) | Bel | Leaves | Leaves are boiled, and the juice is taken |
| | Correa | | | (Mondal et al.,2013) |
| | (Rutaceae) | | | |
| | | | | |
| 14 | Azadirachta indica A. | Neem | Leaves | Raw leaf extracts mixed with little water is |
| | Juss., 1830 | | | taken at a dose of 2-3 tea spoon daily in |
| | (Meliaceae) | | | empty stomach (Swargiary et al, 2013) |
| 15 | Centella asiatica (L). | Manimuni | Whole plant | The leaf of this plant is taken raw or pasted |
| | Urban | | | and taken as chutney. The root is also |
| | (Apiaceae) | | | consumed raw and even cooked. |
| 16 | Erythrina variegata L. | Mandaar | Root | Fresh roots are grounded for obtaining |
| | (Fabaceae) | | | juice. 25 ml of juice is taken for 1 week |
| | | | | without water. |
| 17 | Annona squamosa L. | Aataphol | Bark, leaves | Raw bark or/and leaves are grinded, and |
| | (Annonaceae) | | | the extracts are obtained by squeezing. |
| | | | | The extract is then filtered and used 2 to 3 |
| | | | | spoons full every morning (Swargiary et al, |
| | | | | 2013) |



| 18 | Phyllanthus emblica L. | Amlokhi | Fruit | Fruit juice or raw fruit is taken orally |
|----|---|---------------|------------------------------|--|
| 19 | (Phyllanthaceae) Syzygium cumini (L.) Skeels. (Myrtaceae) | Kolajam | Seed | (Mondal et al,2013) Powdered seeds about 1 teaspoonfull is taken with water in the morning in empty stomach and also in the evening before meals (Swargiary et al,2013) |
| 20 | Kalanchoe pinnata (Lam.) pers. (Crassulaceae) | Pategoja | Leaves | 1g of raw leaves grinded with 100 ml of water is consumed. Leaf extracts ,2-3 tea spoons early in the morning is also taken (Swargiary et al,2013) |
| 21 | <i>Leucas aspera</i> (Lamiaceae) | Dron | Whole plant | It is used as potherb during diabetic treatment (Swargiary et al,2013) |
| 22 | Curcuma Longa L. (Zingiberaceae) | Halodhi | Root | About 8g of raw turmeric is grinded, mixed with water and ½ spoon of honey and taken for 1 month after meal (Swargiary et al,2013) |
| 23 | Cynodon dactylon (L.) pers. (Poaceae) | Duburi | leaves | This leaf is taken raw or sometimes it may be dried up, preserved in a container and taken with water, food or tea. |
| 24 | Ficus religiosa Linn. (Moraceae) | Aahot | Leaves, fruit | Leaf or fruit is consumed orally (Mondal et al.,2013) |
| 25 | Areca catechu L. (Arecaceae) | Tamul | Nut | Dried and soaked with water (Mondal et al.,2013) |
| 26 | Nyctanthes arbor- tristis L. (Oleaceae) | Hewali | Leaves, flowers | Young leaf juice, flower paste is taken orally. (Mondal et al.,2013) |
| 27 | Carica papaya L. (Caricaceae) | Amita | Seed | Powdered seed is mixed with water (Mondal et al.,2013) |
| 28 | Cannabis sativa L. (Cannabaceae) | Bhaang | Leaves, stems, flowers | Leaves and stems are soaked in water. (Mondal et al.,2013) |
| 29 | Alocasia macrorrhizos (L.) G.Don. (Araceae) | Maankochu | Rhizomes | Dried rhizome is used (Mondal et al.,2013) |
| 30 | Gmelina arborea Roxb. (Lamiaceae) | Gomari | Leaves, fruits | Leaf juice is used (Mondal et al.,2013) |
| 31 | Cinnamomum tamala (BuchHam.) T.Nees & C.H.Eberm. (Lauraceae) | Tezpat | Bark, root | Powdered bark is consumed (Mondal et al.,2013) |
| 32 | Moringa oleifera Lam. (Moringaceae) | Sajna | Leaves, fruits | The leaf of this plant is pasted, mixed with water and consumed early in the morning, in empty stomach. Drumsticks are taken as vegetables after cooking. |
| 33 | Abelmoschus esculentus (L.) Moench (Malvaceae) | Bhendi | Vegetable | The vegetable is taken raw and cooked. The vegetable is soaked overnight in water and the same is consumed the next morning. |
| 34 | Aloe vera (L.) Burm.f. (Asphodeloideae) | Ghrita kumari | Leaves | The juice of the plant is taken. |



| 35 | Averrhoa carambola L. (Oxalidaceae) | Kordoitenga | Fruit | The fruit is taken raw and cooked both. |
|----|--|-------------|------------------------|--|
| 36 | <i>Musa acuminata</i> Colla (Musaceae) | Kol | Stem, fruit, flower | The stem, fruit, and flower is cooked and taken. |
| 37 | Cajanus cajan (L.) Millsp. (Fabaceae) | Rohorpaat | Leaves | Leaves are cooked and taken. |
| 38 | <i>Hygrophila auriculata</i> Schumach. (Acanthaceae) | Kulekhara | Leaves | Leaves are taken cooked and raw. |
| 39 | Terminalia arjuna (Roxb.) Wight & Arn. (Combretaceae) | Arjun | Bark | The bark is dipped in water and the water is consumed. |



(1) Murraya koenigii



(2) Chromolaena odorata



(3) Artocarpus heterophyllus



(4) Tinospora cordifolia




(5) Catharanthus roseus



(6) Mikania micrantha





(13) Aegle marmelos

(14) Azadirachta indica





(22) Curcuma Longa

(21) Leucas aspera





ne); ISSN: 2321-3272 (Print) Int J Pharm Biol Sci.

(23) Cynodon dactylon

(24) Ficus religiosa





(25) Areca catechu

(26) Nyctanthes arbor-tristis





(27) Carica papaya

(28) Cannabis sativa





(29) Alocasia macrorrhizos

(30) Gmelina arborea





(31) Cinnamomum tamala

(32) Moringa oleifera





(33) Abelmoschus esculentus

(34) Aloe vera





(35) Averrhoa carambola

(36) Musa acuminate





(37) Cajanus cajan

(38) Hygrophila auriculate



(39) Terminalia arjuna



Conclusion:

Mondal et al., 2013 has reported 141 species of antidiabetic medicinal plant from the North-Eastern part of India and during the study we recorded 39 species of antidiabetic medicinal plant from Bhowraguri which a small village under kokrajhar district is. Bhowraguri is a small village which has a large diversity of antidiabetic medicinal plants. Previously no one has recorded any information on antidiabetic medicinal plants from this small area. Hence the record on antidiabetic medicinal plants will hopefully play an important role and will be very helpful for further studies in the field.

References:

- Aadhan K and Anand sp, Survey medicinal plants used for the treatment of diabetes by the Paliyar's Tribe in Sadhuragiri hills, Tamil Nadu, India, International Journal of Herbal Medicine, 2017; 5(3): 17-25.
- Swargiary 1* Ananta, Boro 2 Hankhray, Brahma 2
 Birendra Kumar, Rahman 1 Seydur, Ethno-Botanical
 study of Anti-Diabrtic Medicinal Plants used by the Local
 people of Kokrajhar District of Bodoland Territorial
 council, India, Journal of Medicinal Plant Studies, Year:
 2013, Volume: 1, Issue: 5, First page: (51) Last page: (58)
- Sarmah Pratul Chandra, Ethno Antidiabetic Plant of Assam, International Journal of Applied And Pharmaceutical Technology, Volume-2, Issue-4,246-251, (2011).

- Mondal* Prodyut, Bhuyan Niroj, Das Sonjit, Kumar Mritunjay, Borah Sudarshana, Mahato Kabita, Herbal Mecicines useful for the treatment of Diabetes in North-East India, International Journal Of Pharmacy And biological Science, Volume-3, Issue- 1,575-589, (2013).
- Ryakala 1* Venkat Kishore, Ali 1,2* Shahin Sharif, Sharanabasava1 Hallihosur, Hasin 3 Naushaba, Sharma4 Pragya, Bora1* Utpal, Ethnobotany of Plants Used to Cure Diabetes by the People of North East India, Medicinal and Aromatic Plant Science and Biotechnology, 4(Special Issue 1),64-68, (2009).
- Singab* Abdel Nasser, Youssef Fadias , Ashour Mohamed L, Medicinal Plants With Potential Antidiabetic Activity and their Assessment, Medicinal & Aromatic Plants, Volume-3,Issue-1,1-12,2014.
- http://www.newstrackindia,com/information/worldinf o/latitudelongitude/CountryCities/India/city-Kokrajhar-1035592.htm
- 3. https://en.m.wikipedia.org/wiki/Diabetes_mellitus
- 9. Handa S.S., and Kapoor V.K., Pharmacognosy. Vallabah Prakashan Publications, New Delhi; pp. 51 (1992).
- Yuan Z., He P., Ciu J., and Takeuchi H., Hypoglycaemic effect of water soluble polysaccharide from Auricularis auricular-Judae Quel. on genetically diabetic KK-AY mice. Biosci. Biotech. and Biochem, 62: 1898-1903, (1998).
- Price K.R., Johnson L.I., and Ferewick H., The chemical and biological significant of saponins in food and feeding staffs. CRC Critical Rev. Food. Sci. Nutr, 26: 127-135, (1987).
- American Diabetes Association (ADA). Eating healthy with the diabetes food pyramid as your guide, (1999).

*Corresponding Author: Tamal Das

Email: creamice29@gmail.com