



## ENZYME ASSISTED BIOACTIVE EXTRACTION FROM *FLACOURTIA MONTANA* AND INVESTIGATION OF ITS *IN-VITRO* ANTIOXIDANT AND ANTI-DIABETIC ACTIVITY

Kavya MV<sup>1</sup>, Debika Chakrabarty<sup>2</sup>, Priyanka Prabhakar<sup>3</sup>, Kirana Shubhasri R<sup>4</sup>, Vishwaprakash Mahadimane<sup>5</sup>, Shobha G<sup>6\*</sup>

<sup>1,2,3,4,6</sup>Department of Biotechnology, Sapthagiri college of engineering (Affiliated to VTU), Bengaluru - 560057, India.

<sup>5</sup>Department of Bioscience, University of Mysore, Hemangotri, Hassan -573220, India.

\*Corresponding Author Email: [shobhag@sapthagiri.edu.in](mailto:shobhag@sapthagiri.edu.in)

### ABSTRACT

*Flacourtia* species are known for medicinal properties since ancient times. Here in we report the efficacy of the enzyme assisted extraction of bioactive compounds from *Flacourtia montana* leaf using three different enzymes and in combination of same enzymes. The extraction was carried out by enzyme formulations which contained cellulase, pectinase and amylase in water bath at a temperature of 50°C for 3 hours. Further the extract were used to determine the phenolic content, flavonoid content, antioxidant and anti-diabetic properties. The combination of enzyme used for extraction showed maximum total phenolic and total flavonoid content. The extract also showed strongest antioxidant activities and antidiabetic activity compared to other methods. The TPC ranged from 54.22± 1.25 to 31.25 ± 1.02 mg GAE/g of DW, TFC ranged from 21.77 ± 0.54 to 8.72 ± 0.2 mg QE/g of DW and TAC was found to be 149.83± 4.4 to 88.16 ± 6.0 mg EAA/g of DW. The IC<sub>50</sub> values for anti-diabetic properties varied from 300 ± 0.01 µg/ml to 910 ± 0.02 µg/ml.

### KEY WORDS

Cellulase, Amylase, Antioxidant activity, Antidiabetic activity, Enzyme extraction, *Flacourtia montana*, Pectinase.

### INTRODUCTION

Plant based medicines are used for combating diseases since ancient times due to the presence of a large number of bioactive compounds [1], hence there is a continuous search for medicinal plants that are of rich in these compounds. It has been reported that among more than 25,000 secondary metabolites that have been identified in plants [2], phenolic compounds found to distributed in all parts of higher plants shown to exhibits high degree of free radical scavenging property which may be the prime reason behind antioxidant activity, anti-tumor, antibacterial, anti-aging, anti-allergic, anti-inflammatory and antidiabetic properties [3,4]. The increased demand for the antioxidants and

antidiabetic activities from natural compounds have encouraged the research studies about enhanced extraction process. The conventional techniques of plant materials extraction are usually based on the choice of solvents and the use of heat to increase the solubility of the desired compounds. Usually, conventional techniques require longer extraction time, thus running a risk of thermal degradation of some of the bioactive compounds [5]. The solvents used in the extraction also increase the risk of environmental pollution. In last few years many new alternative methods have been developed for the extraction of phytochemicals from plants such as ultrasound-assisted extraction(UAE), enzyme assisted extraction(EAE),

microwave-assisted extraction (MAE), supercritical CO<sub>2</sub>(SC-CO<sub>2</sub>), pressurized fluid extraction (PFE). EAE has gained importance since it offers an eco-friendly approach, an improvement in quantity of target compound and enhancement in bioactivity. The extraction by EAE method is based on the ability of enzymes to degrade cell wall components, leaving the intracellular materials more exposed for extraction into solvents [4]. Many studies have been extensively conducted in recent years to identify and quantify phenolic compounds and the relevant antioxidant activity of plants [6], *Flacourtia montana* belonging to family *Flacourtiaceae* is medicinal tree of Indian origin. The species of the family are well known for their pharmacological activities and has been used as a traditional medicine in many countries for centuries [7]. From the findings of above the objective of our study was set to evaluate the efficiency of extraction of bioactive compounds by EAE in comparison to the traditional method. Later the extract were analyzed for invitro antioxidant and invitro antidiabetic studies.

## MATERIALS AND METHODS

### A. Processing of plant material

Barks, leaves, flowers, and nuts from *Flacourtia montana* tree were collected from Hassan district, Karnataka. The plant was authenticated by Prof. Dr. Sharanappa P (Department of Bioscience, Hemagangotri University, Hassan, Karnataka) and voucher specimen (PS27551016) was deposited at the Department. The fresh leaves of *Flacourtia montana* was separated from the stem and washed with running tap water to remove the surface impurities. The leaves were shade dried for a week and ground to fine powder and stored in refrigerator.

### B. Preparation of Extract

The enzyme assisted extraction was carried as described by [8] with slight modification, using commercial enzymes amylase, pectinase and cellulase. The plant extract and distilled water was taken at the ratio of 1:10 (W/V), to it enzyme at the concentration of 0.5% was added and then placed in water bath for 3 hours at 50°C. After the period of incubation, the temperature was raised to 100°C for 20 min to denature the enzymes. The extract was strained through double layered muslin cloth followed by centrifugation at 10,000 rpm for 10 min at 20°C to remove the undigested residue. The resultant supernatant was filtered using whattman filter

paper and concentrated in the oven at 55°C until dryness. The same procedure was repeated for all the enzymes and its absence. Each extract were re-suspended in water to make a 1mg/ml and 1 ml of aliquot of leaf extract were used for all the assays.

### C. Determination of Total Phenolic Content

The total phenolic content of the extracts was determined by using Folin–Ciocalteu's method describe by [9] with slight modification. Briefly, 1 ml of extract was mixed with 5 ml of Folin–Ciocalteu's reagent (diluted tenfold with distilled water) and incubated in room temperature for 2 minute. After the period of incubation, 4 ml of 7.5 % sodium carbonate solution was added and mixed thoroughly and incubated for 120 min in dark at room temperature. The absorbance of sample was measured at 765nm. The appearance of blue colour indicates the reduction of products. Blank was prepared without extract. The total phenolic content in the extract was calculated as Gallic acid equivalent (GAE) per gram of dry weight (DW) established from the calibration curve in mg/ml.

### D. Total flavonoid content

The Total flavonoid content of the unrefined leaf extract was assessed utilizing the aluminum chloride colorimetric method as describe by [10] with a slight alteration. Briefly, a volume of 1 ml of the extract was blended with 2.8 ml of distilled water and after that 0.1 ml of 1 mg/ml potassium acetic acid was added. To this, 0.1 ml of 10% aluminum chloride was added, the blend was allowed to stand for 30 min and absorbance was measured at 415nm. The appearances of yellow colour indicate the presences of flavonoids. The TFC was expressed as mg of quercetin equivalent (QE) per gram of a dry weight.

### E. Total antioxidant captivity (TAC)

The TAC of the extracts were evaluated by the phosphomolybdenum method according to the procedure described previously [6] with some modification. Briefly 1 ml extracts were mixed with 3.0 ml of complex reagent solution (including 0.6 M sulfuric acid, 28 mM sodium phosphate and 4 mM ammonium molybdate, 1:1:1). The reaction solution were incubated at 95°C for 90 min and then cooled to room temperature. Finally, the absorbance of the solution was measured at 695 nm against a blank on a UV-visible spectrophotometer. A typical blank contained 1 ml of the reagent solution along with an appropriate volume of the solvent and incubated under similar conditions.

Antioxidant capacity of ascorbic acid has been used as a reference standard from which plant extracts with potential antioxidant activity are compared. The total antioxidant capacity (TAC) was expressed as mg equivalents of ascorbic acid per gram (EAA/g).

#### F. Antidiabetic activity

##### Inhibition of Amylase Enzyme

Antidiabetic activity was examined by amylase inhibition using dinitro salicylic acid (DNS) method. Briefly, a volume of 500  $\mu$ l of plant extract having concentration of 200-1000  $\mu$ g/ml were taken in 500  $\mu$ l of sodium phosphate buffer containing  $\alpha$ -amylase (0.5 mg/ml) and incubated at 37  $^{\circ}$ C for 10 min. After a time of incubation, 500  $\mu$ l of 1% starch solution in 0.02 M sodium phosphate buffer (pH 6.9) was added. This reaction blend was then incubated for 10 minutes at 25 $^{\circ}$ C, followed by addition of 1.0 ml of DNS. The reaction was ceased by incubating in boiling water for 10 min than diluted with 10 ml distilled water, and the absorbance was measured at 540 nm. The mixture without plant extract was used as a control [11].

The  $\alpha$ -amylase inhibitory activity was expressed as percentage inhibition (%) by using the formula

$$I\% = \frac{(Ac - As)}{Ac} \times 100$$

Where Ac is the absorbance of the control and as is the absorbance of the sample. The IC<sub>50</sub> value was defined as the concentration of the sample extract to inhibit 50% of  $\alpha$ -amylase activity under assay condition.

##### Statistical Analysis

All the assays of TPC, TFC, and TAC were carried out in triplicates and results were subjected to statistical analysis. All data were expressed as mean  $\pm$  standard error (SE) and were subjected to variance analysis,  $p < 0.05$  was considered statistically significant. For antidiabetic assays, IC<sub>50</sub> value was used to determine.

## RESULTS


### Comparison of EAE with aqueous Extraction Method on TPC and TFC

In present study the efficiency of extraction with and without enzyme on TPC and TFC were compared and values are shown in figure. The TPC and TFC values of


the extracts were validated by a linear correlation between the concentration and absorbance. Based on the calibration curve an equation ( $y = 0.003x - 0.0043$ ,  $R^2 = 0.9957$ ) and ( $y = 0.0091x + 0.0429$ ,  $R^2 = 0.9752$ ) were obtained respectively, where Y represents the absorbance and X represents equivalent (mg/g). The TPC and TFC values varied from  $31.25 \pm 1.02$  to  $54.22 \pm 1.25$  (mg GAE/g DW) and  $21.77 \pm 0.54$  to  $8.72 \pm 0.2$  (mg QE/g DW). According to the statistical analysis ( $P \leq 0.05$ ) the TPC and TFC were markedly higher in an enzyme extract than the aqueous extraction. As shown in (Fig. 1 and Fig. 2), the EAE method had the highest extraction yield of TPC and TFC compared to aqueous extraction. Among the EAE extraction mixture of enzymes showed maximum phenolic content  $54.22 \pm 1.25$  mg GAE/g DW and flavonoid content  $21.77 \pm 0.54$  mg QE/g DW followed by cellulase and pectinase. These results indicated the influence of the extraction by enzyme on the total content of phenolic compounds extracted. The order of yield of TPC and TFC extracted with different methods was similar to the results of former studies (12).

### Evaluation of Antioxidant Capacity


The TAC was chosen to determine the antioxidant capacity of extracts obtained from different extraction method. It is based on the reduction of Mo (VI) to Mo (V) by the extract and subsequent formation of green phosphate/Mo (V) complex at acid pH. TAC of the phosphomolybdenum model evaluates both water-soluble and fat-soluble antioxidant capacity (total antioxidant capacity). The results are shown in Fig. 3, all the extract obtained from extraction has showed the antioxidant activity in different range. Among four types of enzyme extraction the combination of enzyme as showed highest antioxidant activities  $149.83 \pm 4.4$  EAA mg/g and lowest activity was shown by amylase. Conversely, extraction without enzyme yielded the lowest antioxidant activities of  $88.16 \pm 6.0$  EAA mg/g. This results were supported by many studies in literature. Feng [4] reported the cellulase extraction of *Trapa quadrispinosa* as showed the TAC of  $45.2 \pm 2.2$  mg GAE/gDW at concentration 2% cellulase at 50 $^{\circ}$  C.




**Figure 1. Total Phenolic content in various leaf extracts of *F. montana***



**Figure 2. Total flavonoid content in various leaf extracts of *F. montana***



**Figure 3. Total antioxidant content in various leaf extracts of *F. montana***



**Figure 4. Invitro Antidiabetic Activity of various leaf extracts of *F. montana***

#### Antidiabetic activity

In the present research, the study has been carried out to screen the effect of EAE and aqueous extracts of *F. montana* plants in inhibiting  $\alpha$ -amylase. The results of the inhibitory assay and with IC<sub>50</sub> values are summarized in Fig. 4. All the above extraction showed varying inhibitory effect on enzyme which varied from the range  $300 \pm 0.01$   $\mu$ g/ml to  $910 \pm 0.02$   $\mu$ g/ml. This may be due to the presence of higher secondary metabolites. These observations produce a scientific

basis for the use of this medicinal plant in traditional medicine for the treatment of diabetes. The dose dependent invitro antidiabetic activity of plant extract from *Nigella sativa*, *Eugenia jambolana*, *Andrographis paniculata* and *Gymnema sylvestre* [13], *C. swietenia* [14], *Ceropegia juncea*[15] was reported in literature.

#### DISCUSSION

*Flacourtia sps* has reported for valuable pharmaceutical properties, as it's contain abundant phenols, and exhibit

strong antioxidant activity [16,17]. These property is due to the phytochemicals, especially plant phenolics and flavonoids [18]. From the previous study the presence of bioactive compounds like alkaloids, Flavonoids, Tannins in aqueous extract of *F. montana* has been reported [7]. The bioactive compounds, such as terpenoids and flavonoids are the principal antioxidants that apply a scavenging effect on active oxygen free radicals by various mechanism including the inhibition of enzymes responsible for free radical generation, the chelation of metal ions, such as iron and copper and inhibit lipid oxidation at an early stage. Hence the extraction of secondary metabolites from the plant is a primary step to enhance the release of phenolic compounds, keep their stability and antioxidant activities [19]. Thus, the extraction parameters such as time, temperature, type of solvent, particle size of extract, type of enzyme, and concentration of the enzyme used influence the extraction efficiency. It has been reported that a long extraction time presented a favourable effect on the production of phenolic compounds. However, excessive extending of extraction time may induce the difference in phenolic compounds molecule structure and bioactivities due to oxidization or hydrolysis [4]. Recent advances in understanding the activity of intestinal enzymes helped in the development of newer pharmacological agents [20].  $\alpha$ -amylase enzymes present in the intestinal human play main roles in the digestion of carbohydrates to degrade starch and oligosaccharides. The inhibition action of enzyme would slows the breakdown and absorption of glucose from the GI tract and leads to reduction of postprandial glucose level elevation. This delay digestion and breakdown may have beneficial effects on insulin resistance and glycemic index control in people with diabetes [21]. It has been also reported phenolic compounds are known to reduce the risk of metabolic syndromes, and the related complications of type 2 diabetes [22]. Gallic and protocatechuic acids have activities that inhibit  $\alpha$ -amylase, which is key enzymes for the digestion of dietary carbohydrates [23].

## CONCLUSIONS

There are several reports which supports the antioxidant activities and antidiabetic activities of *Flacourtia species* but, there are no previous reports, to the best of our knowledge, about *F. montana*. This

paper provides a strong evidence for the antioxidant activities and antidiabetic activities of *F. montana*. The work herein indicates that enzyme assisted extraction has higher antioxidant activity and antidiabetic activities in vitro, in comparison with aqueous extract. The knowledge obtained from this study should be helpful for further exploitation and application of this resource. Hence, further, studies has to be focused on the optimization of extraction process in respective of time, temperature and enzyme concentration and bioactive compounds that are responsible for the inhibition of  $\alpha$ -amylase activity.

## Acknowledgement

The authors are grateful to the Prof. Dr. Sharanappa P, Department of Bioscience, Hemagangotri University, Hassan, Karnataka for helping in the identification of plant.

## REFERENCES

1. Patridge E, Gareiss P, Kinch MS and Hoyer D. An analysis of FDA approved drugs: natural products and their derivatives. Drug Discov Today, (21):204–207, (2016)
2. Acosta-Estrada B A, Gutiérrez-uribe J A and Serna-Saldívar S O. Bound phenolics in foods. A review. Food Chem, (152): 46–55, (2014)
3. Ayoub M, de Camargo A C, Fereidoon S. Antioxidants and bioactivities of free, esterified and insoluble-bound phenolics from berry seed meals. Food Chemistry, (197):221–232, (2016)
4. Feng Li, Yi-Dan Mao, Yi-Fan Wang, Aun Raza, Li-Peng Qiu and Xiu-Quan Xu. Optimization of Ultrasonic-Assisted Enzymatic Extraction Conditions for Improving Total Phenolic Content, Antioxidant and Antitumor Activities In Vitro from *Trapa quadrispinosa* Roxb. Residues. Molecules, (22):396, (2017)
5. Huynh NT, Smagghe G, Gonzales GB, Van Camp J and Raes K. Enzyme-assisted extraction enhancing the phenolic release from cauliflower (*Brassica Oleracea* L. Var. Botrytis) outer leaves. Journal of Agricultural and Food Chemistry, (62): 7468–7476, (2014)
6. Hammi KM, Jdey A, Abdelly C, Majdoub H, Ksouri R. Optimization of ultrasound-assisted extraction of antioxidant compounds from *tunisian zizyphus lotus* fruits using response surface methodology. Food Chemistry, (184): 80–89, (2015)
7. Shobha G, Shashidhara KS, Vishwaprakash Mahadimane and Ananda S. Plant Pathovars Inhibition from Copper Based Nanoparticles Synthesized from Leaf Extract of *Flacourtia montana*. Journal of Bionanoscience, (11):1–8, (2017)

8. Garima Kulshreshtha, Anne-Sophie Burlot, Christel Marty, Alan Critchley, Jeff Hafting, Gilles Bedoux, Nathalie Bourgogne and Balakrishnan Prithiviraj. Enzyme-Assisted Extraction of Bioactive Material from *Chondrus crispus* and *Codium fragile* and Its Effect on *Herpes Simplex Virus* (HSV-1). *Marine Drugs*, 13(1): 558–580, (2015)
9. Valdés A, Vidal L, Beltrán A, Canals A, Garrigós M C. Microwave-assisted extraction of phenolic compounds from almond skin byproducts (*Prunus Amygdalus*): A multivariate analysis approach. *J. Agric. Food Chem.*, (63):5395–5402, (2015)
10. Venkanna Banothu, Chandrasekharnath Neelagiri, Uma Adepally, Jayalakshmi Lingam and Kesavaharshinibommarreddy. Phytochemical screening and evaluation of in vitro antioxidant and antimicrobial activities of the indigenous medicinal plant *Albizia odoratissima*. *Pharmaceutical Biology*, 55(1): 1155–1161, (2017)
11. Shobha G, Soumya C, Shashidhara KS and Vinutha Moses. Phytochemical profile, antibacterial and antidiabetic effects of crude aqueous leaf extract of *Datura stramonium*. *Pharmacophore*, 5 (2): 273–278, (2014)
12. Matthias Onyebuchi Agbo, Philip Felix Uzor, Uchenna Nneamaka Akazie-Nneji, Chidozie Uzoma Eze-Odurukwe, Uchenna Basilia Ogbatue and Emmanuel Chukwunwike Mbaaji. Antioxidant, Total Phenolic and Flavonoid Content of Selected Nigerian Medicinal Plants Dhaka Univ. *J. Pharm. Sci.*, 14(1): 35–41, (2015)
13. Arunachalam Sathivelu, Sundaramoorthy Sangeetha, Rastogi Archit, Sathivelu Mythili. In Vitro anti-diabetic activity of aqueous extract of the medicinal plants *Nigella sativa*, *Eugenia jambolana*, *Andrographis paniculata* and *Gymnema sylvestre*. *International Journal of Drug Development & Research*, 5(2):323–328, (2013)
14. Ramana Murty Kadali SLDV, Mangala Charan Das, Rajagopalan Vijayaraghavan and Ittagi Shanmukha. *In vitro* evaluation of antidiabetic activity of aqueous and ethanolic leaves extracts of *Chloroxylon swietenia*. *National Journal of Physiology, Pharmacy and Pharmacology*, 7(5): 486–490, (2017)
15. Saraswathy M, Kalimuthu K, Chinnadurai V and Sharmila Juliet Y. In vitro Evaluation of Antidiabetic and Anti-Inflammatory Activity of Ethanolic and Methanolic Extracts of *Ceropegia juncea*. *British Journal of Pharmaceutical Research*, 15(2): 1–9, (2017)
16. Gopi Chand Kota, Karthikeyan M, Kannan M and Rajasekar. *Flacourtia indica* (Burm. f.) Merr.-A Phytopharmacological Review. *International Journal of Research in Pharmaceutical and Biomedical Sciences*, 3(1): 78–81, (2012)
17. Sanjeeb Kumar Patro, Padmacharan Behera, Posa Mahesh Kumar, Sasmal D, Ranjan Kumar Padhy and Dash SK. Pharmacological Review of *Flacourtia sepiaria* (Ruxb.). *Sch. Acad. J. Pharm.*, 2(2):89–93, (2013)
18. Sumczynski D, Kotásková E, Družbík H and Miček J. Determination of contents and antioxidant activity of free and bound phenolics compounds and in vitro digestibility of commercial black and red rice (*Oryza sativa* L.) varieties. *Food Chem.*, (211):339–346, (2016)
19. Hossain M A, Shah M D, Gnanaraj C, Iqbal M. In vitro total phenolics, flavonoids contents and antioxidant activity of essential oil, various organic extracts from the leaves of tropical medicinal plant *Tetradlea* from Sabah. *Asian Pacific Journal of Tropical Medicine*, (4):717–721, (2011)
20. Narkhede M B, Ajimire P V, Wagh A E, Mohan M and Shivashanmugam A T. In vitro antidiabetic activity of *Caesalpinia digyna* (R.) Methanol root extract. *Asian J Plant Sci Res.*, 1(2):101–6, (2011)
21. Uddin N, Hasan MR, Hossain MM, Sarker A, Hasan AH and Islam AF. In vitro  $\alpha$ -amylase inhibitory activity and *in vivo* hypoglycemic effect of methanol extract of *Citrus macroptera* Montr. *Fruit Asian Pac J Trop Biomed.*, 4(6):473–9, (2014)
22. Lin D, Xiao M, Zhao J, Li Z, Xing B, Li X, Kong M, Li L, Zhang Q and Liu Y. An overview of plant phenolic compounds and their importance in human Nutrition and management of type 2 diabetes. *Molecules*. (21):1374, (2016)
23. Lee WJ, Ou HC, Hsu WC, Chou MM, Tseng JJ, Hsu SL, Tsai KL, Sheu WH. Ellagic acid inhibits oxidized LDL-mediated LOX-1 expression, ROS generation, and inflammation in human endothelial cells. *Journal of Vascular Surgery*, 52(5): 1290–1300, (2010)

**\*Corresponding Author:**

**Shobha G\***

Email: [shobhag@sapthagiri.edu.in](mailto:shobhag@sapthagiri.edu.in)