



## VALIDATED STABILITY INDICATING METHODS FOR *THYMOQUINONE* USING HPLC AND HPTLC

Mrinalini C. Damle\*, Aparna Zirange and Shivani Jadhav

All India ShriShivaji Memorial Society's College of Pharmacy, (Affiliated to Savitribai Phule Pune University)  
Kennedy Road, Near RTO, Pune-411001.

\*Corresponding Author Email: [mcdamle@rediffmail.com](mailto:mcdamle@rediffmail.com)

### ABSTRACT

A stability indicating assay method was developed and validated according to the ICH guidelines for estimation of Thymoquinone using HPLC & HPTLC. HiQSil C18 column was used with Acetonitrile as mobile phase. Wavelength selected was 253 nm. HPTLC analysis was done using precoated silica plates. Toluene was used as mobile phase & detection wavelength was 253 nm. Drug was subjected to forced degradation studies. Validation was carried according to ICH guidelines.

### KEY WORDS

Thymoquinone, HPLC, HPTLC, Stability indicating.

### INTRODUCTION

Among various medicinal plants, *Nigella sativa* (*N. sativa*) (Family Ranunculaceae) is emerging as a miracle herb with a rich historical and religious background since many researches revealed its wide spectrum of pharmacological potential. *N. sativa* is commonly known as black seed. *N. sativa* is native to Southern Europe, North Africa and Southwest Asia and it is cultivated in many countries in the world like Middle Eastern Mediterranean region, South Europe, India, Pakistan, Syria, Turkey, Saudi Arabia (1). Thymoquinone chemically is 2-Isopropyl-5-methyl-1,4-benzoquinone. It is very important constituent in *Nigella sativa* seeds (2). Besides being used as a spice and a condiment, *N.S.* seeds have been used for medicinal purposes in many Middle Eastern and Far Eastern countries for more than two thousand years. It is very popular in various traditional systems of medicine like Unani and Tibb, Ayurveda and Siddha. Seeds and oil have a long history of folklore usage in various systems of medicines and food. The seeds of *N. sativa* have been widely used in the treatment of different diseases and ailments. In

Islamic literature, it is considered as one of the greatest forms of healing medicine. It has been recommended for using on regular basis in Tibb-e-Nabwi (Prophetic Medicine). It has been widely used as antihypertensive, liver tonic, diuretic, digestive, anti-diarrheal, appetite stimulant, analgesics and in skin disorders. Literature survey reveals that certain pharmacological studies (3,4) have been reported on *Nigella sativa* (5-7). But no stability indicating HPLC & HPTLC methods are reported for marker Thymoquinone.




Fig 1. Structure of Thymoquinone

## MATERIAL AND METHOD

### Plant material

Seeds of *NIGELLA SATIVA* (BLACK CUMMIN) was purchased from Vishal-Chem Mumbai.

### Chemicals

Acetonitrile, Methanol & Toluene were purchased from LOBA Chemie (Mumbai).

### INSTRUMENTS

#### HPLC

JASCO-PU 2080 PLUS intelligent HPLC pump.

MD 2010 PDA detector

Borwin- PDA software (version 1.5)

Rheodyne injector port with 20 µl loop size

Hamilton Syringe (100 µl)

#### HPTLC

CamagLinomat 5 (Semiautomatic Spotting device)

Camag Twin trough chamber (10 x10 cm)

Camag TLC Scanner-3

Camag Wincats Software (version 1.4.3.6336)

Hamilton Syringe (100 µl)

All weighing were done on electronic analytical balance. (Shimadzu AY 120)

### Extraction procedure

50 gm of seeds were soaked overnight in 100 ml methanol in stoppered flask. Filtered through whatmann paper.

### Optimised HPLC Chromatographic Conditions:

The method consisted of mobile phase of Acetonitrile (100 % v/v) with HiQsil C18 HS column (250x4.6mm; 5 µm particle size). The mobile phase was filtered through 0.45 µm membrane filter followed by sonication for 10 min using bath sonicator. The optimum wavelength selected for quantification was 253 nm with a total run time of 10 min & flow rate of 1.0 ml/min.

### Optimised HPTLC Chromatographic Conditions:

Aluminium sheets precoated with silica gel G60 F<sub>254</sub> (10 x10 cm) of layer thickness 0.2mm were used. TLC plates prewashed with Methanol & activated in oven at 60°C for 10 mins. Mobile phase consisting of Toluene with detection wavelength 253 was used. TLC chamber was saturated for 15 min.

### Preparation of solutions

#### Preparation of standard stock solution of drugs

Standard stock solution of drug Thymoquinone of concentration of 1000 µg/ml (1000ppm) was prepared in methanol by dissolving 10 mg of drug in 10 ml of methanol. The standard stock solution was diluted with

the mobile phase to get solution in concentration ranging from 5 µg/ml to 30 µg/ml.

### Forced degradation studies (8)

To provide an indication of the stability-indicating ability and specificity of the proposed method forced degradation studies were performed. Hydrolysis under acidic, alkaline & neutral, oxidative, thermal and photolytic degradation studies were conducted on Thymoquinone.

#### Acid degradation

Thymoquinone working standard was weighed accurately & dissolved in MeOH & volume was made up to obtain 100 µg/ml. From that 1.5 ml of sample was pipetted out. 1ml of 0.1 N HCL was added. Volume was made up to 10 ml & sample aliquots were kept for instant degradation & examined. Sample was neutralized using 0.1 N NaOH & volume was made up with methanol.

#### Alkaline degradation

From 100 µg/ml stock solution 1.5 ml of sample was pipetted out. 1ml of 0.1 N NaOH was added. Volume was made up to 10 ml & sample aliquots were kept for instant degradation & examined. Sample was neutralized using 0.1 N HCL before injection.

#### Oxidative degradation

From 100 µg/ml stock solution 1.5 ml of sample was pipetted out. 1ml of 0.6% H<sub>2</sub>O<sub>2</sub> was added. Volume was made up to 10 ml & sample aliquots were kept for instant degradation & examined.

#### Neutral degradation

From 100 µg/ml stock solution 1.5 ml (15µg/ml) of sample was pipetted out. 1ml of distilled Water was added. Volume was made up & sample aliquots were kept for overnight & instant degradation & examined.

#### Thermal Degradation

Thymoquinone was exposed to 50 °C for 4 & 8 hours. 10 mg of sample was weighed & dissolved in methanol & diluted approximately to get 20 µg/ml of sample was pipetted out examined.

#### Photolytic degradation

Standard of thymoquinone was exposed to UV light for 200-watt hours/square meter and for fluorescence 1.2million Lux/hours. The samples were allowed to cool. 10 mg of standard thymoquinone was and volume was makeup with methanol. Further diluted with methanol to attain the working standards of 15 µg/ml concentration.

## RESULTS AND DISCUSSION




Figure 2. Standard chromatogram of thymoquinone (5ug/ml)

Table 1. System suitability of 5 ug/ml Thymoquinone chromatogram

| Sr.No. | Conc (ug/ml) | Rt    | Area       | Plates    | Resolution | Asymmetry |
|--------|--------------|-------|------------|-----------|------------|-----------|
| 1.     | 5 ug/ml      | 3.280 | 646714.429 | 14937.012 | 1.77       | 1.52      |

Fig 3. Standard densitogram of thymoquinone (500 – 2500 ng/band)




Table 2. Linearity of Thymoquinone

| Sr. no. | Amount spotted (ng/band) | Area    |
|---------|--------------------------|---------|
| 1       | 500                      | 12130.4 |
| 2       | 1000                     | 18473.1 |
| 3       | 1500                     | 22927.6 |
| 4       | 2000                     | 25020.4 |
| 5       | 2500                     | 26195.3 |

**Table 3. Forced degradation studies of Thymoquinone (HPLC)& (HPTLC)**

| Sr. No. | Conditions                                | HPLC  |           |            | HPTLC |           |            |
|---------|-------------------------------------------|-------|-----------|------------|-------|-----------|------------|
|         |                                           | % rec | Peak tail | Peak front | % rec | Peak tail | Peak front |
| 1.      | 15 ug/ml                                  | -     | 633.67    | 991.94     | -     | 0.996     | 0.995      |
| 2.      | 0.01 N HCL (INST)                         | 63.34 | 735.77    | 977.93     | 73.23 | 0.995     | 0.994      |
| 3.      | 0.01 N NaOH (INST)                        | 49.57 | 574.66    | 799.80     | 46.02 | 0.992     | 0.996      |
| 4.      | 0.6% H <sub>2</sub> O <sub>2</sub> (INST) | 52.84 | 644.96    | 521.75     | 60.50 | 0.993     | 0.992      |
| 5.      | Neutral                                   | 82.88 | 773.45    | 705.65     | 85.16 | 0.998     | 0.995      |
| 6.      | Thermal 50 °C (4hrs)                      | 44.53 | 525.33    | 982.61     | 49.24 | 0.999     | 0.998      |
| 7.      | UV (200 watt/ hrs)                        | 60.53 | 700.09    | 883.65     | 61.38 | 0.995     | 0.994      |
| 8.      | Fluorescence<br>(1.2 million lux hours)   | 63.30 | 538.73    | 940.22     | 66.09 | 0.998     | 0.997      |

#### STATISTICAL TREATMENT OF ANALYTICAL DATA:

Chromatographic methods validated statistically by applying student t test, to check statistical difference in these method and results were found as follows,

#### Student t test

Comparison between % recovered after stress degradation by HPLC and HPTLC methods for **Thymoquinone** done by applying paired t test using formula,

$$t = [d/(s/\sqrt{n})].$$

**Table 1.18: Statistical treatment for comparison of HPLC and HPTLC method for estimation of Thymoquinone**

| Parameters      | HPLC (x) | HPTLC (y) | d = (y - x)                        | (d - $\bar{d}$ ) | (d - $\bar{d}$ ) <sup>2</sup>                    |
|-----------------|----------|-----------|------------------------------------|------------------|--------------------------------------------------|
| Acid hydrolysis | 67.34    | 73.23     | 5.89                               | 2.94             | 8.67                                             |
| Base hydrolysis | 49.57    | 46.02     | -3.55                              | -6.50            | 42.25                                            |
| Oxidation       | 52.84    | 60.50     | 7.65                               | 4.70             | 22.17                                            |
| Neutral         | 82.88    | 85.16     | 2.28                               | -0.66            | 0.43                                             |
| Thermal         | 44.53    | 49.24     | 4.70                               | 1.75             | 3.09                                             |
| UV              | 60.53    | 61.38     | 0.85                               | -2.09            | 4.37                                             |
| Fluorescence    | 63.30    | 66.09     | 2.78                               | -0.15            | 0.02                                             |
|                 |          |           | <b><math>\bar{d} = 2.94</math></b> |                  | <b><math>\sum (d - \bar{d})^2 = 81.02</math></b> |

Null hypothesis (H<sub>0</sub>) i.e. there is no significant difference between mean results by two methods.

$$S^2 = (1/n-1) \sum (d - \bar{d})^2$$

$$S^2 = 13.50$$

$$t = [d/(s/\sqrt{n})]$$

$$t = 0.4684$$

**Calculated |t| is less than tabulated t (2.447), H<sub>0</sub> may be accepted at 5% level of significance and conclude that results by two methods do not differ significantly.**  
**METHOD VALIDATION (9)**

Both the methods were validated in terms of linearity, range, precision, accuracy, specificity, limit of detection, limit of quantitation and robustness as per ICH guidelines. The system suitability parameters were evaluated as specified in Indian Pharmacopoeia 2014.

#### Specificity:

The specificity of the method was confirmed by analyzing standard marker Thymoquinone & methanolic extracts. The spot for Thymoquinone in the

sample was confirmed by comparing the R<sub>f</sub> and spectra of the spot with that of sample. The peak purity of Thymoquinone was examined by comparing the spectra at peak start, peak middle and peak end positions of the bands.

#### Linearity & Range

Six different concentrations of (5, 10, 15, 20, 25 and 30 µg/mL) were prepared from standard stock solution of 1000 µg/mL of Thymoquinone and was analysed. For HPTLC, different volumes of 100 µg/ml standard solutions were spotted to obtain 500 to 2500 ng/ band. Linearity equation of HPTLC was found to be  $y = 7.0575x + 10653$   $r^2 = 0.940$  & of HPLC was found to be  $y = 81153x - 34654$   $r^2 = 0.959$  respectively.

#### Precision:

The precision was evaluated with respect to both repeatability and intermediate precision. Repeatability was evaluated by injecting six replicate injections of test solution of the drug Thymoquinone (5 µg/ml). The studies were repeated for three different days to

determine intermediate precision. Peak areas of the drugs were determined and % RSD was calculated

**Table 4. Precision for Thymoquinone (HPLC & HPTLC)**

| Sr. No. | Method | Amount conc ug/ml & ng/band | Mean     | SD        | % RSD    |
|---------|--------|-----------------------------|----------|-----------|----------|
| 1.      | HPLC   | 5 ug/ml                     | Interday | 660588.43 | 14652.27 |
|         |        |                             | Intraday | 421115.36 | 9031.95  |
| 2.      | HPTLC  | 500 ng/band                 | Interday | 4081      | 90.29    |
|         |        |                             | Intraday | 5807.1    | 122.09   |

#### Limit of detection (LOD) and Limit of Quantitation (LOQ):

From the linearity data the LOD and LOQ was calculated, using the formula  $LOD = 3.3 \sigma/S$  and  $LOQ = 10 \sigma/S$

where,  $\sigma$  = standard deviation of the y-intercept of linearity equations and  $S$  = slope of the calibration curve of the analyte.

**Table 5: LOD & LOQ for Thymoquinone**

| PARAMETERS | LOD          | LOQ         |
|------------|--------------|-------------|
| HPLC       | 0.59 ug/ml   | 1.80ug/ml   |
| HPTLC      | 0.11 ng/band | 0.33ng/band |

#### Robustness:

Robustness was evaluated by deliberate variation in parameters like mobile phase and flow rate for HPLC & variation of mobile phase & time from spotting to

development for HPTLC. The proposed method was found to be robust under given experimental conditions.

**Table 6: Summary of validation parameters for Thymoquinone**

| Sr.no. | Validation Parameters       | HPLC                                  | HPTLC                                  |
|--------|-----------------------------|---------------------------------------|----------------------------------------|
| 1.     | Specificity                 | Specific                              | Specific                               |
| 2.     | Linearity                   | $y = 81153x - 34654$<br>$r^2 = 0.959$ | $y = 7.0575x + 10653$<br>$r^2 = 0.940$ |
|        | Range                       | (5 -30 ug/ml)                         | (500- 2500 ng/band)                    |
| 3.     | Precision                   | Interday                              | 500 ng/band                            |
|        |                             | Intraday                              | 500 ng/band                            |
| 6.     | Limit of Detection (LOD)    | 0.59 ug/ml                            | 0.11 ng/band                           |
| 7.     | Limit of Quantitation (LOQ) | 1.80ug/ml                             | 0.33ng/band                            |
| 8.     | Robustness                  | Robust                                | Robust                                 |

#### CONCLUSION:

Thymoquinone was found be prone to degradation in acidic, alkaline, oxidation, neutral & photolytic conditions. The developed method is simple rapid & stability indicating.

#### ACKNOWLEDGEMENT:

Authors are thankful to Savitribai Phule Pune University for funding this work & to the Principal and the management of AISSMS College of Pharmacy, Pune for providing the necessary facilities for research work.

#### REFERENCES:

- EnasM, Shifaa I., A study of the chemical composition and the biological active components of *Nigella Sativa* and *trigonella foenum - graecum* L. seeds: Journal Of Applied Chemistry, 6 (6) : 2278-5736, (2014)
- Aftab A., Asif H. et.al., A review on therapeutic potential of *Nigella sativa*: A miracle herb: Asian Pacific journal Of Tropical Biomedicine, 3(5): 337-352, (2013)
- Ramlah M. I., Nurul S. H. et al., A randomised controlled trial on hypolipidemic effects of *Nigella Sativa* seeds powder in menopausal women: Journal of Translational Medicine, 1-7, (2014)
- Emine O., Muhsine D., Effect of black cumin oil (*Nigella sativa* L.) on fresh fish (*Barbus grypus*) fillets during

- storage at  $2 \pm 1$  °C: Food Science & Technology, 37(1): 148-152, (2017)
5. Muhammad S. Ahamad B. et al., Stability Study of Algerian *Nigella sativa* Seeds Stored under Different Conditions: Journal of Analytical Methods in Chemistry, 1-12, (2017)
  6. Snehalatha B, Momin M, Mishal V and Kale T R., Development and validation of spectrophotometric method for simultaneous estimation of nigella sativa seed oil and ginger extract in the same dosage form: International Journal of Pharmaceutical Sciences & Research, 5 (12): 5235-52 (2014)
  7. Krishnaveni M. and Saranya S., Phytoconstituent Analysis of *Nigella Sativa* Seeds using Analytical Techniques. Bulletin of Environment: Pharmacology & Life Sciences, 5(3): 25-38, (2016).
  8. ICH guidelines, Q1A(R2); Stability Testing of New Drug Substances and Products, International Council for Harmonisation Geneva. (<http://www.ich.org/products/guidelines/quality/article/quality-guidelines/html>), 2003.
  9. ICH guidelines, Q2 (R1): Validation of Analytical Procedures: Text and Methodology, International Conference on Harmonization Geneva. (<http://www.ich.org/products/guidelines/quality/article/quality-guidelines/html>), 2005.

Received:04.08.18, Accepted: 07.09.18, Published:01.10.2018

**\*Corresponding Author:**

**Mrinalini C. Damle\***

Email: [mcdamle@rediffmail.com](mailto:mcdamle@rediffmail.com)