



Orchid Mycorrhizal Studies of *Nervilia crociformis*

Jyothsna B.S.¹, Kavitha K.R.², and Keshamma E.^{3*}

^{1,2}Associate Professor, Department of Botany & PG Studies, Nrupathunga University, Bengaluru, Karnataka, India

³Associate Professor, Department of Biochemistry, Maharani Cluster University, Palace Road, Bengaluru, Karnataka, India

Received: 12 Jul 2021/ Accepted: 4 Aug 2021 / Published online: 1 Oct 2021

*Corresponding Author Email: keshamma.blr76@gmail.com

Abstract

Orchid mycorrhizal associations involve a plethora of distinctive nutrient transport systems, structures and phenomena which have only been observed in the family Orchidaceae. Orchid mycorrhizae are symbiotic relationships between the roots of plants of the family Orchidaceae and a variety of fungi. Hence, current study aimed to isolate and purify the mycorrhizal fungi of orchid from natural habitats of Western Ghats. The terrestrial orchid *Nervilia crociformis*, was collected from natural habitats from various locations of Kemmangundi and Sagar, Karnataka State and were investigated for mycorrhizal association. Freshly collected root/tuber of *Nervilia crociformis* was used for isolation, identification and purification. The fungal isolates were purified by hyphal tip method. Classical identification techniques used in the identification of fungi include colony characters and microscopic characters of hyphae and spores. One fungal isolate was obtained from the tuber of *Nervilia crociformis* which was identified as *Colletotrichum fruticola*.

Keywords

Terrestrial orchid, *Nervilia crociformis*, *Colletotrichum fruticola*, fungal isolate

INTRODUCTION

The term *mycorrhiza* was coined by Frank for the fungus infested roots [1]. Mycorrhizae are symbiotic associations between specialized soil fungi and plants. They are primarily responsible for nutrient transfer and are essential for one or both the partners. Most plants in natural ecosystems have mycorrhizal association when symbiotic fungi inhabit healthy tissues of the root of terrestrial plants and play an important role in their successful colonization. Bernard was the first to isolate *Rhizoctonia*- like fungi by surface sterilizing the root fragments and aseptically plating on nutrient agar medium [2]. Recently isolation from single peloton was attempted which is a tedious process [3].

Isolates of *Rhizoctonia* sp. obtained from terrestrial orchids *Dactylorhiza sambucina*, *D. maculata* and *Platanthera bifolia* collected from Italy were studied for macro and microscopic morphology, growth rate, ultrastructure of septal pore apparatus. On the basis of cultural and morphometric characters the isolates were divided into 3 groups. No perfect stage was produced in culture; group I were identified as species of *Ceratobasidium*, group II as species of *Thanetophorus* and group III as species of *Tulasnella* or *Sebacina*. According to Currah et al *Rhizoctonia panaicula* Currah, sp. now represented by 5 different isolates had a unique morphology of monilioid cells was isolated from *Platanthera obtusata*, *Ceratobasidium obscurum* from *Amerorchis*

rotundifolia, *Phialocephala fortini* from *Calypso bulbosa* [4]. Two new species of fungi mycorrhizal with terrestrial orchids native to Canadian priaries have been reported viz. *Ceratorhiza pernacatena* sp.nov. from mycorrhiza of *Platanthera praeclara*. *Epulorhiza calendulina* sp.nov. from mycorrhiza of *Amerorchis rotundifolia* [5].

These fungi were generally identified based on culture characters, microscopic study of hyphae, their associated structures. A key to the genera of fungi mycorrhizal with orchids was provided by Currah & Zelmer [6]. Sneh et al have given a guide to identify *Rhizoctonia* species [7]. A check list of *Rhizoctonia* epithets was provided by Anderson [8]. Recently orchid mycobionts being identified using molecular biology technique by sequencing the internal transcribed spacer (ITS) of the nuclear ribosomal DNA after PCR amplification using a variety of primer combinations has been the method routinely used to identify orchid mycobionts from cultured fungi or directly from orchid protocorms, roots, tubers and rhizomes [9,10]. Both universal fungal ITS primers and Tulasnaceae- specific PCR primers were being used [11,12]. Suarez *et al.*, (2006), Martos et al used 5.8 S- Tul to amplify the 5' part of 28 S rDNA to work on range of clades of Tulasnaceae and Ascomycete mycobionts [13,14]. ITS sequencing and cloning also revealed many endophytes associated with orchid roots [15-18]. Review of literature revealed that there are only a few reports on the study of orchid mycorrhiza especially from India [19-21]. Hence in the present work we attempted to study terrestrial orchid mycorrhizae of Karnataka province.

MATERIALS AND METHODS

The materials and methods used in the present investigation on the study of mycorrhizae in terrestrial orchids included collection of orchid plants root and tuber of selected ground orchids.

Isolation of associated mycorrhizal fungi

Standard procedures were used for isolation of mycorrhizal fungi by plating portions of surface-sterilized roots/tubers on suitable media [22]. Fresh root/tuber samples of terrestrial orchids were collected from their natural habitats by carefully removing the soil so as to keep the underground system intact. Roots and tuber (if present) of the orchid plants were used separately for the purpose of isolation. The sample surface was first washed with running tap water several times carefully to get rid of soil particles and organic debris. The roots were then surface sterilized by submerging them in 0.1% mercuric chloride solution for 3-5 minutes washed in three changes of sterile distilled water, 5-10 mm

portions of the roots were then cut, aseptically, plated on Potato dextrose agar (PDA) medium containing streptomycin (50µg/mL) and incubated at 25°C. Fungi grown from the cut ends were used for obtaining the pure culture. These isolates were further grown on PDA at 25°C for 4 weeks under normal diurnal conditions to allow sclerotia and chlamydospores to form and mature before identification was attempted.

Identification of associated mycorrhizal fungi by classical method

The taxonomic identification of an organism is an essential part of the investigation of a new species or isolate of a known species. It is imperative that the organism is correctly identified so that when reference is made to published material, the experimental data becomes valid and may be used and compared by other investigators. Classical identification techniques used in the identification of fungi include colony characters and microscopic characters of hyphae and spores. In the present work, attempts were made to identify all the isolates obtained by studying the similarities and differences between the isolates in the following characters: (a) colony characters- including growth rate, colour, shape and texture of fungal colony; sclerotia present or absent; colour changes in the substrate (medium) caused by the fungus (b) Microscopic characters involving colour, size, septation, branching of the hyphae, formation of rhizomorphs, sclerotia, and sporulation if any. Of these characters, the size and shape of chlamydospores, the size and texture of sclerotia, characteristics of hyphae and spores as described by Barnett & Hunter were given special consideration [23].

Purification and maintenance of fungal isolates

The fungal isolates were purified by hyphal tip method. They were maintained on PDA slants in the refrigerator for further studies. This technique was used for species which frequently develop degenerate cultural variants from germinated conidia, or fungi such as *Rhizoctonia* which do not sporulate in culture.

RESULTS AND DISCUSSION

The terrestrial orchid, selected for mycorrhizal study was *Nervilia crocifformis* was collected from Sagar, Shimoga district, Karnataka). They were identified using taxonomic keys published in various Floras like Abraham & Vatsala, Santapau & Kapadia and Ananda Rao [24-26].

Isolation of associated mycorrhizal fungi

The fungal isolates were obtained from the root/tuber/ of the terrestrial orchid. The fungi grew from the cut ends of the root/tuber as white cottony

growth. Pure cultures were obtained by hyphal tip method and were maintained in PDA slants for further studies. These were preliminarily identified using colony characters and morphological characteristics of hyphae and spores if any as described by Barnett & Hunter [23].

Identification of associated mycorrhizal fungal isolates by classical method

The fungal isolate from the tuber of *Nervilia crociformis* showed *Colletotrichum fruticola* (Plate 1


A-E). Colonies on PDA at first white, becoming grey to dark grey at the centre with age, in reverse greyish green with white halo becoming black with age, maximum of 8.3cm in diam. in 7 days. Aerial mycelium pale grey, dense, cottony, without visible conidial masses. Sclerotia absent. Conidia 9.5-14 x 3-4.5µm, one- cell, smooth walled with a large guttule at the centre and surrounded by smaller guttules, hyaline, cylindrical with obtuse to slightly rounded cells, sometimes oblong.



A:Habit of *Nervilia crociformis*




B:Tuber of *Nervilia crociformis*



C:Isolate of *Nervilia crociformis* (root)



D:Isolate of *Nervilia crociformis* (tuber)



E: Conidia of *Colletotrichum fruticola* (400x)

Plate 1

CONCLUSION

In conclusion, the terrestrial orchid identified as *Nervilia crociformis*, was collected from natural habitats from various locations of Kemmangundi and Sagar, Karnataka State and were investigated for mycorrhizal association. The fungal isolate from the tuber of *Nervilia crociformis* was identified as *Colletotrichum fruticola*.

REFERENCES

1. Frank B. Über die auf Wurzelsymbiose beruhende Ernährung gewisser Bäume durch unterirdische Pilze. 1885.
2. Bernard N. Recherches expérimentales sur les Orchidées. Revue générale de Botanique. 1904a;16(405-451):458-76.
3. Kristiansen KA, Taylor DL, Kjølner R, Rasmussen HN, Rosendahl S. Identification of mycorrhizal fungi from single pelotons of *Dactylorhiza majalis* (Orchidaceae) using single-strand conformation polymorphism and mitochondrial ribosomal large subunit DNA sequences. Molecular Ecology. 2001;10(8):2089-93.
4. Currah RS. Thanatephorus pennatus sp. nov. isolated from mycorrhizal roots of *Calypso bulbosa* (Orchidaceae) from Alberta. Canadian Journal of Botany. 1987;65(9):1957-60.
5. Zelmer CD, Currah RS. *Ceratophora pernatatena* and *Epulorhiza calendulina* spp. nov.: mycorrhizal fungi of terrestrial orchids. Canadian Journal of Botany. 1995;73(12):1981-5.
6. Currah RS, Zelmer C. A key and notes for the genera of fungi mycorrhizal with orchids and a new species

- in the genus *Epulorhiza*. Reports of the Tottori Mycological Institute (Japan). 1992.
7. Sneh B, Burpee L, Ogoshi A. Identification of *Rhizoctonia* species. APS press; 1991.
 8. Andersen TP. A check-list of *Rhizoctonia* epithets. Mycotaxon. 1994; 51:437-57.
 9. White TJ, Bruns T, Lee SJ, Taylor J. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. PCR protocols: a guide to methods and applications. 1990;18(1):315-22.
 10. Gardes M, Bruns TD. ITS primers with enhanced specificity for basidiomycetes-application to the identification of mycorrhizae and rusts. Molecular ecology. 1993;2(2):113-8.
 11. Bidartondo MI, Burghardt B, Gebauer G, Bruns TD, Read DJ. Changing partners in the dark: isotopic and molecular evidence of ectomycorrhizal liaisons between forest orchids and trees. Proceedings of the Royal Society of London. Series B: Biological Sciences. 2004;271(1550):1799-806.
 12. Selosse MA, Faccio A, Scappaticci G, Bonfante P. Chlorophyllous and achlorophyllous specimens of *Epipactis microphylla* (Neottieae, Orchidaceae) are associated with ectomycorrhizal septomycetes, including truffles. Microbial Ecology. 2004;47(4):416-26.
 13. Suárez JP, Weib M, Abele A, Garnica S, Oberwinkler F, Kottke I. Diverse tulasnelloid fungi form mycorrhizas with epiphytic orchids in an Andean cloud forest. Mycological research. 2006 Nov 1;110(11):1257-70.
 14. Martos F, Munoz F, Pailler T, Kottke I, Gonneau C, SELOSSE MA. The role of epiphytism in architecture and evolutionary constraint within mycorrhizal networks of tropical orchids. Molecular Ecology. 2012;21(20):5098-109.
 15. Bidartondo MI, Burghardt B, Gebauer G, Bruns TD, Read DJ. Changing partners in the dark: isotopic and molecular evidence of ectomycorrhizal liaisons between forest orchids and trees. Proceedings of the Royal Society of London. Series B: Biological Sciences. 2004;271(1550):1799-806.
 16. Julou T, Burghardt B, Gebauer G, Berveiller D, Damesin C, Selosse MA. Mixotrophy in orchids: insights from a comparative study of green individuals and nonphotosynthetic individuals of *Cephalanthera damasonium*. New Phytologist. 2005;166(2):639-53.
 17. Abadie JC, Püttsepp Ü, Gebauer G, Faccio A, Bonfante P, Selosse MA. *Cephalanthera longifolia* (Neottieae, Orchidaceae) is mixotrophic: a comparative study between green and nonphotosynthetic individuals. Botany. 2006;84(9):1462-77.
 18. Roy M, Yagame T, Yamato M, Iwase K, Heinz C, Faccio A, Bonfante P, Selosse MA. Ectomycorrhizal *Inocybe* species associate with the mycoheterotrophic orchid *Epipogium aphyllum* but not its asexual propagules. Annals of Botany. 2009;104(3):595-610.
 19. Vij SP, Sharma M, Datta SS. Mycorrhizal association in North Indian Orchidaceae: a morphological study. Bibliotheca Mycologia. 1983; 91:467-73.
 20. Sharma J, Ishida M, Yadon V. Mycorrhizal diversity of an endemic terrestrial orchid. Lankesteriana. 2007.
 21. Aggarwal S, Zettler LW. Reintroduction of an endangered terrestrial orchid, *Dactylorhiza hatagirea* (D. Don) Soo, assisted by symbiotic seed germination: First report from the Indian subcontinent. Nature and Science. 2010;8(10):139-45.
 22. Harvais G, Hadley G. The relation between host and endophyte in orchid mycorrhiza. New Phytologist. 1967;66(2):205-15.
 23. Barnett HL, Hunter BB. Illustrated genera of imperfect fungi. Illustrated genera of imperfect fungi. 1972(3rd ed).
 24. Abraham A. Introduction to orchids with illustrations and descriptions of 150 South Indian Orchids. Tropical Botanic Garden and Research Institute; 1981.
 25. Santapau H and Kapadia Z. Orchids of Bombay. Government Press, India. 1966.
 26. Ananda Rao T. Conservation of Wild orchids of Kodagu in the Western Ghats, KAAS, Bangalore. 1988.