

Research Article | Biological Sciences | Open Access | MCI Approved UGC Approved Journal

# Synthesis, Characterization, and Antimicrobial Activity of Schiff Bases Derived From 4, 6 - Difluoro - 2 - Amino Benzothiazole

# Raksha V. Zala and Surbhi V. Upadhyay

Department of Chemistry, Kamani Science College and Prataprai Arts College, Saurashtra University, Amreli-365601, Gujarat, India

> Received: 8 Oct 2018 / Accepted: 9 Nov 2018 / Published online: 1 Jan 2019 Corresponding Author Email: <u>drkdbhatt@oulook.com</u>

# Abstract

Some novel Schiff bases namely *N*-(4-(dimethyl amino)benzylidene)-4,6-difluoro benzothiazole-2-amine; 4-bromo-2-(((4,6-difluoro benzothiazole-2-yl) imino) methyl)phenol and *N*-(3,4dimethoxybenzylidene)-4,6-difluoro benzothiazole-2-amine have been synthesized from condensation reaction of 4,6-difluoro-2-amino benzothiazole with different aromatic aldehydes like 4-(dimethyl amino)benzaldehyde, 5-bromo-2-hydroxy benzaldehyde and 3,4-dimethoxy benzaldehyde . The synthesized Schiff bases were characterized by elemental analysis, FT-IR, Mass and <sup>1</sup>H NMR spectroscopy. Antimicrobial and Antifungal activities of these Schiff bases were tested using MIC determination against gram positive (*S. Aureus, B.Subtilis*), gram negative bacteria (*S. Marcescens, E. Coli*) and *Rhizopus sp. and A. Niger*.

#### Keywords

Schiff base, Benzothiazole derivative, Antimicrobial activity, Antifungal activity, MIC determination.

#### **1. INTRODUCTION**

Schiff bases derived from an amine and aldehyde are an important class of ligands in coordination chemistry [1,2] that have been studied extensively. They were first reported by a German chemist Hugo Schiff [3] Nobel Prize winner in 1864. Schiff bases are characterized by imine or azomethine group (-CH=N) [4] and their general structural formula is R<sub>1</sub>-CH=N-R<sub>2</sub> where R<sub>1</sub> and R<sub>2</sub> are alkyl, aryl or heterocyclic group having different substituents. Schiff bases coordinate to transition metal ions [5-9] via azomethine nitrogen and these complexes are play an important role in the development of coordination chemistry [10]. Presence of azomethine group helps to explain the mechanism of racemization and transamination reaction occurred in biological system [11]. Schiff bases play very important role in industrial and biological field. They are biological active compounds and possess many biological activities such as anticancer [12], antituberculosis [13], antibacterial [14], antimicrobial [15], plant growth inhibitors [16] and antiinflammatory activity [17].

Benzothiazoles are heterocyclic bicyclic ring containing nitrogen and sulfur heteroatoms in five membered ring and its ring made from 1,3-thiazole

DOI: https://doi.org/10.21276/ijpbs.2019.9.1.25

Raksha V. Zala and Surbhi V. Upadhyay 201

-

solvents employed for synthesis were commercially

available (AR grade) and used as received without

The melting points (uncorrected) were determined

using capillary and theils tube filled with paraffin oil.

Samples for infrared spectra were prepared as KBr

pallets; spectra were recorded on tensor Bruker 27

(Ettlingen, Germany) and expressed in cm<sup>-1</sup>. Mass

spectra (GC-MS) were determined using Jeol D-300

spectrometer. NMR spectra were recorded on a

model DPX 200 MHz and Advance II 500 MHz Bruker

Elemental analysis data are in accordance with the

theoretically calculated percentage of C, H, N and S. Percentage of C, H, N and S were derived using

CHN/S/O Analyzer, Perkin Elmer, series11, 2400.

**benzylidene)-4,6-difluoro benzothiazole-2-amine** To a methanolic solution of 4,6-difluoro-2-amino

benzothiazole (0.186gm,0.001mol) with 4-(dimethyl

amino) benzaldehyde (0.149gm, 0.001 mol) was

mixed using catalytic amount of glacial acetic acid.

The reaction mixture was refluxed with constant

stirring at 50-60°C for 10-12 hrs. Schiff base was

obtained by slow evaporation method at room temperature. To remove excess aldehyde, the product was washed with sodium bisulphite and then

washed with chilled methanol to remove other

impurities. The product was isolated, dried by ether

and recrystallized from hot methanol. Yield: 62.95%.

(Scheme-1) The Schiff base ligand exists in crystalline

form and golden yellow in color.

FT-NMR instruments (Ettlingen, Germany).

[A] Synthesis of N-(4-(dimethyl amino)

2.3. Synthesis of Schiff base

further purification.

2.2 Instrument



ring fused with benzene ring. These moieties are found large number of interests in medicinal and pharmaceutical field [18,19,20,21] due to their biological applications. Various substituted benzothiazoles have shown pharmacological activities such as antitumor [22,23], antimicrobial [24], antibacterial [25,26], antifungal [27], antiinflammatory [28], anticonvulsive [29], analgesic [30] and Parkinson's disease [31].

Due to their importance in pharmaceutical utilities, in the present work, synthesis of Schiff base from benzothiazole derivative amine is of considerable interest. Apart from biological applications these Schiff bases have another area of applications as corrosion inhibitor [32], dye manufacture [33], perfumery [34], plant growth regulators[35] and in analytical chemistry where some of these Schiff base compounds were used as ligand in complexometry [36] and intermediate for various synthesis[37,38].

Here an attempt has been made to synthesize and characterize novel Schiff bases derived from 4,6-difluoro benzothiazole. The biological activity of these new synthesized compounds were tested against two gram positive bacteria *S. Aureus, B.Subtilis* and two Gram negative bacteria *i.e. S. Marcescens, E. Coli* and fungal strains *Rhizopus sp.* and *A. Niger* in solvent DMF/CHCl<sub>3</sub>.

# 2.EXPERIMENTAL

#### 2.1 Chemicals and reagents:

All starting materials, reagents and other chemicals were obtained from Sigma-Aldrich and used as received. Silica gel and fluorescence active TLC plates (F-2009) were purchased from E.Merck. All the





4,6-difluorobenzothiazol-2-amine

4-(dimethylamino)benzaldehyde

DBT-I

СН<sub>3</sub> СН<sub>3</sub>

#### Scheme-1

For C<sub>16</sub>H<sub>13</sub>F<sub>2</sub>N<sub>3</sub>S, Melting point is 152-155°C and Elemental analysis data found: %C 60.45; %H 4.15; %N 13.1; %S 9.98. In FT-IR, bands observed are at 1610 cm<sup>-1</sup> v (HC=N, azomethine group), 2815 cm<sup>-1</sup> v -N(CH<sub>3</sub>)<sub>2</sub>, 1252 cm<sup>-1</sup> v(C-F), 722cm<sup>-1</sup> v(C-S-C). <sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$ N-(CH<sub>3</sub>)<sub>2</sub>, 1.63 (s, 6H);  $\delta$ Ar-H, 6.85-7.65 (d, 6H);  $\delta$ N=CH, 9.26 (s, 1H) (SI 1) and Mass spectra (GC-MS) *m/z* values of DBT-I correspond to the M<sup>-</sup>adduct (317) are shown in SI 2 respectively.

# [B] Synthesis of 4-bromo-2-(((4,6-difluoro benzothiazole-2-yl) imino) methyl) phenol

To a methanolic solution of 4,6-difluoro-2-amino benzothiazole (0.186gm,0.001mol) with 5-bromo-2hydroxy benzaldehyde (0.201 gm, 0.001 mol) was mixed using catalytic amount of glacial acetic acid. The reaction mixture was refluxed with constant stirring at 50-60°C for 10-12 hrs. Schiff base was obtained by slow evaporation method at room temperature. To remove excess aldehyde, the product was washed with sodium bisulphite and then washed with chilled methanol to remove other



impurities. The product was isolated, dried by ether and recrystallized from hot methanol. Yield: 78.65%.

**(Scheme-2)** The Schiff base ligand exists in amorphous form and yellow in color.

To a methanolic solution of 4,6-difluoro-2-amino

dimethoxy benzaldehyde (0.166 gm, 0.001 mol) was

mixed using catalytic amount of glacial acetic acid.

The reaction mixture was refluxed with constant

stirring at 30-40°C for 5-7hrs. Schiff base was

obtained by slow evaporation method at room

temperature. To remove excess aldehyde, the

product was washed with sodium bisulphite and then

washed with chilled methanol to remove other impurities. The product was isolated, dried by ether

and recrystallized from hot methanol. Yield: 55.39%.

(Scheme-3) The Schiff base ligand exists in crystalline

form and pale yellow in color.

(0.186gm,0.001mol)



+ Br

4,6-difluorobenzothiazol-2-amine

5-bromo-2hydroxybenzaldehyde

сно

OH

benzothiazole



DBT-II

with

3,4-

#### Scheme-2

For C<sub>14</sub>H<sub>7</sub>BrF<sub>2</sub>N<sub>2</sub>OS, Melting point is 178-180°C and Elemental analysis data found: %C 45.22; %H 1.93; %N 7.48; %S 8.55. In FT-IR, bands observed are at 1604 cm<sup>-1</sup> v(HC=N, azomethine group), 3127 cm<sup>-1</sup> v(OH), 1226 cm<sup>-1</sup> v(C-F), 753 cm<sup>-1</sup> v(C-S-C). <sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$ Ar-H, 6.82-7.13 (d, 5H);  $\delta$ N=CH, 7.26 (s, 1H);  $\delta$ Ar-OH , 5.72 (s, 1H) (SI 3) and Mass spectra (GC-MS) *m/z* values of DBT-II correspond to the M<sup>-</sup>adduct (368), is shown in SI 4 respectively.

# [C] Synthesis of N-(3,4-dimethoxybenzylidene)-4,6difluoro benzothiazole-2-amine





3,4-dimethoxybenzaldehyde

F N N C OCH3 F S N N C OCH3

4,6-difluorobenzothiazol-2amine

# Scheme-3

For C<sub>16</sub>H<sub>12</sub>F<sub>2</sub>N<sub>2</sub>O<sub>2</sub>S, Melting point is 192-194 °C. Elemental analysis data found: %C 57.62; %H 3.60; %N 8.15; %S 9.45. In FT-IR, bands are observed at 1611 cm<sup>-1</sup> v (HC=N, azomethine group), 1226 cm<sup>-1</sup> v(C-F), 751 cm<sup>-1</sup> v(C-S-C). <sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$ Ar(-OCH<sub>3</sub>)<sub>2</sub>, 3.95 & 3.98 (s, 6H);  $\delta$ Ar-H, 6.81-7.10 (d, 5H);  $\delta$ N=CH, 7.14 (s, 1H) (SI 5) and Mass spectra (GC-MS) m/z values of DBT-III correspond to the Madduct (335), is shown in SI 6 respectively.

# 3. RESULTS AND DISCUSSION

#### 3.1. Spectral studies of ligands

The novel Schiff base compound DBT-I, DBT-II and DBT-III are synthesized according to described process in scheme-1,2 and 3 respectively. The structure of all the compounds were evaluated on the basis of elemental analysis, FT-IR, mass and <sup>1</sup>H NMR spectral data. The ligands were insoluble in water but soluble in methanol, ethanol, chloroform, 1,4-dioxane and DMF.

In the IR spectra, Schiff bases DBT-I, DBT-II and DBT-II exhibit strong peaks in the region 1610  $\rm cm^{-1}, 1604$   $\rm cm^{-1}$  and 1611  $\rm cm^{-1}$  respectively, which can be



assigned as v(HC=N) vibration of azomethine group. Aromatic v(C-H) stretching at 3045-3040 cm<sup>-1</sup> and v(C=C) bands at 1510 cm<sup>-1</sup>, 1505 cm<sup>-1</sup>, 1511 cm<sup>-1</sup> proved existence of aromatic rings. The bands v(C-S-C) region 722-753 cm<sup>-1</sup>. The band of hydroxy functional group v(OH) of DBT-II Schiff base observed at 3127 cm<sup>-1</sup>. The band of N, N-dimethyl functional group v(N(CH<sub>3</sub>)<sub>2</sub>) of DBT-I Schiff base observed at 2815 cm<sup>-1</sup>.

The mass spectrum of Schiff bases DBT-I, DBT-II and DBT-III showed the molecular ion peak at m/z 318.1, 368.9 and 335.2 that corresponds to the molecular weight of Schiff base.

In the <sup>1</sup>H NMR spectra of benzothiazole Schiff bases, multi signals are assigned to Aromatic protons observed in the range from 6.8 to 7.65  $\delta$ ppm. A sharp singlet is observed for Schiff bases DBT-I, DBT-II and DBT-III at 9.26  $\delta$ ppm, 7.26  $\delta$ ppm and 7.14  $\delta$ ppm respectively in spectrum which corresponds to the azomethine proton. Methyl group protons for DBT-I appears as singlet at 1.63  $\delta$ ppm. The hydroxyl group proton for DBT-II is observed as a broad singlet at 5.72  $\delta$ ppm in the spectrum. In the <sup>1</sup>H NMR spectra of



DBT-III, methoxy group protons appear as singlet at 3.98  $\delta$  ppm.

#### 3.2. Antimicrobial activity

Synthesized Schiff bases were tested for their impact on some microorganisms in terms of minimum inhibitory concentration (MIC) using suspended Luria Broth in sterile double distilled water as a media. Culture for bacteria gram negative and positive was incubated for 24 hrs at 35°C, respectively. Control test with no active ingredient was also performed by adding just an equivalent amount of solvent. MIC was determined using double fold serial dilution in liquid media containing varying concentration of tested compounds from 0.1-10,000 µM. Bacterial growth was measured by the turbidity of the culture after 15 h. If particular concentration of the compound inhibited bacterial growth, half the concentration of the compound was tried. This procedure was carried out at a concentration that bacteria grow normally. The lowest concentration that inhibits the bacterial growth totally was determined as MIC value. All equipment and culture media employed during the process were sterile.

The efficiencies of the ligands and the complexes have been tested against two Gram-positive (S. aureus, B. subtilis), two Gram-negative (S. marcescens, E. coli) bacteria and two fungus (Rhizopus sp., A. niger). The data reveals that all the synthesized Schiff bases have higher antimicrobial activity.

Among the various compounds, DBT-III has been found to be the most effective against these microbes showing maximum clarity of zones. Its antibacterial activity was found maximum against Gram positive bacteria (S. aureus, B. subtilis) and Gram-negative bacteria (*E. coli*) and DBT-I also exhibited maximum inhibition against S. marcescens, and DBT-II exhibited minimum inhibition.

The growth of fungus was measured by recording diameter of fungal colony. Following relation was used to calculate the fungal growth inhibition:

Fungal growth inhibition (%) =  $[(A - B)/A] \times 100$ where A is the diameter of the fungal colony in the control plate and B is the diameter of the fungal colony in the test plate. Results are represented in figure-10. Among all compounds DBT-II, has shown maximum antifungal activity compared to DBT-I, DBT-II against A. niger while DBT-III exhibited maximum anti-fungal activity against Rhizopus sp. The comparison chart against standard antibiotics is as given in Figure-1.



Figure-1: Comparison diagram of antimicrobial activity

#### CONCLUSION

In this paper, we have described new Schiff bases, synthesized by condensation of 4,6-difluoro-2-amino benzothiazole and different aldehydes like *N*, *N*-dimethyl benzaldehyde, 5-bromo-2-hydroxy benzaldehyde and 3,4-dimethoxy benzaldehyde in an

alcoholic medium using glaciel acetic acid with good yield. The structure of Schiff bases has been confirmed by various physicochemical and spectral analyses. In this study, these Schiff base derivatives DBT-I, DBT-II and DBT-III are found active antimicrobial compounds. Among the three Schiff

www.ijpbs.com or www.ijpbsonline.com



base tested, DBT-III exhibited the promising antimicrobial activity and DBT-II is the most active against A. niger.

#### ACKNOWLEDGMENT

The authors are thankful to The principal Dr. Atul H. Patel Sir, Kamani Science College and Prataprai Arts College, Amreli (Gujarat), for providing facilities and laboratories.

#### REFERENCES

- Carla Sousa, Cristina Freire, and Baltazar De Castro, "Synthesis and characterization of benzo-15-crown-5 ethers with appended N2O Schiff bases," Molecules 8 (12), 894-900 (2003).
- Zvjezdana Cimerman, Snežana Miljanić, and Nives Galić, "Schiff bases derived from aminopyridines as spectrofluorimetric analytical reagents," Croatica Chemica Acta 73 (1), 81-95 (2000).
- Hugo Schiff, "Mittheilungen aus dem Universitätslaboratorium in Pisa: eine neue Reihe organischer Basen," Justus Liebigs Annalen der Chemie 131 (1), 118-119 (1864).
- A Xavier and N Srividhya, "Synthesis and study of Schiff base ligands," Journal of Applied Chemistry 7 (11), 06-15 (2014).
- K. D. Bhatt, Disha J Vyas, Bharat A Makwana, Savan M Darjee, Vinod K Jain, and Hemangini Shah, "Turnon fluorescence probe for selective detection of Hg (II) by calixpyrrole hydrazide reduced silver nanoparticle: Application to real water sample," Chinese Chemical Letters 27 (5), 731-737 (2016).
- K. D. Bhatt, Disha J Vyas, Hrishikesh S Gupte, Bharat A Makwana, Savan M Darjee, and Vinod K Jain, "Solid phase extraction, preconcentration and sequential separation of U (VI), Th (IV), La (III) and Ce (III) by Octa-O-methoxy resorcin [4] arene based Amberlite XAD-4 Chelating Resin," World J. Anal. Chem 2 (2), 31-41 (2014).
- K. D. Bhatt, Bharat A Makwana, Disha J Vyas, Divya R Mishra, and Vinod K Jain, "Selective recognition by novel calix system: ICT based chemosensor for metal ions," Journal of Luminescence 146, 450-457 (2014).
- K. D. Bhatt, Hrishikesh S Gupte, Bharat A Makwana, Disha J Vyas, Debdeep Maity, and Vinod K Jain, "Calix receptor edifice; scrupulous turn off fluorescent sensor for Fe (III), Co (II) and Cu (II)," Journal of fluorescence 22 (6), 1493-1500 (2012).
- Hrishikesh S Gupte, K. D. Bhatt, Vinod K Jain, Disha J Vyas, and Bharat A Makwana, "Azo resorcin [4] calixpyrrole grafted Amberlite XAD-2 polymer: an efficient solid phase extractant for separation and preconcentration of La (III) and Ce (III) from natural geological samples," Journal of Inclusion Phenomena and Macrocyclic Chemistry 81 (3-4), 409-422 (2015).
- HZ Kou, ZH Ni, BC Zhou, and RJ Wang, "Synthesis and characterization of Co (II), Ni (II) and Cu (II) complexes containing Heterocyclic Nitrogen donor ligands," Inorg. Chem. Commun 7, 1150 (2004).

- Muhammad Aqeel Ashraf, Karamat Mahmood, Abdul Wajid, Mohd Jamil Maah, and Ismail Yusoff, "Synthesis, characterization and biological activity of Schiff bases," IPCBEE 10, 1-7 (2011).
- Aderoju A Osowole, Ingo Ott, and Oladunni M Ogunlana, "Synthesis, spectroscopic, anticancer, and antimicrobial properties of some metal (II) complexes of (substituted) nitrophenol Schiff base," International Journal of Inorganic Chemistry 2012 (2012).
- Elham Pahlavani, Hadi Kargar, and Nahid Sepehri Rad, "A study on antitubercular and antimicrobial activity of isoniazid Derivative," Zahedan Journal of Research in Medical Sciences 17 (7) (2015).
- Rajeev Johari, Gajendra Kumar, Dharmendra Kumar, and Shailandra Singh, "Synthesis and antibacterial activity of M (II) schiff-base complex," J. Ind. Council Chem 26 (1), 23-27 (2009).
- Aysegul Golcu, Mehmet Tumer, Havva Demirelli, and R Alan Wheatley, "Cd (II) and Cu (II) complexes of polydentate Schiff base ligands: synthesis, characterization, properties and biological activity," Inorganica Chimica Acta 358 (6), 1785-1797 (2005).
- Adhikar D Prakash KA, "Application of schiff bases and their metal complexes-A Review," International Journal of Chem Tech Research 3 (4), 1891-1896 (2011).
- Mohammad Sayed Alam, Jung-Hyun Choi, and Dong-Ung Lee, "Synthesis of novel Schiff base analogues of 4-amino-1, 5-dimethyl-2-phenylpyrazol-3-one and their evaluation for antioxidant and antiinflammatory activity," Bioorganic & medicinal chemistry 20 (13), 4103-4108 (2012).
- Ian Hutchinson, Sharon A Jennings, B Rao Vishnuvajjala, Andrew D Westwell, and Malcolm FG Stevens, "Antitumor benzothiazoles. 16. Synthesis and pharmaceutical properties of antitumor 2-(4aminophenyl) benzothiazole amino acid prodrugs," Journal of medicinal chemistry 45 (3), 744-747 (2002).
- Eiji Kashiyama, Ian Hutchinson, Mei-Sze Chua, Sherman F Stinson, Lawrence R Phillips, Gurmeet Kaur, Edward A Sausville, Tracey D Bradshaw, Andrew D Westwell, and Malcolm FG Stevens, "Antitumor benzothiazoles. 8. Synthesis, metabolic formation, and biological properties of the C-and Noxidation products of antitumor 2-(4-aminophenyl) benzothiazoles," Journal of medicinal chemistry 42 (20), 4172-4184 (1999).
- Yang Zuo, Qiongyou Wu, Sun-wen Su, Cong-wei Niu, Zhen Xi, and Guang-Fu Yang, "Synthesis, herbicidal activity, and QSAR of novel N-benzothiazolylpyrimidine-2, 4-diones as protoporphyrinogen oxidase inhibitors," Journal of agricultural and food chemistry 64 (3), 552-562 (2016).
- 21. Valentin Aleksandrovich Lopyrev, Ludmila Ivanovna Larina, and Tamara I Vakul'skaya, "Quantitative Estimation of the Electronic Effects of Substituents in Five-membered Nitrogen-containing Aromatic



Heterocycles," Russian Chemical Reviews 55 (5), 411-425 (1986).

- R Stojkovic, Grace Karminski-Zamola, Livio Racane, V Tralic-Kulenovic, L Glavas-Obrovac, S Ivankovic, and M Radacic, "Antitumor efficiency of novel fluorosubstituted 6-amino-2-phenylbenzothiazole hydrochloride salts in vitro and in vivo," Methods and findings in experimental and clinical pharmacology 28 (6), 347-354 (2006).
- Valérie Bénéteau, Thierry Besson, Jérôme Guillard, Stéphane Léonce, and Bruno Pfeiffer, "Synthesis and in vitro anti-tumour evaluation of benzothiazole-2carbonitrile derivatives," European journal of medicinal chemistry 34 (12), 1053-1060 (1999).
- 24. MA Basser and NA Mote, "Synthesis and antimicrobial activity of some Schiff bases from benzothiazoles," Asian Journal of Chemistry 13 (2), 496 (2001).
- Chanada S Baluja S, Kulshrestha A., "Synthesis, physical characterization and antibacterial activity of some derivatives of 2-amino benzothiazoles," Acta Chim Pharam Indica 7 (4) (2017).
- FD Karia and PH Parsania, "Biologically active Co (II) and Ni (II) complexes of N-(2-thienylmethylene)-2aminothiadiazole," Asian J Chem 11, 991 (1999).
- HS Hothi, A Makkar, JR Sharma, and MR Manrao, "Synthesis and antifungal 8 potential of Schiff bases of 2'-hydroxyacetophenone and their Cu (II) complexes," Ind. J. Agri. Chemi 41, 53-58 (2008).
- E Pontiki and D Hadjipavlou-Litina, "Synthesis and pharmacochemical evaluation of novel aryl-acetic acid inhibitors of lipoxygenase, antioxidants, and anti-inflammatory agents," Bioorganic & medicinal chemistry 15 (17), 5819-5827 (2007).
- 29. Sheryl J Hays, Michael J Rice, Daniel F Ortwine, Graham Johnson, Roy D Schwarz, Denise K Boyd, Laura F Copeland, Mark G Vartanian, and Peter A Boxer, "Substituted 2-benzothiazolamines as sodium flux inhibitors: Quantitative structure–activity relationships and anticonvulsant activity," Journal of pharmaceutical sciences 83 (10), 1425-1432 (1994).
- 30. BHM Mruthyunjayaswamy and BK Shanthaveerappa, "Synthesis and pharmacological evaluation of 3, 5-

disubstituted indole-2-[N $\beta$ -(substituted benzopyran-2'-one-3'-carboxyl)] carboxy hydrazides and 2H-3-(various substituted indol-3'-yl) methyl-1, 3-benzothiazoles," (2000).

- Christinsonie D. M. and Assunta I. Alain B., "Pharmaceutical compositions containing 2-amino benzothiazole for treating Parkinson's disease and Parkinsonian Syndromes," Patent C A pharmaceuticals 1, 63 (1997).
- Zohreh Salarvand, Mehdi Amirnasr, Milad Talebian, Keyvan Raeissi, and Soraia Meghdadi, "Enhanced corrosion resistance of mild steel in 1 M HCl solution by trace amount of 2-phenyl-benzothiazole derivatives: Experimental, quantum chemical calculations and molecular dynamics (MD) simulation studies," Corrosion Science 114, 133-145 (2017).
- Karema Masoude Abuamer, Abdussalam Ali Maihub, Marei Miloud El-Ajaily, Abdunnaser Mohamed Etorki, Mortaja Mohamed Abou-Krisha, and Majda Albashir Almagani, "The role of aromatic Schiff bases in the dyes techniques," International Journal of Organic Chemistry 4 (01), 7 (2014).
- Marczewski Z. Pilecki M., Gora J., "Schiff bases and their application in perfumery," Kosmetyki 28, 223-227 (1984).
- 35. Shi T. Zhang W., Ding G., et al., "Nano Silica Schiff-Base Co:er(II) complexes with sustainable Antimicrobial Activity against Bacteria and Reduced Risk of Harm to Plant and Environment," ACS Sustainable Chem Eng 5, 502-509 (2017).
- MC Rodriguez, AG Liwes, and DR Frnando, "Acenaphtheneqninom thiosemicarbazones and it is transition metal complexes; synthesis, structure and biological activity," Inorg. Biochem 98 (2004).
- K Lata and A Prakash, "Synthesis and biological evaluation of 4-thiazolidinones and their ketoazomethines," ORIENTAL JOURNAL OF CHEMISTRY 22 (3), 717 (2006).
- Srivastava SK. Saxsena RK., "Synthesis and Antibacterial activity of azomethines and thiazolidinone derivatives of benzimidazoles," J Ind Chem Soc 64, 446-448 (1987).