



Preparation of Terbinafine Hydrochloride Cubosomes: Preformulation Studies

Samuel AJ * and Mulla SA

M.C.E Society's Allana College of Pharmacy, Savitribai Phule Pune University, Pune-411001, Maharashtra, India.

Received: 10 Oct 2018/ Accepted: 8 Nov 2018/ Published online: 01Jan 2019

Corresponding Author Email: aneyjoice2004@yahoo.com

Abstract

Cubosomes are self-assembled liquid crystalline particles that can be used for carrier potential of hydrophilic, lipophilic and amphiphilic drugs as compared to free drug directly to the particular site of action, thus allow drug targeting and the sustained or controlled release of conventional medicines. Terbinafine hydrochloride (TB-HCl) is a synthetic allylamine antifungal agent. The present research was aimed at preformulation characterization of TB-HCl. The drug was characterized for its micromeritic properties, DSC, XRD, FTIR etc. The standard plot indicates high linear correlation between concentration of drug and absorbance. This data further provide a basis for the development of topical cubosomal formulation.

Keywords

Differential Scanning Calorimetry, Micromeritics, Preformulation, Terbinafine Hydrochloride, X-Ray Diffractometry.

INTRODUCTION

The objective of preformulation studies is to develop a portfolio of information about the drug substance to serve as a set of parameters against which detailed formulation design can be carried out. Preformulation investigations are designed to identify those physicochemical properties of drug substances and excipients that may influence the formulation design, method of manufacture, and pharmacokinetic-biopharmaceutical properties of the resulting product [1, 2]. Topical drug delivery is the term used for localized treatment of dermatological condition where the medicine has effects only in specific area, not throughout the body. Cubosomes are discrete, sub-micron, nano-

structured particles of bicontinuous cubic liquid crystalline phase. Cubosomes possess the same microstructure as the parent cubic phase but have larger specific surface area and their dispersions have much lower viscosity in comparison to the bulk cubic phase. Cubosomes are nanoparticles, more accurately nanostructure particles of a liquid crystalline phase with cubic crystallographic symmetry formed by the self-assembly of amphiphilic or surfactant like molecules. However, the cubic phases possess a very high solid like viscosity, which is a unique property because of their intriguing bicontinuous structures which enclose two distinct regions of water separated by a controlled

bilayer of surfactant. As a result, the cubic phases can be fractured and dispersed to form particulate dispersions that are colloidally and/or thermodynamically stable for longer time. Cubosomes have a great potential in drug nanoformulations [3-6].

Cubosomes can be manufactured by two distinct methods [7]:

Top down Technique: It is the most widely used procedure in which bulk cubic phase is first produced and by application of high energy such as high pressure homogenization it is processed into cubosomes nanoparticles.

Bottom up Technique: It is more recently developed technique of cubosome formation, allowing cubosomes to form and crystallize from precursors on the molecular length scale.

Advantages of cubosomes [7]:

- High drug payloads due to high internal surface area and cubic crystalline structures.
- Relatively simple method of preparation
- Biodegradability of lipids
- Capability of encapsulating hydrophilic, hydrophobic and amphiphilic substances.
- Targeted release and controlled release of bioactive agents

Terbinafine is an allylamine which has a broad spectrum of antifungal activity in fungal infections of the hair and skin such as Pityriasis versicolor. It shows oral bioavailability of about 40% because of first pass hepatic metabolism. So Terbinafine is increasingly administered by topical route may increase the bioavailability. Terbinafine hydrochloride is highly lipophilic in nature and tends to accumulate in skin, nails, and fatty tissues. Like other allylamines, terbinafine inhibits ergosterol synthesis by inhibiting the fungal squalene monooxygenase (squalene 2, 3-epoxidase), an enzyme that is part of the fungal cell wall synthesis pathway. So terbinafine prevents conversion of squalene to lanosterol, ergosterol cannot be synthesized. This is thought to change cell membrane permeability, causing fungal cell lysis. It is an anti-fungal which diffuses rapidly into the skin from topical applications and effectively treats athlete's foot i.e. tinea pedis. Terbinafine hydrochloride interferes with the integrity and growth of the fungal cell wall by weakening the cell wall and eventually leading to fungal cell death. It is effective at low levels and diffuses rapidly into the skin to effectively cure tinea. [8, 9, 10].

MATERIALS AND METHODS

Materials:

Terbinafine Hydrochloride was purchased from Yarrow chemicals, Mumbai, India and Ethanol analytical grade was purchased from Himedia Mumbai, India.

Drug Profile [11]

Chemical structure and 3D structure of Terbinafine hydrochloride is shown in figure: 1

[<https://pubchem.ncbi.nlm.nih.gov>]

Physicochemical properties of Drug is given in table: 1

Pre-formulation studies [12-16]

Micromeritic properties

a) Bulk Density:

Bulk Density was determined by pouring the powder into graduated cylinder. The bulk volume (V_b) and weight of powder (M) was recorded and the bulk density was calculated by following formula.

$$\text{Bulk density} = \frac{\text{Mass}}{\text{Bulk volume}}$$

b) Tapped Density:

The cylinder containing powder was tapped for fixed time. The minimum volume (V_t) occupied in the cylinder after tapping and weight (M) of blend was measured. Tapped density was calculated by using formula

$$\text{Tapped density} = \frac{\text{Mass}}{\text{Tapped volume}}$$

c) Compressibility Index:

An indirect method of measuring powder flow from densities was developed by Carr.

Carr's index of the powder was calculated according to equation:

$$\text{Carr's index} = \frac{\text{tapped density} - \text{bulk density}}{\text{tapped density}} \times 100$$

d) Hausner's Ratio:

$$\text{Hausner's ratio} = \frac{\text{tapped density}}{\text{bulk density}}$$

e) Angle of Repose:

Angle of repose was determined by using fixed funnel method. The powder was allowed to flow freely through funnel on to the surface. The diameter of powder heap was measured and angle of repose was calculated by following formula.

$$\theta = \tan^{-1} h/r$$

Where h=height of pile of powder
r= radius of the powder pile

Melting point:

Melting point of TB-HCl was determined with the help of capillary melting point apparatus.

UV visible spectroscopy [17]**a) Determination of λ max**

Accurately weighed amount (10mg) of TB-HCl was transferred in to 10 ml volumetric flask and volume was made up to 10 ml with solvent. From this solution, 1 ml was withdrawn and added to the 10 ml volumetric flask and diluted up to 10 ml with solvent. Finally, sample was scanned in the range of 200-400 nm. The wavelength of the maximum absorption was noted, and UV spectrum was recorded.

b) Calibration Curve of Terbinafine hydrochloride:**Preparation of calibration curve in ethanol:**

Accurately weighed amount (10mg) of TB-HCl was added to 10 ml volumetric flask and the volume was made up to 10 ml with ethanol (1000 μ g/ml). From this solution 1 ml was withdrawn and added into 10 ml volumetric flask and volume was made to 10 ml with ethanol (100 μ g/ml). This solution was used as stock solution. Standard solutions were prepared in the concentration range of 2-10 μ g/ml by suitable dilutions of the stock solution in ethanol and absorbance was taken at 223 nm by UV Spectroscopy (Shimadzu).

Preparation of calibration curve in pH 6.4 buffer:

Accurately weighed amount (10mg) of TB-HCl was added to a volumetric flask and the volume was made up to 10 ml with phosphate buffer (pH 6.4). From this solution 1 ml was withdrawn and added into 10 ml volumetric flask and volume was made to 10 ml with pH 6.4 buffer (100 μ g/ml). This solution was used as stock solution. Standard solutions were prepared in the concentration range of 2-10 μ g/ml by suitable dilutions of the stock solution in pH 6.4 buffer and absorbance was taken at 223 nm by UV Spectroscopy.

FTIR Spectroscopy [18]

FTIR spectrum of drug and excipients sample was recorded for its identification. The spectrum of sample was recorded over the wave number 4000 to 400 cm^{-1} in an FTIR spectrophotometer. The identified peaks were compared with the principle peaks of reported IR spectrum and the sample was identified.

X-Ray Diffractometry (XRD)

The physical nature of the drug was identified by using X-Ray Diffractometry studies. The pure drug sample was analyzed by using Brooker D2 Phaser powder X-Ray Diffractometer.

Differential Scanning Calorimetry (DSC) [19]

Thermogram of pure TB-HCl were taken for DSC study. An empty aluminum pan was used as a reference. DSC measurements were performed at a heating rate of 5 $^{\circ}$ C/min from 50 to 400 $^{\circ}$ C using aluminium sealed pan. The sample size was 4.5 mg of pure drug for measurements. During the measurement, the sample cell was purged with nitrogen gas at 40 ml/min.

RESULTS AND DISCUSSIONS**Micromeritic properties**

The micromeritic properties of the drug powder are given in Table: 2

Melting Point

The melting point of the pure drug Terbinafine hydrochloride was found in the range of 195 $^{\circ}$ C-198 $^{\circ}$ C. The melting point was found to be similar as given in literature. While comparing the melting point of drug with the standard, no significant difference were observed this indicated the purity of the drug.

UV Spectroscopy Analysis**a) Determination of λ max of TB-HCl by UV spectroscopic method**

The maximum absorption value of pure drug, Terbinafine hydrochloride was found to be 223nm wavelength. The observed λ max value of drug was found to be similar as given in literature. Hence the drug was considered to be pure. The UV spectrum of Terbinafine hydrochloride was showed in figure 2.

b) Calibration Curve of Terbinafine hydrochloride in ethanol and in pH 6.4 buffer

Standard graph of TB-HCl in ethanol and phosphate buffer (pH6.4) was plotted. Linearity was observed and good correlation was obtained with R^2 value of 0.9978 & 0.9975 for ethanol & phosphate buffer respectively (table 3 and figure 3, 4)

FTIR Spectroscopy

The FTIR spectrum of the drug, Terbinafine hydrochloride, was obtained using FTIR spectrophotometer. The spectrum is shown in figure 5.

X-Ray Diffractometry (XRD)

The X-Ray diffraction patterns of pure Terbinafine hydrochloride were illustrated in Figure 6. For the crystalline nature of drug powder the sharp intensity peaks were observed around 20 $^{\circ}$, 22 $^{\circ}$ and 25 $^{\circ}$. The characteristic peaks of TB-HCl appeared at a diffraction angle of 24.16 $^{\circ}$ and maximum intensity of 1543 and several sharp diffraction peaks suggesting that the drug is present in crystalline form.

Differential Scanning Calorimetry (DSC)

DSC curves obtained for pure TB-HCl (4.5mg) is shown in figure 7. Pure Terbinafine hydrochloride showed a sharp melting endotherm at (212.59°C).

Table 1: Physicochemical properties of Terbinafine hydrochloride

| | |
|---------------------|--|
| Chemical Name | (2E)-N,6,6-trimethyl-N-(naphthalen-1-ylmethyl)hept-2-en-4-yn-1-amine hydrochloride |
| Chemical Group | Allylamine class |
| CAS Registry Number | 78628-80-5 |
| Molecular Weight | 327.9 |
| Molecular Formula | C ₂₁ H ₂₅ N.HCl |
| Melting Point | 195°C-198°C |
| Description | White Crystalline powder, odourless |
| pKa | 8.94 |
| Solubility | Very slightly or slightly soluble in water, freely soluble in anhydrous ethanol and in methanol, slightly soluble in acetone |

Table 2: Micromeritic properties of the drug

| Sr no. | Parameters | Observations |
|--------|-----------------|--------------|
| 1. | Bulk Density | 0.3 gm/ml. |
| 2. | Tapped Density | 0.508 gm/ml |
| 3. | Carr's index | 41% |
| 4. | Hausner's ratio | 1.695 |
| 5. | Angle Of Repose | 35.1° |

Table 3: Calibration curve of Terbinafine hydrochloride in ethanol

| Concentration (µg/ml) | Absorbance | |
|-----------------------|------------|-----------|
| | In ethanol | In pH 6.4 |
| 2 | 0.1782 | 0.126 |
| 4 | 0.3698 | 0.254 |
| 6 | 0.5198 | 0.398 |
| 8 | 0.7151 | 0.515 |
| 10 | 0.8561 | 0.678 |





Figure1: Chemical structure and 3D structure of Terbinafine hydrochloride




Figure2 : UV spectrum of Terbinafine hydrochloride in ethanol




Figure3: Calibration curve of TB-HCl in ethanol




Figure 4: Calibration curve of TB-HCl in pH 6.4 buffer




Figure 5: IR spectra of TB-HCl




Figure 6: XRD of pure TB-HCl




Figure 7: DSC thermogram of TB-HCl

CONCLUSION

Preformulation studies have played a significant role in anticipating formulation problems and identifying procedures to rectify and develop a safe, effective and stable formulation. This research article provides details of preformulation studies of Terbinafine

hydrochloride which would be helpful to use as a drug candidate.

ACKNOWLEDGEMENT

We are thankful to Principal and Management of Allana College of Pharmacy for providing the laboratory facilities to carry out the studies.

REFERENCES

1. Yadav CS, Lariya N. Preformulation studies of carbamazepine for tablet dosage form. *J Sci Res Pharm.* 2017; 6(1): 1-4.
2. Ansel CH, Popovich GN and Allen VL. *Pharmaceutics dosage form and drug delivery system.* 4th Edition. Lippincott Williams and Wilkins; New York; 2005.
3. Patrick TS, Matthew I.L. Bicontinuous cubic liquid crystalline phase and cubosome personal care delivery systems.
4. Madhurila T, ParuchuriSK, Suriaprabha K. Overview of cubosomes: a nano particle. *International journal of research in pharmacy and chemistry,* 2011; 1(3): 535-541.
5. Spicer P. Cubosome Processing Industrial Nanoparticle Technology Development. *Chemical Engineering Research and Design,* 2005; 83(A11): 1283–1286.
6. Rizwan SB, Dong YD, Boyd BJ, Rades T, Hook S. Characterization of bicontinuous cubic liquid crystalline systems of phytantriol and water using cryo field emission scanning Electron microscopy (cryo FESEM) *Micron,* 2007; 38(5): 478–485.
7. Ramya Sri V, Madhusudhan Reddy A, Karthikeyan R, Srinivasababu P. A review on: Cubosomes drug delivery system. *Indian Journal of Drugs,* 2017, 5(3), 104-108.
8. Balfour JA, Faulds D. Terbinafine. A review of its pharmacodynamic and pharmacokinetic properties, and therapeutic potential in superficial mycoses. *Drugs* 1992; 43(2): 259-84.
9. Ryder NS. Terbinafine: mode of action and properties of the squalene epoxidase inhibition. *Br J Dermatol* 1992; 126 (suppl 39): 2-7.
10. Dollery C, *Therapeutic drugs,* second edition, 2, 1999, T-41.
11. *British Pharmacopoeia,* The department of health, Britain, Published by The Stationery Office on behalf of MHRA, London, 2016; Vol II: 1001.
12. Yadav P, Lariya N. Preformulation studies of hydroquinone for nanocarrier drug delivery system. *J Sci Res Pharm.* 2017; 6(1):5-8.
13. Gupta MK, Swarnkar SK. Preformulation studies of diltiazem hydrochloride from tableted microspheres. *Journal of Drug Delivery and Therapeutics.* 2018; 8(1): 64-69.
14. Khemchand G, Roy SB, Singhvi I. Preformulation study in the development of a tablet formulation for the treatment of ulcerative colitis. *Int. J. Pharm Drug Anal.* 2015; 6(3): 214-222.
15. Brown RJC, Brown RFC. Melting point and molecular symmetry. *Journal of Chemical Education.* 2000;77(6): 724
16. Kumar RP. Physico-chemical characterization, UV spectrophotometric method development and validation studies esomeprazole Magnesium Trihydrate. *J.Chem Pharm.Res.* 2012; 2(3):484.
17. Uddin SA, Rauf A, Kazi TG et al, Highly sensitive spectrometric method for determination of Hydroquinone in skin lightening creams: application in cosmetics. *Int. J Cosmetic Sci* 2010; 1-8.
18. Thermo Nicolet Corporation. *Introduction to Fourier Transform Infrared spectrometry.* A thermo electron business. 2001.
19. Lukas K, Lemaire P, *Differential scanning calorimetry fundamental overview.* *Resonance.* 2009; 14 (8) 807-817.