

COMPARATIVE STUDY OF ACUTE PHASE PROTEINS IN CASE OF ANEMIA OF CHRONIC DISEASE (ACD) AND IRON DEFICIENCY ANEMIA (IDA) AND ITS RELATIONSHIP WITH ERYTHROPOIETIN

Subodh Kumar¹, Dr. U. N. Singh², Mrs. Kiran Saxena³, Ravi Saxena⁴

¹PhD Medical Biochemistry, Assist. Prof. Dept. of Biochemistry, Major S D Singh Medical College & Hospital, Fatehgarh, Farrukhabad (UP), India

²Asso. Prof. Dept. of Biochemistry, Major S D Singh Medical College & Hospital, Fatehgarh, Farrukhabad (UP), India

³Prof. & HOD Biochemistry, Chirayu Medical College & Hospital, Bhopal (MP)

⁴Prof. & HOD Physiology, Chirayu Medical College & Hospital, Bhopal (MP)

*Corresponding Author Email: somya2011.dsk@gmail.com

ABSTRACT

Acute phase proteins play a significant role in anemia of chronic disease and thought to be beneficial in maintain homeostasis. In both the condition, anemia of chronic disease as well as iron deficiency anemia alteration in iron homeostasis takes place. The study includes patients of pulmonary tuberculosis patients (ACD), Iron deficiency anemia (IDA) patients and control subjects. Serum CRP, Serum Haptoglobin (HPT), Serum Transferrin (TRF) and Serum Erythropoietin (EPO) levels were estimated in both the study groups. These values were compared to control subjects. The result of the present study reveals that alteration of acute phase protein takes place in the study groups which leads to disturbance in iron metabolism, resulting anemia. However the changing pattern in Acute Phase Proteins in Anemia of Chronic Disease (ACD) and Iron Deficiency Anemia is different (in ACD, Serum level of CRP, HPT, FRT increases whereas serum level of TRF decreases. But in case of IDA Serum level of CRP and TRF increases whereas serum level of FRT decreases and there was no significant change ($p > 0.05$) observed in Serum Haptoglobin level, when this value was compared with the value of controls).

KEY WORDS

Acute Phase Proteins, Anemia of Chronic disease, Iron deficiency anemia, Erythropoietin, dysregulation of iron metabolism

INTRODUCTION

Acute Phase Proteins are a class of diverse Proteins whose blood plasma concentrations increase (positive acute phase protein), or decreases (negative acute phase protein) during the response to inflammation in the acute phase. They are produced within a few hours by the liver, responding to inflammatory cytokines such as IL-1, TNF- α and in particular IL-6.^{1, 2}
³ The function of positive APPs is important in opsonization and trapping of microorganisms and their products, in activating the complement system, in binding cellular remnants like nuclear fractions, in neutralizing enzymes, scavenging free hemoglobin and radicals, and in modulating the host's immune

response. It has been reported that Acute Phase Protein, hepcidin, synthesized the liver is increased during inflammation which leads to disturbance in iron homeostasis, resulting anemia. The aim of the present study is to compare the abnormalities in Acute Phase Proteins and Erythropoietin in Anemia of Chronic Disease (ACD) and Iron Deficiency Anemia, and to find out what is the probable cause of anemia in such patients?

REVIEW OF LITERATURE

The most common type of anemia observed is Iron-Deficiency Anemia (IDA), followed by thalassemia and anemia of chronic disease (ACD).⁴ Tuberculosis is a

chronic infectious disease, so anemia of inflammation may contribute significantly.⁵ However, anemia of inflammation is considered a major contributor to anemia observed in developing countries.⁶ It has been reported that a small peptide hormone, hepcidin might be involved in the pathogenesis of Anemia of Chronic Disease (ACD).⁷ In addition to its effect on iron metabolism, hepcidin might contribute to EPO resistance by directly inhibiting erythroid-progenitor proliferation and survival.⁸ C - reactive protein (CRP) an acute phase protein, recognizes foreign pathogens and damaged host cells and initiates their elimination.⁹ Other Scientist reported that CRP is a modulator of innate and adaptive immunity. Depending on the circumstances, CRP may enhance or dampen inflammation.¹⁰ Haptoglobin (HPT) are acute phase reactants and bind to free hemoglobin released from destructed red blood cells and form Haptoglobin-Hemoglobin complexes which are removed from circulation by the liver and thus it helps in recycling of iron part of hemoglobin.¹¹ Transferrin- receptor expression is negatively affected by inflammatory cytokines so the level of soluble TfR proceeded at the normal level.¹² It has been reported previously that in the setting of anemia, serum ferritin is the most sensitive lab test for Iron Deficiency Anemia.¹³ Erythropoietin (EPO) is a glycoprotein hormone first purified from human urine.¹⁴ It stimulates the production of Red Blood Cells (RBCs) by stem cell in Bone Marrow. Produced mainly by the kidneys, it is released in response to decreased levels of oxygen in body tissue. Anemia in pulmonary tuberculosis may also occur as a consequence of chronic inflammation, and without apparent loss of blood or bone marrow suppression.¹⁵ Blunted responses of erythropoietin due to release of tumor necrosis factor or other cytokines have been observed.¹⁶

Insufficient endogenous production of erythropoietin (EPO) is probably one of the pathogenic mechanism of the Anemia of Chronic Disease (ACD).¹⁷ In vitro and In vivo evidence suggests that hypoxia and anemia are the most important stimuli of increased EPO

production.¹⁸ Inverse correlation between serum EPO levels and red blood cell mass has been reported earlier.¹⁹

MATERIAL AND METHOD

The present study was conducted in Department of Biochemistry in Shri Aurobindo Institute of Medical Sciences, Indore during March 2008 to March 2010. The study includes 101 subjects in the age groups of 20 to 58 years including 35 control subjects selected from staff and students of SAIMS Medical College, Indore and 30 Patients of Iron deficiency Anemia (IDA) and 36 patients of Anemia of chronic disease (ACD) were selected from the Out Patient Department (OPD) of Medicine in Shri Aurobindo Institute of Medical Sciences, Manorma Raje Tuberculosis Hospital & from different DOTS Centers of Indore. In Anemia of chronic disease patients, only newly diagnosed Pulmonary Tuberculosis Patients falling in DOTS (CAT I) were selected.

Inclusion Criteria for selection of patients (i) Anemic Patients were selected as per WHO criteria (for Male < 13gm/dL & for Female < 12gm/dL) and on the basis of RBC Picture.²⁰ (ii) IDA Patients were selected on the basis of Serum Ferritin level (< 50 ng/mL). Prior to start of study Informed consent was taken from each subject, and then they were enrolled in the study. The Ethical Committee of SAIMS Medical College, Indore approved the study.

Method

Blood samples of all the subjects were analyzed for hemoglobin by automated cell counter (Sysmex KX-21), Serum C- reactive protein by the method of Votila M et al 1981)²¹ serum ferritin by the method of Ronald H et al²² Serum Haptoglobin and Serum Transferrine estimated Turbidimetry by kit method (Giese Diagnostics) and serum erythropoietin (EPO) by the method of Garcia J.F (1982)²³ by using kit acquired from DRG International Inc. USA. Statistical Analysis: Unpaired t- test were used for statistical assessments with SPSS Version 10 to evaluate mean levels of variables in study groups. Values were expressed as Mean $\pm$ SD.

RESULTS & DISCUSSION

Table 1: COMPARISON OF PARAMETERS BETWEEN CONTROL GROUP & ACD

Parameters	Control (35)	ACD (36)	t-value	p-value
CRP ($\mu\text{g/mL}$)	2.95 $\pm$ 0.72	37.99 $\pm$ 7.21	28.58	< 0.001**
HPT (mg/dL)	263.48 $\pm$ 40.25	375.66 $\pm$ 67.46	8.47	< 0.001**
Ferritin (ng/mL)	161.31 $\pm$ 40.2	385.58 $\pm$ 80.08	14.84	< 0.001**
Transferrin (mg/dL)	267.97 $\pm$ 33.34	192.55 $\pm$ 18.1	11.89	< 0.001**
EPO (mU/mL)	25.77 $\pm$ 3.23	40.44 $\pm$ 11.88	7.05	< 0.001**

Values are in Mean $\pm$ SD **highly Significant $p < 0.001$ DF = 69

Table 1. In biochemical parameter the increase in the level of CRP, HPT, Ferritin & EPO were found to be highly significant ($p < 0.001$). Transferring level was decreased and these parameter was statistically highly significant ($p < 0.001$) when these values were compared with control group.

Table 2: COMPARISON OF PARAMETERS BETWEEN CONTROL GROUP & IDA

Parameters	Control (35)	IDA (30)	t-value	p-value
CRP ($\mu\text{g/mL}$)	2.95 $\pm$ 0.72	9.93 $\pm$ 2.87	13.86	< 0.001**
HPT (mg/dL)	263.48 $\pm$ 40.25	261.9 $\pm$ 35.8	0.166	> 0.05
Ferritin (ng/mL)	161.31 $\pm$ 40.2	35.96 $\pm$ 6.95	16.84	< 0.001**
Transferrin (mg/dL)	267.97 $\pm$ 33.34	371.86 $\pm$ 35.64	12.13	< 0.001**
EPO (mU/mL)	25.77 $\pm$ 3.23	61.83 $\pm$ 8.4	23.45	< 0.001**

Values are in Mean $\pm$ SD **highly Significant $p < 0.001$ $p > 0.05$ = Non significant DF = 63

Table 2. In biochemical parameter the increase in the level of CRP, Transferrin, & EPO were found to be highly significant ($p < 0.001$). Ferritin level was decreased and these parameters were statistically highly significant ($p < 0.001$). Significant change in the level of HPT was not recorded ($p > 0.05$) when these values were compared with control group.

Table 3 COMPARISON OF PARAMETER BETWEEN ACD & IDA

Parameters	Control	ACD		IDA	
CRP ($\mu\text{g/mL}$)	2.95 $\pm$ 0.72	37.99 $\pm$ 7.21	↑	9.93 $\pm$ 2.87	↑
HPT (mg/dL)	263.48 $\pm$ 40.25	375.66 $\pm$ 67.46	↑	261.9 $\pm$ 35.8	↓
Ferritin (ng/mL)	161.31 $\pm$ 40.2	385.58 $\pm$ 80.08	↑	35.96 $\pm$ 6.95	↓
Transferrin (mg/dL)	267.97 $\pm$ 33.34	192.55 $\pm$ 18.1	↓	371.86 $\pm$ 35.6	↑
Erythropoietin (mU/mL)	25.77 $\pm$ 3.23	40.44 $\pm$ 11.88	↑	61.83 $\pm$ 8.4	↑

Table 3. The above table showed increased (↑) and decreased (↓) level of biochemical parameters in comparison to control subjects. In both the study groups (ACD & IDA), Erythropoietin level was increased but the level was more raised in IDA subjects.




CHART 1: Comparison of Biochemical Parameter value in Control, ACD and IDA subjects

DISCUSSION

Pulmonary tuberculosis is a chronic infective disease occurring predominantly in socio-economically deprived populations. Therefore, both anaemia of inflammation as well as iron deficiency may coexist in pulmonary tuberculosis. In the present study majority of ACD patients had normocytic normochromic anemia and few cases were of normocytic hypochromic anemia. In Iron deficiency anemia (IDA) patient's subnormal hematological picture is due to low body iron store which influenced Hb synthesis adversely. The blood picture of IDA patients represents hypochromic microcytic pattern. Similar findings were reported earlier.²⁴ In ACD patients a highly significant rise was noted in serum levels of C-reactive proteins (CRP), Haptoglobin (HPT) and Ferritin (FRT) while serum level of transferrin (TRF) was significantly decreased as compared to control group (Chart 1 & Table 1). This finding is consistent with the earlier finding.^{25, 26} In pulmonary tuberculosis ferritin synthesis is stimulated by the inflammatory process.^{27,28,29} Earlier report suggested that certain acute phase proteins: α -2 macroglobulin, AAGP, CRP, Ceruloplasmin and Haptoglobin (HPT) were significantly increased in tuberculosis except transferrin.³⁰ Finding of other scientist suggested that a decrease in concentration of transferrin with severity of disease in pulmonary tuberculosis.³¹ In the present study the change in acute phase protein levels in case of ACD patients might be due to

increase level of proinflammatory cytokine levels during infection. Ferritin synthesis is stimulated by inflammatory process regardless of iron status, serum ferritin, a non invasive indicator of iron store is an acute phase response. In fact this acute phase response is thought to be beneficial to the organism by preventing microbial growth and helping to restore homeostasis. In Anemia of chronic disease (ACD) patients, highly significant rise was noted in serum levels of EPO as compared to control group. The increase in EPO level in ACD patients might be due to increased level of proinflammatory cytokines & acute phase proteins which leads to blunted erythropoietin resistance (Table 1 & Chart 1). Raised level of CRP is a marker of inflammation which causes blunted erythropoietin resistance in Anemia of chronic disease (ACD). The data based on the result of the present and previous study suggest that blunted erythropoietin resistance occurred during infection with rise in acute phase protein level mainly CRP in anemic pulmonary tuberculosis patients.

In Iron deficiency anemia (IDA) patients CRP & TRF were significantly increased, Ferritin (FRT) decreased while change in level of Haptoglobin was non significant ($P > 0.05$). The decreased level of serum Ferritin level, in Iron deficiency anemia (IDA) was due to the depletion of stored iron & poor nutrition (Table 2 & Chart 1). In IDA patients our findings are consistent with the earlier finding.^{32, 12}

In patients of IDA fall in serum iron level causes reduction in ferritin concentration in blood is considered to be a specific indicator of body iron stores.^{33, 34}

Serum TfR measurement is a reliable index of iron depletion and importance in the diagnosis of iron-deficiency anemia.^{32, 12}

Highly significant rise of serum level of EPO was observed in Iron deficiency anemia as compared to control (Table 2). Since Iron deprivation stimulates hypoxia, a most potent stimulus to increases EPO production and is a compensatory mechanism to restore iron in IDA patients. In Iron deficiency anemia (IDA) patients, low serum iron and Hb causes hypoxia that stimulates EPO production. Moreover sub clinical infection in IDA patients increases CRP and other acute phase proteins. This finding is the consistent with earlier study.¹⁷ The report of earlier finding suggested that for any hemoglobin level, the response of erythropoietin was significantly higher in anemic patients with iron deficiency.^{35, 36} The results of the present study suggest that, in both the study groups (ACD & IDA), Erythropoietin level was increased but the level was more raised in IDA subjects, when these values were compared with control. Estimation of Serum Ferritin and Transferrin level is essential for discriminate between ACD with IDA (Table 3).

CONCLUSION

The present study reveals that inflammation and acute phase response interact with iron metabolism. The alteration in Acute Phase Protein during infection is the major leading cause of anemia in such patients. However the changing pattern in Acute phases Proteins in ACD and IDA is variable.

Acknowledgement

I wish to express my deepest gratitude to Dr. Vinod Bhandari (Chairman), Dr.S.S. Bose (former dean) & Dr. Sudhakar Bharti (ethical committee member) SAIMS Medical College, Indore, who provided a healthy environment for study. This work was supported by grants obtained from SAIMS Medical College, Indore, India.

REFERENCES

1. TRE JE et al. The acute phase response and the haematopoietic system: The role of cytokines. Crit Rev Oncol Hemato 1995; 21:1-18.
2. Epstein FH: Acute Phase Proteins and other systemic responses to inflammation. N Engl J Med. 1999; 340:448-454.
3. Acute Phase Reaction and Acute Phase Proteins J. Zhejiang Univ Sci B 2005; Nov 6 (11): 1054-56.
4. Djulbegovic B. Red blood cell Problems. In: Djulbegovic B, ed. Reasoning and Decision Making in Hematology. Edinburgh, UK: Churchill Livingstone; 1992:13.
5. Jurado RI. Iron, infection, and anemia of inflammation. Clin Infect Dis 1997; 25, 888-895.
6. Abshire TC. The anaemia of inflammation: a common cause of childhood anaemia. Pediatr Clinics North America 1996; 43, 623-638.
7. David A. Weinstein, Cindy N. Roy, Mark D.Fleming, Massimo F. Loda, Joseph I.Wolfsdort, and Nancy C. Andrews. Inappropriate expression of hepcidin is associated with iron refractory anemia: implication for the anemia of chronic disease. Blood.2002; 100: 3776-3781.
8. Dallalio G et al. Hepcidin inhibits in vitro erythroid colony formation at reduced erythropoietin concentrations. Blood 2006; 107: 2702-2704.
9. Kilpatrick JM and Volanakis JE Molecular genetics, structure, and function of C-reactive protein. Immunologic Research 1991; 10: 43-53.
10. Du Clos TW and Mold C C-reactive protein: an activator of innate immunity and a modulator of adaptive immunity. Immunologic Research 2004; 30(3): 261-277.
11. Allison A.C, Aprees A.P. The binding of hemoglobin by plasma proteins. Brit Med J. 1957; ii; 1137.
12. Weiss G. Iron and immunity: a doubleedged sword. Eur J Clin Invest 2002; 32: Suppl 1:70-8.
13. Greene MD. "Iron and ADHD" Alan (2004-12-17)
14. Miyake T, Kung CK, Goldwasser E. Purification of human erythropoietin J Biol Chem. 1977; 252 (15): 5558-64
15. Baynes RD, Flax H, Bothwell TH, Bezwoda WR, MacPhail AP, Atkinson P, Lewis D. Haematological and iron-related measurements in active pulmonary tuberculosis. Scand J Haematol 1986; 36: 280-7.
16. Ebrahim O, Folb PI, Robson SC & Jacobs P. Blunted erythropoietin response to anaemia in tuberculosis. Eur J Haematol 1995; 55, 251-254.
17. Igor Theurl, Verena Mattle, Markus Seifert, Mariagabriella Mariani, Christian Marth, and Gunter Weiss. Dysregulated monocyte iron homeostasis and

- erythropoietin formation in patients of chronic disease. *Blood*. 2006;107 (10)
18. Kendall RG: Erythropoietin. *Clin Lab Haematol* 2001, 23:71-80
 19. Spivak, J.L. The Mechanism of Action of Erythropoietin. *Int J Cell Cloning* 1986; 4: 139-166.
 20. Iron deficiency anemia. WHO Tech Rep Ser 1959; 182:4
 21. Votilla M et al. *Immunol Method*, 1981; 42 :11
 22. Ronald H et al, *Clin chem.* 1983; 2916 : 1109 – 1113
 23. Gracia JF, Ebbe SN, Hollander L, Cutting HO, Miller ME, Cronkites EP. Radioimmunoassay of Erythropoietin: Circulating level in Normal and Polycythemic Human Beings. *J Lab Clin Med*. 1982; 99: 624-635.
 24. Cook JD. The measurement of serum transferrin receptor. *Am J Med Sci* 1999; 318(4):269-76.
 25. Rao, Sukhesh, Bernhardt, Vidya. Serum C - reactive protein in Pulmonary Tuberculosis: Correlation with Bacteriological Load and Extent of Disease. *Infectious Diseases in Clinical Practice*: 2009; 17 (5):314-316.
 26. Davood Attaran, Shahrzad M, Mohammad Khajehdaluae, Hossein Ayatollahi, Md. Towhidi Highly sensitive C-reactive protein levels in Iranian patients with pulmonary complication of sulfur mustard poisoning and its correlation with severity of airway diseases *Hum Exp Toxicol* December 2009; 28 (12) : 739-745
 27. Morris CD, Bird AR, Nell H. The haematological and biochemical changes in severe pulmonary tuberculosis. *Q J Med* 1989; 73: 1151-9.
 28. Henrik Friss, Nyagosya Range, Camilla Braendgaard Kristensen, Pernille kaestel. Acute phase response and iron status marker study in Mwanza, Tanzania. *British journal of Nutrition*. 2009; 102: 310- 317.
 29. Subodh Kumar, U. N.Singh, Kiran Saxena, Ravi Saxena. Hematological and Biochemical abnormalities in case of pulmonary tuberculosis patients in Malwa region (Indore) . *Int J Pharm Bio Sci*, 2013; 3 (3) : 237 -241
 30. Grange JM, Kardjito T, Setiabudi I. A study of acute phase reactant proteins in Indonesian patients with pulmonary tuberculosis. *Tubercle* 1984; 65 (1): 23 – 39.
 31. Immanuel C, Acharyulu GS, Kannapiran M, Segaran R, Sarma GR. Acute phase proteins in tuberculous patients. *Indian J Chest Dis Allied Sci* 1990; 32:15–23.
 32. Ferguson BJ, Skikne BS, Simpson KM, Baynes RD, Cook JD. Serum transferrin receptor distinguishes the anemia of chronic disease from iron deficiency anemia. *J.Lab.Clin. Med* 1992; 19:385-90.
 33. Jacobs A, Miller F, Worwood M, Beamish MR, Wardrop CA. Ferritin in the serum of normal subjects and patients with iron deficiency and iron overload. *Br. Med J*. 1972; 4: 206.
 34. Lipschitz DA, Cook JD, Finch CA. A clinical evaluation of serum Ferritin as an index of iron stores. *N Engl J Med* 1974; 290: 1213-1216.
 35. Joosten E, Van Hove L, Lesaffre E, Goosens W, Dereymaeker L, Van Goethem G, Pelemans W. Serum erythropoietin levels in elderly inpatients with the anemia of cancer disorder and iron deficiency anemia. *J Am Geriatr Soc*. 1993; 41 (12):1301-4.
 36. Pamela J kling, Dragsten PR, Roberts RA, Dos Santos B, Brooks, DJ Hedlund BE, Taetle R. Iron deprivation increases erythropoietin production in vitro, in normal subjects and patients with malignancy. *Oncology Digest* 1997; 3:13-15.



***Corresponding Author:**

Subodh Kumar *

Assistant Prof.

Department of Biochemistry,

Major S D Singh Medical College Fatehgarh,

Doctors Residence- G/005 College Campus,

Farrukhabad, UP, India. Pin-209601