

FT-IR STUDIES OF ETHANOLIC EXTRACT OF *IPOMOEA DIGITATA*

Asha Jyothi V* and Satyavati D¹

*Shadan Womens College of Pharmacy, Khairatbad, Hyderabad.

¹Brilliant college of Pharmacy, Koheda, Near Ramoji film city, Hyderabad.

*Corresponding Author Email: ashajyothivadlapudi@gmail.com

ABSTRACT

In the present study an attempt has been made to establish FT-IR profile and identify the functional components of *Ipomoea digitata*. FTIR method was performed on a Thermo Scientific Spectrophotometer system which was used to detect the characteristic peak values and their functional groups. The results of *Ipomoea digitata* roots FTIR analysis confirmed the presence of alkyne, alkane and C-F bonding which shows major peaks at 2360-2312 cm^{-1} , 1616 cm^{-1} , peak between 1456 and 1076 cm^{-1} respectively. The results of the present study produced the FTIR spectrum profile for the medicinally important plant *Ipomoea digitata*.

KEY WORDS

Ipomoea digitata, FTIR, Spectroscopy, Functional groups and Phytoestrogens.

INTRODUCTION

Fourier transform infrared spectrometry is a physico-chemical analytical technique that does not resolve the concentrations of individual metabolites but provides a snapshot of the metabolic composition of a tissue at a given time¹. FTIR can be employed to determine the structure of unknown composition and the intensity of the absorption spectra associated with molecular composition or content of the chemical group^{1, 2}. The FT-IR method measures the vibrations of bonds within chemical functional groups and generates a spectrum that can be regarded as a biochemical or metabolic "fingerprint" of the sample. By attaining IR spectra from plant samples, it might possible to detect the minor changes of primary and secondary metabolites^{2, 3}. At present, particularly in phytochemistry, FT-IR has been exercised to identify the concrete structure of certain plant secondary metabolites⁴⁻⁶. But, on pharmacognosy front FT-IR is still a novel tool to characterize and identify the commercial components from the adulterant. FT-IR method has been successfully utilized in the characterization of bacterial, fungal and plant species⁷⁻¹⁹. FT-IR is one of the most widely used methods to identify the chemical constituents and elucidate the compounds structures, and has been used as a requisite method to identify medicines in Pharmacopoeia of many countries¹². At the pharmacological front FT-IR is helpful tool to identify

the compounds responsible for respective pharmacological activities.

Ipomoea digitata popularly called Vidarikanda is the dried tuber belonging to family *Convolvaceae*, a large, perennial climber with tuberous roots, upto 60 cm long and 30 cm thick, even weighing upto 35 kg, from about 5 or 10 kg; they are distributed nearly throughout India. *Ipomoea digitata* has various pharmacological properties like Spasmogenic effect, Revitalizing effect, Antioxidant activity, Antidiabetic activity, Hepatoprotective activity, Cardioprotective effect and Antihypertensive activity²¹.

MATERIALS AND METHODS

Collection and processing of plant material:

Ipomoea digitata was procured from the authorized botanist Dr. Madhukar Reddy of Heritage bionaturals, Habsiguda, Hyderabad. Shade dried samples were grounded to fine powder using pulverizer. The powdered samples were then stored in a refrigerator for further use.

Extraction of plant material:

The powdered bulbil of *Ipomoea digitata* were extracted using ethanol with gentle stirring for 72 h separately at room temperature. The extracts were then filtered through Whatmann No. 1 filter paper and concentrated using rotaevaporator.

FTIR Spectroscopic Analysis:

The FT-IR spectrophotometer used was Shimadzu at Osmania University. KBr is an important sample matrix for FTIR scanning. The KBr used was of IR grade (SD Fines). About 500 mg of KBr was placed into a mortar and grind it until there is no evidence of crystallinity. The KBr powder was transferred into the drying box at a temperature of 40°C. 10 mg of solid sample was placed into the mortar and again grind it until a fine powder is formed. Weigh 1mg of solid fine powder of sample (as per requirement of the die) and 200mg of dry fine powder of KBr. Weighed quantities were transferred into a mortar and mix well with the help of a spatula. Bottom and top portion of KBr were assembled at press assembly and one of the 13 mm die with the polished surface up inside the press. The KBr sample mixture was transferred to KBr press assembly. Second die was transferred inside the KBr press assembly with polished side down so that KBr sample mixture was sandwiched between the polished surfaces of the each die. The KBr was transferred to press assembly to press. Vacuum line

was connected to evacuate air from the KBr press assembly with a vacuum pump. The die is slowly compressed in KBr press assembly until a pressure of 2000 kg/cm² is achieved on gauge with the vacuum on. Making sure that pressure release valve is closed. After 60 s, slowly the pressure release valve is open to release the pressure and also the vacuum line is disconnected. The disc is checked if it is translucent and the sample is homogeneously distributed in the disc²⁰. The prepared disc is then subjected for scanning between 500-4000⁻¹ cm.

RESULTS AND DISCUSSION

The FTIR spectrum was used to identify the functional group of the active components based on the peak value in the region of infrared radiation. The bulbs powder and ethanolic extracts evaporated powder of *Ipomoea digitata* was passed into the FTIR and the functional groups of the components were separated based on its peak ratio.




Fig 1: Showing the peaks of FT-IR scan of ethanolic extract *Ipomoea digitata*.

The results of *Ipomoea digitata* roots FTIR analysis confirmed the presence of alkyne, alkane and C-F bonding which shows major peaks at 2360-2312 cm⁻¹, 1616 cm⁻¹, peak between 1456 and 1076 cm⁻¹ respectively. While the peak at 2090-1992 cm⁻¹ was observed was unidentified for any organic constituents. The results of the present study produced the FTIR spectrum profile for the medicinally important plant *Ipomoea digitata*.

CONCLUSION

Many researchers applied the FTIR spectrum as a tool for distinguishing closely associated plants and other organisms⁽¹⁻¹⁹⁾. The results of the present study coincided with the previous observations observed by various plant biologist and taxonomist. The results of the present study developed novel phytochemical marker to identify the medicinally important plant. Further advanced spectroscopic studies are required

for the structural elucidation and identification of active principles present in the leaves of *Ipomoea digitata*.

REFERENCES

1. Griffiths PR, de Haseth JA. Fourier transform infrared spectroscopy. New York: Wiley, 1986.
2. Surewicz WK, Mantsch HH, Chapman D. Determination of Protein Secondary Structure by Fourier Transform Infrared Spectroscopy: A Critical Assessment. *Biochemistry* 1993; 32(2): 389-393.
3. McCann MC, Hammouri M, Wilson R, Belton P, Roberts K. Fourier transform infrared microspectroscopy is a new way to look at plant cell walls. *Plant Physiol.* 1992; 100: 1940- 1947.
4. Yang J, Yen HCE. Early Salt Stress Effects on the Changes in Chemical Composition in Leaves of Ice Plant and Arabidopsis. A Fourier Transform Infrared Spectroscopy Study. *Plant Physiology* 2002; 130: 1032-1042.
5. Stehfest K, Toepel J, Wilhelm C. The application of micro-FTIR spectroscopy to analyze nutrient stress-related changes in biomass composition of phytoplankton algae. *Plant Physiology and Biochemistry* 2005; 43: 717-726.
6. Ivanova DG, Singh BR. Nondestructive FTIR monitoring of leaf senescence and elicitin-induced changes in plant leaves. *Biopolymers* 2003; 72(2): 79-85.
7. Helm D, Labischinski H, Schallehn G, Naumann D. Classification and identification of bacteria by Fourier transform infrared spectroscopy. *J. Gen. Microbiol.* 1991; 137:69-79.
8. Timmins EM, Howell SA, Alsberg BK, Noble WC, Goodacre R. Rapid differentiation of closely related *Candida* species and strains by pyrolysis-mass spectrometry and Fourier transform-infrared spectroscopy. *J. Clin. Microbiol.* 1998; 36: 367-374.
9. Goodacre R, Shann B, Gilbert RJ, Timmins EM, McGovern AC, Alsberg BK, Kell DB, Logan NA. Detection of the dipicolinic acid biomarker in *Bacillus* spores using Curie-point pyrolysis mass spectrometry and Fourier transform-infrared spectroscopy. *Anal. Chem.* 2000; 72:119-127.
10. Johnson HE, Broadhurst D, Goodacre R, Smith AR. Metabolic fingerprinting of salt-stressed tomatoes. *Phytochemistry* 2003; 62:919-928.
11. Martín JA, Solla A, Woodward S, Gil L. Fourier transform-infrared spectroscopy as a new method for evaluating host resistance in the Dutch elm disease complex. *Tree Physiology* 2005; 25, 1331-1338.
12. Liu H, Sun S, Lv G, Chan KKC. Study on Angelica and its different extracts by Fourier transform infrared spectroscopy and two-dimensional correlation IR spectroscopy. *Spectrochimica Acta Part A* 2006; 64: 321-326.
13. Duraes N, Bobos I, Ferreira Da Silva E. Chemistry and FT-IR spectroscopic studies of plants from contaminated mining sites in the Iberian Pyrite Belt, Portugal. *Mineralogical Magazine* 2008; 72(1): 405-409.
14. Wei ZL, Dong L, Tian ZH. Fourier transform infrared spectrometry study on early stage of cadmium stress in clover leaves. *Pak. J. Bot.* 2009; 41(4): 1743-1750.
15. Thenmozhi M, Bhavya PK, Rajeshwari Sivaraj. Compounds identification using HPLC and FTIR in *Eclipta alba* and *Emilia sonchifolia*. *International Journal of Engineering Science and Technology* 2011; 3(1):292-298.
16. Janakiraman N, Sahaya Sathish S, Johnson M. UV-VIS and FTIR Spectroscopic Studies on *Peristrophe bicalyculata* (Retz.) Nees. *Asian Journal of Pharmaceutical and Clinical Research* 2011; 4(4): 125-129.
17. Dorado J, Almendros G, Field JA, Sierra-Álvarez R. Infrared spectroscopy analysis of hemp (*Cannabis sativa*) after selective delignification by Bjerkandera sp. at different nitrogen levels. *Enzyme Microb. Technol.* 2001; 28:550-559.
18. Gidman E, Goodacre R, Emmett B, Smith AR, Jones DG. Investigating plant-plant interference by metabolic finger printing. *Phytochemistry* 2003; 63: 705-710.
19. Hori R, Sugiyama J. A combined FT-IR microscopy and principal component analysis on softwood cell walls. *Carbohydr. Polym.* 2003; 52: 449-453.
20. Kumar S, Joshi P, Rajvanshi A, Scanning Of Forensic Samples on Ftir – Sample Preparation, The.Internet.J.For.Sci. 2008. (4)1:1-4.
21. Jyothi. VA, Pippalla. S.R, Satyavathi.D. *Ipomoea digitata* – An Update, IJPRR, Jan 2013, Volume 1, Issue 1, PP 13-21.

***Corresponding Author:**

Asha Jyothi V* & Satyavathi D

Email: ashajyothivadlapudi@gmail.com