



Guillain-Barré Syndrome as a Neurological Disorder and Its Association with Post-COVID-19 Vaccination: A Review

Aditya Singh¹, Saief Ansari¹, Neha Sharma², Arushi Purva², Harshit Gautam², and Sandeep Singh²

¹University Institute of Pharmaceutical Education and Research (UIPER), University of Kota, Rajasthan

²Assistant Professor, University Institute of Pharmaceutical Education and Research (UIPER), University of Kota, Rajasthan

Received: 20 Feb 2026 / Accepted: 25 Mar 2026 / Published online: 01 April 2026

*Corresponding Author Email: sharmaneha2706@gmail.com

Abstract

Guillain-Barré syndrome (GBS) is a rare autoimmune neurological disorder affecting the peripheral nervous system. It is commonly triggered by infections and has also been reported rarely after vaccination. During the COVID-19 pandemic, cases of GBS following COVID-19 vaccination were reported, raising safety concerns. This review summarizes recent literature (2016–2025) on the pathophysiology, clinical features, diagnosis, and management of GBS, with special importance on post-COVID-19 vaccination cases. Evidence suggests a temporal association, particularly with adenovirus-based vaccines; however, the overall incidence remains very low. Early diagnosis and rapid treatment result in favourable outcomes. The benefits of COVID-19 vaccination overshadow the rare risk of neurological complications.

Keywords

Guillain-Barré syndrome, COVID-19 vaccine, neurological disorder, autoimmune disease.

1. INTRODUCTION

Guillain Barre Syndrome (GBS) is an acute disorder in which immune system attacks body's own tissues leading to inflammation and damage to peripheral nervous system (PNS) usually triggered by an infection or sometimes vaccinations. It is also a most common cause of acute flaccid paralysis (sudden weakness of all four limbs) worldwide. This syndrome was first referred over 100 years ago by Guillain, Barre, and strohl, who identified the characteristic finding of albumin cytological dissociation in cerebrospinal fluid (CSF). It affects about 1-2 people per 100,000 per year. More common in men as compared to women and risk increases with age, but occurs at any age. GBS is now

considered a miscellaneous group of related disorders, including Miller Fisher syndrome (MFS) and several regional variants. GBS shows a wide range of clinical patterns, which involves mild nerves up to severe paralysis and cause problems in respiratory functions.

The coronavirus disease 2019 (COVID-19) pandemic has posed a unique challenge to global public health system, resulting in many diseases, deaths and socio-economic disturbance worldwide. Since its arrival in late 2019, severe acute respiratory syndrome coronavirus-2 (SARS-COV-2) has led to repeated waves of infection, massive overload on healthcare infrastructures and demanding the rapid development of preventive strategies. Among these,

vaccination is the most effective method to reduce severe illness and COVID-19 related deaths. Vaccination campaigns were later implemented across both high income and low to-middle-income countries on large scale, leading to the administration of billions of vaccine doses worldwide.

Recent research suggests a possible temporal association between COVID-19 vaccination and onset of GBS in some individuals, but causality remains unproven and the overall risk is low compared to the public health benefits of vaccination.

2. EPIDEMIOLOGY PARAMETER

Description

Global incidence 1–2 per 100,000 persons/year

Gender More common in males

Vaccine association Rare, mostly adenovirus-vector vaccines
Onset time 7–21 days post-vaccination

3. PATHOPHYSIOLOGY

GBS is autoimmune in nature. Molecular mimicry — where immune cells mistake nerve components for vaccine or viral proteins — may play a role. In COVID-19 vaccination, especially with adenovirus-based vaccines, the immune response may sometimes trigger antibodies that attack nerve tissue.

Main mechanisms discussed in literature:

- Autoimmune attack on myelin/axon of peripheral nerves
- Possible involvement of pro-inflammatory cytokine response
- Vaccination may trigger immune activation in rare sensitive individuals




Figure 1: Immunopathogenesis of Guillain-Barre Syndrome

4. Clinical Features

- Ascending muscle weakness
- Loss of deep tendon reflexes
- Sensory disturbances
- Facial nerve involvement
- Autonomic dysfunction in severe cases
- Paresthesia (Numbness)




Figure 2: Clinical Course of Guillain-Barre Syndrome

5. DIAGNOSIS

Diagnosis is based on: -

- **Clinical Symptoms**
- **Nerve conduction Studies**
- **CSF analysis:** shows albumin cytologic dissociation (high protein, normal cells)

6. TREATMENT AND OUTCOME

Treatment Effect IV Immunoglobulin (IVIg) Most commonly used

Plasmapheresis: Effective in severe cases

Physical therapy: Important for recovery
Most patients respond well if treated early, but recovery varies by case severity.




Figure 3: Timeline of Guillain-Barre Syndrome following COVID-19 vaccination

7. GBS FOLLOWING COVID-19 VACCINATION


Although rare, several reviews and systematic analyses reported GBS cases after COVID-19 vaccination:

- **AIDP (Acute Inflammatory Demyelinating Polyneuropathy) subtype** was the most common form post-vaccination.
- Multiple case series reported onset typically within ~14 days after vaccination.

- It appears to be **higher association with adenovirus-vector vaccines** than with mRNA vaccines.

Post-vaccination GBS is rare and usually presents within two weeks of vaccination. Most reported cases respond well to standard treatment.

Graph constructed based on data reported by Hanson et al. (2022)



Flowchart 1: Selection process of studies included in the review

8. CONCLUSION

Guillain-Barré syndrome is a rare autoimmune neurological disease that can follow infections and, very rarely, vaccinations. While some cases have been reported after COVID-19 vaccines, the overall

risk remains extremely low and significantly less than the risk of neurological complications from natural SARS-CoV-2 infection. Early recognition, diagnosis, and appropriate immunotherapy can lead to good recovery in most patients. Continued monitoring and

research are essential to improve understanding and vaccine safety.

Institutional Review Board Statement

Not applicable

Informed Consent Statement

Not applicable

Funding

Not applicable

Acknowledgments

Thanks to University institute of pharmaceutical education and research institute (UIPER), University of kota, Kota for providing favorable environment to review this article.

Conflicts of Interest

The author declares no conflict of interest.

REFERENCES

- Willison HJ, Jacobs BC, van Doorn PA. Guillain-Barré syndrome. *Lancet*. 2016;388(10045):717-727.
- Finsterer J. Neurological side effects of SARS-CoV-2 vaccinations. *Acta Neurol Scand*. 2022;145(1):5-9.
- Abu-Rumeileh S, Abdelhak A, Foschi M, Tumani H, Otto M. Guillain-Barré syndrome spectrum associated with COVID-19: an up-to-date systematic review. *Neurol Sci*. 2021;42(11):4397-4407.
- Shahrizaila N, Lehmann HC, Kuwabara S. Guillain-Barré syndrome. *Lancet*. 2021;397(10280):1214-1228.
- Patone M, Handunnetthi L, Saatci D, et al. Guillain-Barré syndrome following COVID-19 vaccination. *Nat Med*. 2021;27(12):2145-2153.
- Keh RYS, Scanlon S, Datta-Nemdharry P, et al. COVID-19 vaccination and Guillain-Barré syndrome: analyses using the National Immunoglobulin Database. *Brain*. 2022;145(2):739-748.
- Maramattom BV, Krishnan P, Paul R, et al. Guillain-Barré syndrome following ChAdOx1-S/nCoV-19 vaccine. *Neurol India*. 2021;69(5):13021304.
- Luijten LWG, van Kammen MS, van Oijen MGH, et al. Guillain-Barré syndrome after COVID-19 vaccination: a systematic review. *J Neurol*. 2023;270(5):2581-2594.
- Keddie S, Pakpoor J, Mousele C, et al. Epidemiological and cohort study finds no association between COVID-19 and Guillain-Barré syndrome. *Brain*. 2021;144(2):682-693.
- Leonhard SE, Mandarakas MR, Gondim FAA, et al. Diagnosis and management of Guillain-Barré syndrome in ten steps. *Nat Rev Neurol*. 2019;15(11):671-683.
- Sejvar JJ, Kohl KS, Gidudu J, et al. Guillain-Barré syndrome and Fisher syndrome: case definitions for immunization safety data. *Vaccine*. 2019;37(42):6161-6173.
- Hanson KE, Goddard K, Lewis N, et al. Incidence of Guillain-Barré syndrome after COVID-19 vaccination. *JAMA Netw Open*. 2022;5(4):e228879.
- Li X, Ostropolets A, Makadia R, et al. Incidence of Guillain-Barré syndrome after COVID-19 vaccination. *Clin Infect Dis*. 2022;75(7):e1030e1036.
- Woo EJ, Mba-Jonas A, Dimova RB, et al. Guillain-Barré syndrome after Ad26.COV2.S vaccination. *JAMA*. 2021;326(16):1606-1613.
- Shao SC, Wang CH, Chang KC, et al. Guillain-Barré syndrome associated with COVID-19 vaccination. *Front Immunol*. 2021;12:739215.
- Kim JE, Min YG, Shin JY, et al. Guillain-Barré syndrome and COVID-19 vaccination: a nationwide cohort study. *Eur J Neurol*. 2023;30(4):11251133.
- Fragiel M, Miró Ò, Llorens P, et al. Guillain-Barré syndrome in COVID-19 patients. *Clin Microbiol Infect*. 2021;27(10):1440-1446.
- Toscano G, Palmerini F, Ravaglia S, et al. Guillain-Barré syndrome associated with SARS-CoV-2. *N Engl J Med*. 2020;382(26):2574-2576.
- Dalakas MC. Guillain-Barré syndrome: COVID-19-triggered autoimmune neurologic disease. *Neurol Neuroimmunol Neuroinflamm*. 2020;7(5):e781.
- Lunn MP, Cornblath DR, Jacobs BC, et al. COVID-19 vaccine and Guillain-Barré syndrome. *Brain*. 2021;144(2):357-360.
- Allen CM, Ramsamy S, Tarr AW, et al. Guillain-Barré syndrome variant after SARS-CoV-2 vaccination. *Ann Neurol*. 2021;90(2):315-318.
- Hippisley-Cox J, Patone M, Mei XW, et al. Neurological complications after COVID-19 vaccination. *BMJ*. 2021;374:n1786.
- Andrews N, Stowe J, Ramsay M, et al. Risk of Guillain-Barré syndrome after COVID-19 vaccination. *Lancet Infect Dis*. 2022;22(5):623-631.
- Waheed S, Bayas A, Hindi F, et al. Guillain-Barré syndrome after Pfizer vaccination. *Cureus*. 2021;13(2):e13426.
- Razok A, Shams A, Almeer A, et al. post-COVID-19 vaccine Guillain-Barré syndrome. *IDCases*. 2021;24:e01187.
- van den Berg B, Walgaard C, Drenthen J, et al. Guillain-Barré syndrome: pathogenesis and treatment. *Nat Rev Neurol*. 2014;10(8):469-482.
- Nagy A, Alhatlani B. Molecular mimicry in autoimmune disease. *Int J Mol Sci*. 2020;21(21):8477.
- Lucchese G, Flöel A. Molecular mimicry between SARS-CoV-2 and human proteins. *Autoimmun Rev*. 2020;19(8):102548.
- Kaulen LD, Doubrovinskaia S, Mooshage C, et al. Autoimmune neurological diseases following vaccination. *Front Neurol*. 2022;13:832558.
- Finsterer J. Guillain-Barré syndrome after SARS-CoV-2 vaccination: a systematic review. *J Neurol Sci*. 2022;440:120340.
- Shasha D, Bareket R, Sikron FH, et al. Real-world safety of BNT162b2 vaccine. *Clin Microbiol Infect*. 2022;28(10):1533.e1-1533.e6.
- García-Grimshaw M, Michel-Chávez A, Vera-Zertuche JM, et al. Guillain-Barré syndrome following

- COVID-19 vaccination. *Neurology*. 2021;97(24):e2376-e2383.
33. Márquez Loza AM, Holroyd KB, Johnson SA, et al. Guillain-Barré syndrome in the COVID-19 era. *Neurol Clin Pract*. 2021;11(2):e142-e147.
34. Keh RYS, Scanlon S, Datta-Nemdharry P, et al. Temporal association of Guillain-Barré syndrome with COVID-19 vaccination. *Brain*. 2022;145(2):739-748.
35. Woo EJ. An overview of Guillain-Barré syndrome and vaccination. *Vaccine*. 2023;41(5):1043-1050.