

EVALUATION OF HEPATOPROTECTIVE ACTIVITY OF ETHANOLIC EXTRACT OF ACACIA CATECHU WILD IN PARACETEMOL INDUCED HEPATOTOXICITY IN ALBINO RATS

Sheshidhar G Bannale¹, Yasmeen A Maniyar², Arati Chikaraddy³, Sangappa V kashinakunti⁴, Pundarikaksha HP⁵, Vijay Dumble⁶, Manjula R⁷

^{1,2,3} Department of Pharmacology S Nijalingappa Medical College Bagalkot 587102

⁴ Department of Biochemistry S Nijalingappa Medical College Bagalkot 587102

⁵ Department of Pharmacology Kempegowda institute of medical sciences Bangalore 560070

⁶ Department of Pathology S Nijalingappa Medical College Bagalkot 587102

⁷ Department of Community medicine S Nijalingappa Medical College Bagalkot 587102

*Corresponding Author Email: drshashibannale@yahoo.co.in

ABSTRACT

Introduction: *Acacia catechu* is a deciduous, thorny tree which grows up to 12-15 meter in height. In Hindi is called as Khair and in kannada Kachu. It has been used as used for the treatment of inflammation, diarrhoea, fever.

Aims: to evaluate the hepatoprotective effect of ethanolic extract of acacia catechu in paracetamol induced liver damage in experimental animals, rats was studied. **Methodology:** The ethanolic extract of acacia catechu was studied for its hepatoprotective effect on paracetamol induced liver damage on Wistar albino rats. The degree of hepatoprotection was measured by biochemical (serum gultamic pyruvic transaminase, serum gultamic oxaloacetic transaminase, alkaline phosphatase) markers and histological changes. Silymarin was used as positive control. **Results:** Pretreatment with acacia extract significantly prevented the biochemical enzyme levels and histological changes induced by paracetamol in the liver. The effects of acacia catechu were comparable to that of the standard drug silymarin. Thus ethanol extract of acacia was found to exhibit marked hepatoprotective activity ($P < 0.05$). **Discussion and conclusion:** These results indicate that acacia catechu could be useful in preventing chemically induced acute liver injury. Hence from this study it can be concluded that ethanol extract of acacia catechu possess significant hepatoprotective activity.

KEY WORDS

Acacia catechu, hepatotoxicity, paracetamol, silymarin

INTRODUCTION

Liver disorders constitute serious health problems in India. Currently there are various drugs of allopathic medicine are available which show significant hepatoprotective action, yet plant products/extracts play a vital role in this scenario. India has a rich source of traditional medicines, many of which are of plant origin.

Liver disorders are mainly inflammatory in nature i.e. acute, sub-acute and chronic hepatitis of various aetiologies like viral infection- hepatitis A, B, C, E, chronic alcohol consumption, toxin induced, drug

induced and malignancy associated disorders. Liver disorders can vary from asymptomatic elevation of biochemical markers to symptomatic presentation like jaundice, pain abdomen, vomiting, fatigue and even go for severe organ failure and coma [1].

India is being rich source of traditional medicines, many of which are plant origin. Herbal drugs have gained importance and popularity in recent years because of their safety, efficacy and cost effectiveness. There are more than one hundred compounds of plant origin, which have shown hepatoprotective potential.

In this study, we evaluated the hepatoprotective action of acacia catechu wild, in paracetamol induced hepatitis in albino rats. Pale catechu or katha (in Hindi) or cutch is an important product of acacia catechu (leguminosae). Catechu the extract prepared from the acacia catechu wood has been used for treating fever, diarrhoea, leucorrhoea, piles, erysipelas. It has also shown hypoglycaemic activity [2].

The juice from the fresh bark has been used in treatment of haemoptysis and gonorrhoea. It also has expectorant activity [3]. Recently its antibacterial action has been evaluated [4].

Catechu contains – catechuic acid, catechu tannic acid (25-33%), acacatechin (10-12%), catechu red, quercetin, catechin (2-12%), epicatechin, phlebotanin (25-33%), gummy matter, quercitrin, quercetin and moisture. Quercitrin is a phenolic flavonoid and catechu of acacia is a pseudotannin. Catechin and epicatechin usually accompany other flavonoids [2,3]. Considering the above facts, the present study was undertaken to evaluate the hepatoprotective property of acacia catechu in experimental animal model i.e. paracetamol induced hepatotoxicity in albino rats.

MATERIALS AND METHODS

Collection and Preparation of extract

Catechu (550 gm) was collected from the local market during the month of August 2012 and differentiated from uncaria gambier by fluorescein test. The plant was identified by botanist and voucher specimen was numbered and deposited in the Herbarium of Department of Pharmacology of S. Nijalingappa medical college, Bagalkot, Karnataka.

The ethyl extract of acacia catechu obtained by making powder of catechu with the help of grinder. Pale coloured powder is obtained, which is deflated with ether to remove the non-polar and phenolic compounds. Then catechu was extracted with 95% ethanol. Then air dried the extract for 48 hours and it was again extracted with ethanol to have concentrated extract.

Acute toxicity Study

The acute toxicity study of heartwood powder of *Acacia catechu* was carried out on Swiss mice with a

dose of 2, 4 and 6 g/Kg body weight orally. A single dose of *Acacia catechu* was administered in the form of slurry (water as vehicle). The animals were observed for changes in parameters like body weight, food and water intake and were reported. There is no sign of toxicity. There was no mortality recorded even at the highest dose level of 6gm/kg body weight. [14-16].

Phytochemical analysis

The extract material was subjected for phytochemical analysis. It showed the presence of alkaloids, tannins, proteins, carbohydrates, resins were present.

Study drugs

Paracetamol, Silymarin, acacia catechu extract were used in this study.

METHODOLOGY

Study animals

24 albino wistar rats of either gender weighing 100-150gm were selected from the central animal house of S Nijalingappa Medical College, Bagalkot, Karnataka, India. The experimental protocol was approved by the institutional animal ethics committee and by the animal regulatory body of the Indian Government. (Registration No: 829/AC/04/CPCSEA). No protocol related procedures were undertaken before taking animal ethics committee approval. All procedures carried out in the study were as per CPCEA guidelines. Selected animals were examined and screened for general health condition including vital parameters. All the animals were acclimatized to laboratory for one week before starting the study. The animals were housed under standard laboratory conditions maintained at $25 \pm 5^{\circ}\text{C}$ and exposed to 12 hr dark and 12 hr light cycle and fed with standard pellet diet and water *ad libitum*.

Induction of hepatotoxicity

These animals were grouped randomly into 4 groups i.e. 6 animals in each group randomly (n=6).

Groups include group –I: Vehicular control group received water (5ml/kg p.o.) for 7 days served as control group

Group-II: Paracetamol group received paracetamol (250mg/kg p.o.) daily for 7 days along with water (5ml/kg p.o.)

Group-III: Silymarin group received standard drug Silymarin (25mg/kg p.o.) daily for 7 days along with paracetamol (250mg/kg p.o.) daily.

Group-IV: Test group received acacia extract (250mg/kg p.o.) daily for 7 days along with paracetamol (250mg/kg p.o.) daily.

On the 7th day of study, blood was drawn from all animals in separate containers for analysis. Laboratory values i.e. SGOT, SGPT and serum alkaline phosphatase (ALP) of all animals is obtained and tabulated.

Histopathological study

Then animals were sacrificed and a portion of hepatic tissue is obtained and washed with normal saline. Then these liver tissues were fixed in 10% buffered neutral formalin for 48 hours. Then thin 5mm sections were made by using microtome and stained with haematoxylin and eosin. Then evaluation was done under light microscope.

Statistical analysis:

Data were analysed using the software SPSS version 19.0 to calculate standard deviation, mean and standard error of mean. Comparison between the groups was done using one way ANNOVA with post hoc Turkey's multiple comparison test using SPSS

version 19 software. Difference between groups was considered to be significant if Null hypothesis rejected > 95% confidence interval ($p < 0.05$).

RESULTS

Table 1 depicts the changes in biochemical parameters in the control and various experimental groups. Group I shows values within physiological range for parameters like SGOT, SGPT and ALP. Whereas there was a significant raise in these values in group II who received paracetamol indicating hepatic injury caused by paracetamol. Group III showing reduction in these values because of the standard control i.e. Silymarin. Group IV which consists of, acacia extract shows decrease in mean values comparable to that of group III.

Table 2 shows comparison among various groups using ANNOVA and post hoc Turkey test. There was significant difference between control group and paracetamol group showing raise in lab values was statistically significant. Silymarin control group (III) values and acacia extract group (IV) were comparable statistically, there was no significant difference.

Table: 1 Effect of acacia catechu extract on paracetamol induced hepatotoxicity in rats

Groups	SGOT	SGPT	ALP
I (Vehicle control)	171.17±01.2	70.92± 00.76	177.85±2.09
II (Paracetamol control)	284.23±11.82	229.57±06.61	202.40±3.18
III (Silymarin +Paracetamol)	222.85±16.38	127.23±15.71	188.30±5.06
IV(Acacia extract + Paracetamol)	223.32±04.94	128.53±00.72	190.07±1.00

Values are mean ± SEM; n= 06 in each group p was considered significant 95% CI

Table 2 depicting comparison between groups with p value* = <0.05

			ALP	SGPT	SGOT
		Group	P	p	P
Tukey HSD	Control (I)	paracetamol	0.000*	0.000*	0.000*
		silymarine and paracetamol	0.130*	0.001*	0.011
		Acacia extract and paracetamol	0.062	0.001*	0.010*
	Paracetamol (II)	control	0.000	0.000	0.000
		silymarine and paracetamol	0.026*	0.000	0.002*
		Acacia extract and paracetamol	0.058	0.000	0.003*
	silymarine and paracetamol (III)	control	0.130	0.001*	0.011*
		paracetamol	0.026*	0.000	0.002*
		Acacia extract and paracetamol	0.979	1.000	1.000
	Acacia extract and paracetamol (IV)	control	0.062	0.001*	0.01*
		paracetamol	0.058	0.000	0.003*
		silymarine and paracetamol	0.979	1.000	1.000




Fig 1, 2 and 3: light microscopic view of rat liver tissue after treatment with paracetamol





Fig 4 and 5: light microscopic view of rat liver tissue after treatment with Silymarin





Fig 6 and 7: light microscopic view of rat liver tissue after treatment with acacia catechu extract

In **Figure 1, 2 & 3** shows histological morphology of rat liver which are treated with paracetamol. There is congestion, centrilobular hepatocyte necrosis and mild ballooning degeneration. In **Figure 3**, portal tract shows lymphocytic and plasmacytic infiltration, ballooning degeneration of hepatocyte along with generalised congestion. **Figure 4 & 5** group III rats which were treated with Silymarin along with paracetamol show mild congestion of central vein and mild ballooning degeneration of hepatocyte with normal looking portal triad. **Figure 6 & 7** group IV rats which were treated with acacia catechu extract show central vein with normal looking hepatocyte and portal tract with mild congestion.

DISCUSSION

Paracetamol, one of the most commonly used analgesic and antipyretic drug. Liver injury can be induced in experimental animals using paracetamol, carbontetrachloride, ethanol or any other hepatotoxic agent. There are several animal models have been shown to be effectively induce hepatotoxicity in animals [5].

In our study paracetamol was used to induce hepatotoxicity in experimental animal i.e. rats. After administration of paracetamol there was significant increase in serum enzyme SGOP, SGPT and ALP levels and characteristic histological changes as depicted in **Figures 1, 2 And 3**, thus supporting the evidence of paracetamol induced hepatotoxicity.

Paracetamol is primarily metabolised in the liver by conjugation with glucuronic acid (about 60%) and sulphuric acid (35%) or cysteine (~3%). Small amounts of hydroxylated and deacetylated metabolites also have been detected. Paracetamol also undergoes cytochrome (CYP) enzyme mediated N-hydroxylation to form highly reactive metabolite N-acetyl-p-benzoquinoneimine (NAPQI). Normally, harmful actions of this metabolite is nullified by reaction with sulphhydryl groups present in glutathione, but in higher doses of paracetamol there is excess production of NAPQI leading glutathione depletion from hepatocyte. Depletion of glutathione levels renders hepatocytes highly susceptible to oxidative stress and apoptosis. Hence there is NAPQI mediated lipid peroxidation leading to dysfunction of enzymatic

systems, structural cellular changes leading to hepatic damage [6, 7].

Histological examination of liver sections from group –II (paracetamol group) show congestion, centrilobular hepatocyte necrosis and mild ballooning degeneration as depicted in figures 1,2 and 3 confirming the hepatotoxic effects of paracetamol. These changes were positively co-related with the increase in serum enzyme levels ($p < 0.05$) compared to control group-I. SGPT & SGOT and ALP are more commonly used biomarkers to study the hepatic injury. SGOT and SGPT are sensitive indicators of hepatocyte function, hence marked increase indicative of hepatic tissue injury by paracetamol in group –II [8, 9]. Silymarin a proven hepatoprotective was used as positive control [10]. Silymarin a flavonoid known to be a hepatoprotective but exact mechanism of action is not known. It has been proposed to be an antioxidant, regulator of intracellular glutathione levels and cell membrane stabilizer [11]. It was administered at the dose of 25mg/kg orally in the form of suspension, showed a significant decrease in the serum transaminase levels as compared to group-II paracetamol alone ($P < 0.05$). Histopathological study of Silymarin group (group-III) shows mild congestion as depicted in figures 4 & 5. Thus, confirming its hepatoprotective action. Group-IV animals which were administered with the ethanolic extract of acacia catechu at the dose of 250mg/kg orally, showed a marked decrease in serum enzyme levels as compared to paracetamol alone (group-II) with a significant level of difference $p < 0.05$. Thus showing hepatoprotective action which was also supported by Histopathological changes i.e. normal looking hepatocyte, portal tract with mild congestion as shown in figure 6 & 7. The laboratory values were comparable with that of the Silymarin group without any statistical significant difference. Thus, confirming the hepatoprotective action of acacia catechu.

Acacia catechu is a deciduous, thorny tree which grows up to 12-15 meter in height. In Hindi is called as Khair, Kathu in kannada, kattha in urdu, khadir in Sanskrit and in English called as Indian peninsular catch [12]. Traditionally it is been used in healing wounds, as antimicrobial, anti-inflammatory and antifungal [13]. Also has been used in treatment of bleeding, treatment of diarrhoea and as astringent.

Concentrated extracts mainly contain tannins (2-20%), catechins (25-33%), phlobotannins like catechutannic acid 20-50%, flavonoids, gums, resins and pigments.

Tannins have astringent action, whereas catechins have hepatoprotective, anti-oxidant, antidiarrhoeal and hypoglycaemic action [2]. In our study phytochemical analysis showed presence of tannins, alkaloids, proteins and resins. From the study results, which shows a significant hepatoprotective activity, which was comparable with that of the standard control Silymarin. This significant action is supported by serum biochemical markers and histopathological examination. Thus the protective action can be attributed towards the presence of tannins, catechins and flavonoids which are known to have anti-oxidant and free radical scavenging action leading to hepatoprotection [14, 15, 16].

CONCLUSION

Thus it can be concluded that the ethanol extract of acacia catechu possesses a significant hepatoprotective effect. Limitation of our study was we could not isolate active molecule and its structural analysis. Hence studies to establish the exact mechanism of hepatoprotective action at cellular level can be undertaken.

REFERENCES

1. Bunchorntavakul C., Reddy KR., Review article ; herbal and dietary supplement hepatotoxicity Alimentary pharmacology and therapeutics, 13(1):3-17,(2013)
2. Ray D, Sharatchandra KH., Thokchom IS., Antipyretic, antidiarrhoeal, hypoglycaemic and hepatoprotective activity of ethyl catechu extract of acacia catechu wild in albino rats. Indian Journal of Pharmacology, 38(6):408-413(2006)
3. Ewans W C In Trease and Evans, Ed Pharmacognosy 15th Edn Rajkamal Electric press Publisher, 214-252,(2007)
4. Sharma A., Patel VK., Chaturvedi AN., Vibrocidal activity of certain medicinal plants used in Indian folklore medicine by tribals of Mahakoshal region of central India, Indian J Pharmacol, 41(3):129-133(2009)
5. Recknagel RO., Glende EA., Dolak JA., Waller RL., Mechanism of carbontetrachloride toxicity, Pharmacol Ther, 43:139-54(1989)

6. Burke A., Smyth EM., Fitzgerald GA., Analgesics antipyretics, in; Brunton LL., Lazo JS., Parker KL (Eds), Goodman and Gilman's the pharmacological basis of therapeutics, 11th ed, McGraw-Hill Publishers New Delhi 2006, p.671-716.
7. Parmer SR., Varshambhai PH., Kalia K., Hepatoprotective activity of some plants extract against paracetamol induced hepatotoxicity in rats. Journal of Herbal medicine and toxicology, 4(2):101-106(2010).
8. Mahesh A., Shaheetha J., Thonagadurai D., Rao DM., Protective effect of Indian honey on acetaminophen induced oxidative stress and liver toxicity in rats, Biologia,64:1225-31(2009).
9. Martin FL., McLean AEM., Comparison of Paracetamol-Induced Hepatotoxicity in the Rat in vivo with Progression of Cell Injury in Vitro in Rat Liver Slices, Drug and chemical toxicology, 21(4):477-494(1998).
10. Fraschini F., Demartini G., Esposti D., Pharmacology of Silymarin Clinical Drug investigation,22(1):51-65(2002).
11. Panda V., Asher H and Srinath S., Antioxidant and hepatoprotective effect of garcinia indica fruit rind in ethanol induced hepatic damage in rodents, Interdisciplinary Toxicology, 5(4)207-13(2012).
12. Chauhan G., Singh SP., Singh OP., Trigunayat A., Khadir (acacia catechu) a unique ayurvedic remedy, Indian journal of Research, 5:1-4(2011).
13. Patel JD., Kumar V., Bhatt SA., Antimicrobial screening and phytochemical analysis of the resin part of *Acacia catechu*, Pharmaceutical Biology,47(1):34-37(2009).
14. Paya M., Ferrandiz ML., Sanj MJ., Alcaraz MJ., Effects of phenolic compounds on bromobenzene mediated hepatotoxicity in mice, Xenobiotica, 23:327-33(1993).
15. Nithianantham K., Shyamala M., Chen Y., Latha LY., Jothy SL., Sasidharan S., Hepatoprotective Potential of Clitoria ternatea Leaf Extract Against Paracetamol Induced Damage in Mice, Molecules, 16:10134-45(2011).
16. Chao J., Liao J., Peng W., Lee MS., Pao LH., Cheng HY., Antioxidant, Analgesic, Anti-Inflammatory and Hepatoprotective Effects of the Ethanol Extract of Mahonia oiwakensis Stem, Int J Mol Sci, 14:2928-45(2013).



***Corresponding Author:**

Dr. Sheshidhar G Bannale

Assistant professor,

Department of pharmacology

S Nijalingappa Medical College

Bagalkot, Karnataka India 587103

Email: drshashibannale@yahoo.co.in

© 2013; JP RESEARCH Publishers

This is an Open Access article distributed under the terms of the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.—IJPBS--