

EVALUATION OF LANTANA INDICA FOR ANTI ULCER ACTIVITY IN EXPERIMENTAL RATS

Ram Mohan Manda^{1*} & Ganapaty Seru²

¹Dept.of Pharmacognosy & Phytochemistry, Talla Padmavathi Pharmacy College, Warangal-506002, Telangana, India.

²Dept.of Pharmacognosy & Phytochemistry, Githam Institute of Pharmacy, Githam University,
Vishakhapatnam- 530003, Andhra Pradesh, India.

*Corresponding Author Email: rammohanmanda56@gmail.com

ABSTRACT

The leaves of Lantana indica used in the treatment of variety aliments such as ulcer, eczema, antiseptic and in rheumatism. In this study the antiulcer activity of various extracts of Lantana indica leaves were evaluated in pyloric ligation induced gastric ulcer in albino rats. The antiulcer activity was assessed by determining and comparing the ulcer index in the test drug group with that of the vehicle control and standard Ranitidine. The parameters taken to assess antiulcer activity were volume of gastric juice, free acidity, total acidity and ulcer index. The result indicated that Ranitidine, ethanol and aqueous extract significantly (p<0.01) decreased the volume of gastric acid secretion, free acidity, total acidity and ulcer index.

KEY WORDS

Lantana indica, Ranitidine, Anti-ulcer activity, pyloric ligation.

INTRODUCTON

Lantana indica Linn, a member of Verbenaceae is commonly known as "Indian Lantana" is a shrub native to India. Traditionally, the plant was reported to be used in the treatment of a variety of ailments all over the India. It was reported that the leaves of Lantana indica are used in traditional medicine for the treatment of the various human ailments such as eczema, antiseptic, antifungal and in rheumatism^{1, 2} .The volatile oil obtained from the leaves of Lindica was containing transcaryophyllene, α-selinene, globulol, transα-guaienevalencene, caryophyllene oxide, humulene and β-eudesmene³⁻⁵. The sterol fraction of the leaves of *L. indica* contained β-sitosterol and cholesterol⁶. From the roots, oleanolic acid, ursolic acid, 3b-24-dihydroxyolean12-en-28-oic acid and 24-formyl-3-oxoolean-12-en-28-oic acid were isolated. 24-Hydroxy-3-oxooleanolic acid and methyl-24-hydroxy-3-oxoursolate have also been identified in *L. indica*^{7, 8}. Since the plant was reported to have many medicinal uses, the author has studied antiulcer activity leaf extracts.

Plant material:

Fresh leaves (1 kg) of *Lantana indica* was collected from young matured plants from the Etunagaram region, Warangal district during early summer and authenticated by Prof. V. S. Raju, Taxonomist, Kakatiya University, Warangal. A voucher specimen (MRM/11/2012) was deposited in the A. U. College of Pharmaceutical Sciences, Andhra University and Visakhapatnam. The material was dried and powered.

International Journal of Pharmacy and Biological Sciences (e-ISSN: 2230-7605)



Preparation of the extract:

The powdered plant material (500 g) of *Lantana indica* was extracted separately with 1 Liters of distilled water and1 Liter of ethanol by maceration process for 72 h. The solvents were then removed under reduced pressure and dried in a desiccator (aqueous extract yield 11.18%w/w and ethanol extract yield 4.92%w/w with respect to dry material). The extracts were suspended in tween20 (20%v/v in distilled water) and used for the present study.

Anti-ulcer activity of Aqueous and the ethanolic leaf extracts of L.indica

The antiulcer activity of the ethanolic extract of *L.indica* was evaluated on selected albino rats by pyloric ligated ulceration model.

Pyloric ligated ulceration

This test was performed as suggested by Shay et al.. The selected animals were divided into four groups of six in each. Each group of the animals received one of the following test samples through oral route: 20%v/v tween 20 in distilled water (2 ml/kg), ranitidine (20 mg/kg), aqueous extract (200 mg/kg), ethanolic extract (200.mg/kg) After one hour, pylorus ligation was made under ether anesthesia. The animals were then returned to the observation chamber. After 4 h, the animals were sacrificed by decapitation, the abdomen of each animal was opened and the stomach was isolated after suturing the lower esophageal end. The gastric juice was collected by giving a small cut to the pyloric region just above the knot in a measuring cylinder and stomach was opened along the greater curvature. The mucosal layer was washed with 1 ml distilled water and the washings were added to the gastric secretions. The gastric contents were centrifuged at 2000 rpm for 10 min. The supernatant fluid (1 ml) was diluted with 9 ml of distilled water and then titrated against 0.01N sodium hydroxide solution using Topfer's reagent till the solution turns to orange colour. The volume of sodium hydroxide required corresponds to free acidity. The solution was further titrated till the solution regained pink colour. The volume of sodium hydroxide required corresponded to the total acidity.

RESULTS AND DISCUSSION

The findings of the study revealed that the leaf extracts possess significant anti-ulcer activity. All the test samples were found to reduce the volume of gastric acid to a significant extent (p< 0.01), The total acidity and the free acidity also registered significant decrease in a similar manner. The ulcer index was significantly reduced with all test samples. The order of reduction of ulcer score observed was ranitidine < ethanol extract < agueous extract, therefore expressed that the leaves of Lindica exerts anti-ulcer activity which may be due to anticipated inhibition of H₂ receptors resulting in inhibition of gastric acid secretion elicited by histamine and gastrin. The work justifies its use in the traditional system of medicine.

The results were expressed as mean \pm S.E.M and tabulated in Table-2.3.1. Significance of differences between control and treated groups was determined using Student's t-test.

Table-2.3.1: Anti-ulcer activity of the aqueous extract and ethanolic extract of the leaves of *L.indica on* pyloric lignated rats.

Groups	Treatment	Dose	Volume of gastric juice(ml.)	Total acidity (meq/lit)	Free acidity (meq/lit)	Ulcer index
I	Vehicle (20% v/v Tween 20)	2 ml/kg	4.21 <u>+</u> 0.16	16.43 <u>+</u> 0.33	3.05 <u>+</u> 0.09	3.82 <u>+</u> 0.21
II	Ranitidine	20 mg/kg	2.03 <u>+</u> 0.18**	6.52 <u>+</u> 0.22**	0.52 <u>+</u> 0.03**	0.68 <u>+</u> 0.11**
III	Aqueous extract of <i>L.indica</i>	200 mg/kg	3.56 <u>+</u> 0.17*	12.26 <u>+</u> 0.29**	1.3 ± 0.14**	1.85 <u>+</u> 0.26**
IV	Ethanolic extract of <i>L.indica</i>	200 mg/kg	3.38 <u>+</u> 0.23*	10.08 <u>+</u> 0.49**	0.99 <u>+</u> 0.04**	1.37 <u>+</u> 0.28**

Results expressed as Mean \pm SEM from six observations * P < 0.01, ** P < 0.001

CONCLUSION

Based on the findings, it can be concluded that *Lantana indica* possesses significant Anti ulcer activity at all tested dose levels. These activities are dose-related. The results corroborate the basis for the traditional use of the plant in folk medicine.

ACKNOWLEDGEMENT

The authors are thankful to the management and faculty of Talla Padmavathi Pharmacy Colleges Warangal for providing necessary facilities to carry out the present research work.

REFERANCES

- Singh, S.K., V.J. Tripathi and R.H. Singh. A New pentacyclic triterpenre acid from Lantana indica, J. Nat. Prod., 1998;54: 755-758.
- Tripathi, B.K. Rana and V.Taneja. Antifungal activity of Triterpenoids, *Fitoterapia*, 1998, 69: 188-189.
- Akhtar MS, Ali MM, Mir SR, Chemical composition of essential oil of Lantana indica Roxb Leaves . J. Essential Oil Res 2006; 18: 611-612.
- 4. Ram P. Yadav, KiranLata and Ajay Singh, Toxicity of Alcoholic Leaf Extracts of *Lantana indica* Plant:

- Effect on Haematological and Physiological Parameters in Non-target Fish Heteropneustes fossilis International Journal of Fisheries and Aquatic. Sciences, 2013, 2(1): 13-17.
- Javidiqbalpandith, Antifungal activity and phytochemical screening of lant species from northern india world journal of pharmacy and pharmaceutical sciences volume 1(2)810-819.
- Goyal MM, Kamal K, High content of cholesterol in the leaves of Lantana indica Roxb Indian Drugs, 1984;22:41
- Singh, S.K., V.J. Tripathi and R.H. Singh. 3p, 24dihydroxyolean-12-en-28-oic acid, a pentacyclic triterpene acid from lantana indica (1990) *Phytochemistry*, 29: 3360-3362.
- Singh, S.K., V.J. Tripathi and R.H. Singh.Triterpenoids of Lantana indica Roxb Indian Drugs, 1984; 26:395-400.
- Manish srvastava,V.K Srvastava and Ajay singh,Molluscicidal and mosquito larvicicidal efficacy of Lantana indica Roxb.leaf extracts,Natural product radiance vol(2),2007,122
- Hidayat hussain, Javid hussain, Ahmed al-harrasi and Zabta khan shinwari, chemistry of some species genus *lantana*, pak. j. bot., 2011, 43: 51-62.
- 11. Vangoori Y, Klnrr D, Gadekal M, Ramesh P. Evaluation of antiulcer activity of ethanolic

International Journal of Pharmacy and Biological Sciences (e-ISSN: 2230-7605)



Available Online through

www.ijpbs.com (or) www.ijpbsonline.com

extract of leaves of Vitex negundo on pylorus ligature induced and aspirin induced ulcer in albino rats. IJPPS 2013; 5: 476-78.

IJPBS | Volume 5 | Issue 1 | JAN-MAR | 2015 | 01-04

12. Nwafor, P.A., F.K. Okwuasaba , andAnti diarrheal and anti ulcerogenic effects of G. Melchor, 2002.rats. J. Ethanopharmacol., 72: 421-427.



*Corresponding Author: Manda Rammohan

Dept of Pharmacognosy & Phytochemistry Email:rammohanmanda56@gmail.com