



# SALMONELLA ENTERITIDIS – FOOD BORNE PATHOGEN – A REVIEW

Ramya Putturu\*, Tirupathireddy Eevuri<sup>1</sup>, Bindukiranmayi Ch<sup>2</sup>, Krishnaiah Nelapati<sup>3</sup>

 \*Assistant Professor, Dept. of Veterinary Public Health & Epidemiology, College of Veterinary Science, Proddatur,
<sup>1</sup>Assistant Professor, Dept. of Poultry Science, College of Veterinary Science, Proddatur.
<sup>2</sup>Assistant Professor, Dept. of Veterinary Public Health & Epidemiology, College of Veterinary Science, Gannavaram.
<sup>3</sup>Professor & Head, Dept. of Veterinary Public Health & Epidemiology, College of Veterinary Science, Rajendranagar.
\*Corresponding Author Email: puttururamya@gmail.com

# ABSTRACT

Salmonellosis has been a notifiable disease in the U.S since 1943, and S.enteritidis is the second most prevelant cause of infection after Salmonella typhimurium in humans. Salmonella is the leading cause of produced related outbreaks accounting for 30%. Because of the severity of this disease and apparent low infective dose (<10 cells) Salmonella is considered one of the most serious of known food borne pathogens. It is mainly pathogenic to human but in animals it does not induce any clinical symptoms except diarrohea. So the animals act as carriers to Salmonella infection. The majority of transmission is through eating of undercooked eggs, chicken, raw milk, raw vegetables and fruits, contaminated water etc. The conventional isolation procedure includes preenrichment in Buffered peptone water for isolation of S.enteritidis at  $37^{\circ}C$  for 16 h to recover stressed cells and selective enrichment in Rappaportvassilidias broth and tetrathionate broth at  $42^{\circ}C$  for 24 h. Since the infection primarily occurs via faeco-oral route, the preventive measures include proper cooking of meat and eggs, consumption of pasteurized milk, washing fruits and vegetables and drinking chlorinated water apart from personnel hygiene measures.

# KEY WORDS

Food borne, Salmonella, Salmonella enteritidis.

# INTRODUCTION

Salmonella enteric is one of the major bacterial agents that cause food borne infections in humans all over the world (1). Salmonellosis is a major economic problem for the food industry and a public health hazard in many countries (2). Salmonella infected samples must be quickly identified so that they can be isolated and spread of contamination can be controlled (3). Salmonella are among the major bacterial pathogens of poultry in the world (4). Prevention of Salmonella infection is important for poultry health and for the food industry, and prevention can be achieved only by good monitoring and screening programmes (5).

In many countries human salmonellosis is mainly due to consumption of eggs followed by poultry, pork, beef, and dairy products (6). *Salmonella enteritidis* (*S.enteritidis*) has emerged as a pathogen of poultry in mid 1970's, but later became an important human pathogen. *S.enteritidis* is a major cause of human food borne illness and is the most frequent serovar detection in outbreaks of human salmonellosis (7). The increased consumption of fast food of

 ${}^{\rm Page}86$ 

International Journal of Pharmacy and Biological Sciences (e-ISSN: 2230-7605)



animal products and the international food trade between countries have also played an important role in spreading *S.enteritidis* (8). It can also spread via the environment through faecal contamination by humans and animals (9).

The number of *Salmonella*, to be swallowed inorder to cause infection is rather small that is fewer than 10 (10) and it is a must for the livestock products to be tested for the presence of *Salmonella* due to it's potentially low infective dose (11). The symptoms in human beings includes diarrhea, nausea, abdominal pain, mild fever, chills, vomition, prostration, headache, malaise etc. and the diarrhea varies from thin vegetable soup like stools to a massive evacuation with accompanying dehydration (12).

# Prevelence of Salmonella *enteritidis* and zoonotic importance

The first report of food poisoning linked to *Salmonella*, occurred in Germany in 1888, in which 50 persons, who consumed raw ground beef became ill and 1 died (13). Infection in poultry flocks, which is assymptomatic was first noticed in the late 1970's and in 1980's spread rapidly throughout UK, US, South America and other areas and it's incidence increased substantially, showing a 275 fold increase in Argentina and becoming the predominant cause of this disease in U.S (14).

The incidence of *Salmonella typhimurium* infections declined and *S.enteritidis* grew from a serotype constituting only 6% of all *Salmonella* isolates in 1980 to become the most frequently reported *Salmonella* serotype isolated in U.S in 1990, constituting 21% of all *Salmonella* isolates that year (15). The incidence of *S.enteritidis* have dramatically increased from 5% (1976) to 26% (1994) among the salmonellosis cases reported to CDC (USDA, 1995) and in the last few decades, *S.enteritidis* has emerged as a major health problem worldwide (1).

## IJPBS |Volume 5| Issue 1 |JAN-MAR|2015|86-95

Between 1975-1987 the Centers for Disease Control (CDC) determined that no.of S.enteritidis infections in North Eastern U.S had increased about 7 fold (17) and it has expanded in Europe and U.K. In 1990, S.enteritidis became the most frequently reported Salmonella serovars isolated in the U.S (18). Benouda et. al. (19) reported that S.enteritidis represented 56.54% of the total isolated Salmonella strains in Moraco between 1990 and 1997; whereas it was 44% between 1994 and 2002 at Casablanca (20). S. enteritidis and S.typhimurium accounted for ~70% of all human and non human isolates of Salmonella reported worldwide between 1985 and 2008. Infact S.enteritidis, alone accounted for 61.4% of the 1.5 million human isolates of Salmonella reported during this period (21). From 2000 to 2002 a survey of global distribution of Salmonella serovars proved Salmonella enteric sub spp enteric serovar enteritidis to be the far most isolated from common serovar humans accounting for 65% of all isolates, more pronounced in Europe, where it is associated with 85% of all human cases (22). In Thailand the incidence of S.enteritidis among salmonellosis has increased in both human beings and animals in 2006 compared to 2005 (23).

Salmonella is still one of the most important food borne pathogen in industrialized countries with an estimated incidence of salmonellosis per 1,00,000 people ranging from 157 in 1999 to 92 in 2004, Belgium is one of the countries with the highest incidence compared with other European countries (24).

# Pathogenesis

The virulence of *Salmonella* relates to their ability to invade host cells, replicate in them and resist both digestions by phagocytes and destruction by the complement components of plasma (25). Adhesion of *S.enteritidis* to cell surface by fimbriae is an essential stage in colonization and

 ${}^{\rm Page}87$ 

International Journal of Pharmacy and Biological Sciences (e-ISSN: 2230-7605)



pathogenesis of salmonellosis (26 & 27). The association and penetration of the bacterium in the digestive mucosa is a prerequisite for systemic infection (28). Following adherence, probably through fimbrial attachment to the surface of intestinal mucosal cells, the bacteria induce ruffling of cell membranes and releases effector proteins through type III secretion system (25).

In a susceptible host, S.enteritidis replicates primarily in the mucosa of digestive tract after oral challenge and localises in mucosa of ileum, caecum, colon and mesenteric lymph nodes and then spreads to the spleen, liver and various other organs and tissues (29). S. enteritidis colonizes the reproductive organs of infected birds without causing discernable illness and survives host defences during the formation of the egg (30). The production of a capsule like O antigen structure by certain wild type strains of S.enteritidis has been associated with reproductive tract tropism and improved survival within eggs (31). Grade A shell eggs have been attributed to the human S.enteritidis related illness because it can infect ovarian tissues of laying hens and be deposited into the developing egg, resulting in contamination of egg contents (32).

## Transmission

Salmonella can be introduced into poultry houses by means of contaminated feeds, particularly those that contain animal raw materials (33) or by water or vectors such as insects, rodents, persons and equipment (34). Wild rodents and wild life are known as vector for *Salmonella*, flies may be involved in the dissemination of *Salmonella* in the environment as carrier of microorganisms because flies were positive for *Salmonella* on the highly contaminated farms (35). The transmission of *S.enteritidis* to humans by internally contaminated table eggs possesses a challenge to

#### IJPBS |Volume 5| Issue 1 |JAN-MAR|2015|86-95

public health, even though efforts to control this problem are at higher side (36).

*S.enteritidis* only rose to its predominant position in man and poultry in the mid 1980's as a result of vertical and horizontal transmission within and between large poultry organizations in many parts of the world (37). Grade A shell eggs have been attributed to the human *S.enteritidis* related illness because *S.enteritidis* can infect ovarian tissues of laying hens and be deposited into the developing egg, resulting in contamination of egg contents (38&32). Ingestion is the main route of infection in Salmonellosis although, it can occur through the mucosa of the upper respiratory tract and conjunctiva (39).

## Human salmonellosis

Salmonellosis has been a notifiable disease in the U.S since 1943, and S.enteritidis is the second most prevalent cause of infection after Salmonella typhimurium in humans (21). S.enteritidis emerged as an important pathogen in poultry and human (27). Human S.enteritidis infections showed a dramatic increase since 1980's, particularly in developing countries and has became the most commonly isolated serotype in many countries (40). In Canada, S.enteritidis has been among the top 3 reported Salmonella serovars resulting in human disease since 1998, accounting for between 12-21% of infections caused by Salmonella (41). S.enteritidis emerged as a major cause of human salmonellosis worldwide throughout the 1980's-1990's and out of 309 S.enteritidis outbreaks with a confirmed source, 214 (78%) were associated with raw or undercooked shell eggs between 1990-2001 (36). S. enteritidis is clinically prevalent serovar associated with the consumption of foods containing eggs or poultry meat from systemically infected chicken (42) and it is the most common serovar in poultry and humans in Finland (43). S.enteritidis infects fowls such as chicken without

severe illness, but when transmitted to humans,

International Journal of Pharmacy and Biological Sciences (e-ISSN: 2230-7605)

Ramya Putturu\* et al



it causes severe diarrhoea and bacteraemia (44). The infection dose of non-typhoidal Salmonella for humans generally described as  $10^2$  to  $10^3$ organisms; however doses of 10 to 20 organisms and even fewer than 10 organisms have caused illness (10). Forshell and Wierup (12) reported clinical signs include diarrhoea, nausea. abdominal pain, mild fever and chills. The diarrhoea varies from a few thin vegetable soup like stools to a massive evacuation with accompanying dehydration. Vomiting, prostration, anorexia, headache and malaise may also occur with syndrome lasts for 2 - 7 d. Gantois et. al. (30) reported that S.enteritidis, upon infection of a human host causes self limiting gastroenteritis, similar to that caused by other non typhoidal Salmonella serovars. Invasive complications like osteomyelitis, pyoarthrosis, nephrosis (45), lung abscess, appendicitis, cholecystitis, salphingitis, peritonitis, pneumonia, pleurisy, cerebral abscess (46), urinary tract infection (47), endocarditis (48), empyaema, purulent meningitis, intestinal perforation (49), hepatitis (50) and haemolytic ureamic syndrome (51) have been reported. Average fatality rate in Salmonella infections have been reported to be 5.3% (45). But certain serotypes can cause a fatality rate of upto 21% (52).

## Food borne salmonellosis

Salmonella is the most common food poisoning bacteria associated with refrigerated poultry (53). As all Salmonella serovars are primarily pathogenic for animals, the most common source of Salmonella infection for humans is contaminated food of animal origin (54). Salmonella enterica serovar enteritidis is the main cause of food borne salmonellosis (55) and it has been a major causative agent of food borne gastroenteritis in humans (56). The majority transmission is through eating of undercooked contaminated chicken, meat and consumption of raw eggs, egg products, salads, vegetables, fruits, chocolate, maize snack and also through bean sprouts (57&58). Hence the prevalence and occurrence of *S.enteritidis* in foods of animal origin are detailed here under.

## Eggs

More than 70% of S.enteritidis infections reported, the source of infection has been traced to grade A table eggs (38). Eggs are contaminated internally with S.enteritidis probably as a result of infection in the hens ovaries and oviducts (59). Egg contamination can originate before oviposition by direct contamination of the yolk, albumen, or egg shell membranes with bacteria from the infected reproductive organs of the birds or after or during the oviposition by penetration of bacteria from contaminated faeces through the egg shell (30). Hogue et. al. (60) reported that 10% to 40% of pasteurized liquid eggs at commercial egg breaking plants contain S.enteritidis in USA.

Salmonella enterica serotype enteritidis has been associated with food borne outbreaks in which contaminated egg-based foods such as homemade icecream, cookie batter, caesar salad and hollandaise sauce, were consumed (61). S.enteritidis emerged as a major cause of human salmonellosis worldwide throughout the 1980's-1990's and out of 309 S.enteritidis outbreaks with a confirmed source, 214 (78%) were associated with raw or undercooked shell eggs between 1990-2001 (36). Infected hens can deposit S.enteritidis into the edible contents of developing eggs which poses a continuous problem for controlling salmonellosis (62 & 63). Eggs cooled had 91-100% penetration by this organism compared to uncooled eggs which had 48% penetration (64).

## **Poultry Meat**

Boonmar *et. al.* (65) reported that some *Salmonella* infections in humans are due to the consumption of contaminated broiler chicken meat. Hoop and Pospischil (66) tested 7 samples

 ${}^{\rm Page}89$ 

International Journal of Pharmacy and Biological Sciences (e-ISSN: 2230-7605)

Ramya Putturu\* et al



taken from ovary and oviduct of poultry which are positive by culture method, 6 were shown positive for S.enteritidis by immune histochemical method. Recently it has been identified that poultry, consumption of raw eggs, poultry meat, processed products, fast food and international food trade between the countries are the major sources of infection and transmission of S.enteritidis in humans (67). Wang et. al. (68) tested 285 samples of chicken meat, eggs and swabs of chicken related samples and reported that only 1% of the samples were positive for Salmonella enteric serovar enteritidis. Hassanien et. al. (69) examined 75 samples of beef, chicken and reported that Salmonella was detected in 20% of minced frozen beef, 36% of frozen chicken leg and 52% of the frozen chicken fillets and the prominent Salmonella serovars were S.enteritidis and Salmonella kenchuki.

## Milk

Salmonella organisms has been isolated from milk and its products. Most milk borne salmonellosis has been associated with raw, heat treated, or inadequately pasteurized milk or with milk contaminated after pasteurization (70). Ryan et. al. (71) reported a large outbreak of salmonellosis in U.S with an estimated number of 1,80,000 cases caused by the consumption of 2 brands of pasteurized milk produced at the same dairy. Bean et. al. (72) reported an outbreak of salmonellosis involving 1,50,000 persons in the U.S attributed to pasteurized milk. Milk products like cheese have also been implicated in a number of salmonellosis outbreaks (73). Postprocessed skimmed milk (74), fresh cream cakes (75) and spray dried milk powder (76) have also been reported to cause salmonellosis.

## Pork and Beef

Zaidi *et. al.* (77) studied the incidence of nontyphoidal *Salmonella* in pork and beef from retail shops in Mexico and reported that out of 339 and 126 pork and beef samples tested 197 (58.1%) and 68 (54%) were positive for *Salmonella* organisms. *Salmonella* organisms has been isolated from beef (78), chevon (79), mutton (80). Several outbreaks of *Salmonella* food poisoning associated with eating beef have also been reported (81).

## Other Foods

Human salmonellosis has also been associated with fresh fruits, vegetables and vegetable salads (82) and fresh fruit juices (80). Foods commonly associated with outbreaks of Salmonella are milk based products (chocolate, cheese, salads, ice cream etc.), sliced cold meat, vegetables and fruits (57). A wide variety of foods are however known to be associated with S.enteritidis including fresh and dry fruits and vegetables, contamination can during where occur fertilization, irrigation, harvesting or handling under non- optimal hygiene conditions (32). Evans et. al. (83) and Arumugaswamy et. al. (84) reported salmonellosis outbreaks consequent to the consumption of fish and shell fish. In a study conducted by Wilson and Moore (85), Salmonella serotypes were recovered from 8% of 433 mollusks examined.

# Faeces and environmental samples Poultry faeces

Prescott and Gellner (86) reported that 18% of chicken flocks examined contained *S.enteritidis* in the chicken's colon at the time of slaughter. *S.enteritidis* may be transmitted between the poultry via faecal contamination and when *S.enteritidis* was orally infected, was found in the birds livers, caeca, oviducts and ovaries after 3 wk (87). Turkyilmaz *et. al.* (88) used PCR method for isolation and identification of *S.enteritidis* and reported that out of 587 cloacal swabs collected from turkeys 86 (14.7%) were *Salmonella* positive and out of 86 *Salmonella* 11 (1.9%) were *S.enteritidis*.

International Journal of Pharmacy and Biological Sciences (e-ISSN: 2230-7605)

Ramya Putturu\* et al



He et. al. (89) estimated the no. of S. enteritidis after oral challenge in different organs of poultry and reported that S.enteritidis reached a peak at 24 h - 36 h post inoculation, with the liver and spleen containing high concentrations of S.enteritidis, where as the blood, heart, kidney, pancreas, gall bladder had low concentrations. S.enteritidis populations began to decrease and were not detectable at 3 d post inoculation, but were still present upto 12 d post inoculation in the gall bladder, 2 wk for liver and 3 wk for spleen without causing apparent symptoms.

## Human faeces

The number of Salmonella organisms present in the faeces of an infected humanbeings will be approximately  $10^{9}$ /g, which will fall gradually (90) and 5% of the patients upon recovery from this disease may become carriers who shed the organisms in their faeces (91). Hassanien et. al. (69) examined 28 children stool cultures collected from hospitalized children with diarrhoea, fever and reported that 10.71% (21) were positive for Salmonella.

#### Other birds faeces

Vanhoosar and Welsh (92) isolated Salmonella from ratites and reported that 46 out of 248 ostrich samples and 34 out of 99 emu samples and 16 out of 60 rheas were positive for various spp. of Salmonella. Turkyilmaz et. al. (88) isolated Salmonella enteric sub. spp. enteric serovar enteritidis, a cause of food poisoning in humans from turkeys and reported that of the 587 cloacal swabs, 86(14.7%) were Salmonella positive and of these, 11(1.9%) were serovar enteritidis.

## **Environmental Samples**

Gast et. al. (63) used electrostatic air sampling device for collection of air samples 3 times a week from experimentally infected laying hens for isolation of S.enteritidis and reported that electrostatic collection of air was superior over traditional methods for detecting S.enteritidis in environmental samples.

## DIAGNOSIS

The detection of Salmonella in foods is problematic due to presence of fewer no.of organisms, together with larger no.of competing microflora and also due to injured organisms by different food processing methods (93). The conventional culture method, which is routinely used for isolation of Salmonella is time consuming, laborious and may not be suitable for viable but non culturable (VBNC) state of the organisms (11). PCR is rapid, specific and sensitive method for the detection of food borne pathogens (94). The increase in the frequency of the outbreaks and the no.of cases has prompted public health officials for accurate detection within few hours.

The conventional isolation procedure involves pre enrichment in Buffered peptone water (BPW) for isolation of S. enteritidis at 37°C for 16h to recover stressed cells (95) and selective enrichment in Rappaportvassilidias broth and tetrathionate broth at 42°C for 24h The most commonly used selective enrichment for isolation and identification of *S.enteritidis* are Rapaport Vassilidias soya broth, Selenite cystene broth and Tetrathionate broth (3). Then streaked on selective agar like BGA, XLD, HEA, BSA at 37°C for 24h. Salmonella will give black centered colonies.

## PREVENTION AND CONTROL

Since the infection primarily occurs via faeco-oral route, the preventive measures include food hygiene measures like proper cooking of meat and eggs, consumption of pasteurized milk, washing fruits and vegetables especially those to be eaten raw and drinking chlorine treated water and personnel hygiene measures like washing hands after toilet visits.

#### REFERENCES

Herikstad, H., Motarjemi, Y. and Tauxe, R.V. (2002). 1. Salmonella surveillance: a global survey of public health serotyping. Epidemiol. Infect., 129: 1-8.

International Journal of Pharmacy and Biological Sciences (e-ISSN: 2230-7605)

Ramya Putturu\* et al



- Wan J, King K, Carven H, Mcauley C, Tan S.E, Coventry M.J. (2000) Probelia<sup>™</sup> PCR system for rapid detection of *Salmonella* in milk powder and ricotta cheese. Lett Appl Microbiol 30: 267-271.
- Soumet, C., Ermel, G., Rose, N., Rose, V., Drouin, P., Salvat, G. and Colin, P. (1999). Identification by a multiplex-PCR-based assay of *Salmonella typhimurium* and *Salmonella enteritidis* starins from environment swabs poultry houses. *Lett. Appl. Microbiol.*, 29: 1-6.
- Carli, K.T., Unal, C.B., Caner, V. and Eyigor, A. (2001). Detection of *Salmonellae* in chicken faeces by a combination of tetrathionate broth enrichment, capillary PCR, and capillary gel electrophoresis. *J. Clin. Microbiol.*, 39(5): 1871-1876.
- Khuri-bulos N.A, Abu Khalaf M, Shahabi A, Shami K. (1994) Foodhandler-associated *Salmonella* outbreak in a university hospital despite routine surveillance cultures of kitchen employees. Infection Control and Hospital Epidemiol 15: 311-314.
- Carraminana, J.J., Yanguela, J., Blanco, D., Rota, C., Agustin, A., Arino, A. and Herrera, A. (1997). Salmonella incidence and distribution of serotypes throughput processing in a Spanish poultry slaughterhouse. J. Food Prot., 60: 1312-1317.
- Ochoa-Reparaz, J., Sesma, B., Alvarez, M., Renedo, M.J., Irache, J.M. and Gamazo, C. (2004). Humoral immune response in hens naturally infected with *S.enteritidis* against outer membrane proteins and other surface structural antigens. *Vet. Res.*, 35:291-298.
- Landeras, E.A., Havia, G., and Mendoza, C. (1998). Molecular epidemiology of *Salmonella serotype enteritidis* relationship between food water and pathogenic strains. *Int. J. Food Microbiol.*, 43: 81-90.
- Okafo C.N, Umoh V, Galadima M. (2003) Occurence of pathogens on vegetables harvested from soils irrigated with contaminated streams. Sci Total Environ 311(1-3): 49-56.
- Matsui, T., Suzuuki, S., Takahashi, H., Ohyama, T., kobayashi, J., Izumiya, H., Watanabe, H., Kasuga, F., Kijima, H., Shibata, K. and Okabe, N. (2004). Salmonella enteritidis outbreak associated with a school lunch dissert: cross contamination and a long incubation period, Japan, 2-1. Epidemiol. Infect., 132: 873-879.
- Bennett, A.R., Greenwood, D., Tennant, C., Banks, J.G. and Betts, R.P. (1998). Rapid and definitive detection of *Salmonella* in foods by PCR. *Lett. Appl. Microbiol.*, 26:437-441.
- 12. Forshell, L.P. and Wierup, M. (2006): Salmonella contamination: A significant challenge to the global marketing of animal food products. Scientific and

## IJPBS |Volume 5| Issue 1 |JAN-MAR|2015|86-95

*Technical Review of the Office International des Epizooties* 25 (2): 541-554.

- 13. Karlinski, J. (1889). Zur kenntis des *Bacillus enteritidis* gaertner. Zentralbl. *Bakteriol. Parasitenkd.,* 6: 289-292.
- Agron, P.G., Walker, R.L., Kinde, H., Sawyer, S.J., Hayes, D.C., Wollard, J. and Andersen, G.L. (2001). Identification of Substractive Hybridization of sequences specific for *Salmonella enteric serovar enteritidis*.
- 15. Rodrigue, D.C., Tauxe, R.V. and Rowe, B. (1990). International increase in *Salmonella enteritidis* : A new pandemic?. *Epidemiol. Infect.*, 105: 21-27.
- USDA. (1995). Tracking food borne pathogens from farm to table: data needs to evaluate control points. *Conference proceedings.*, 9-10.
- 17. Guthrie, R.H. (1992). *S.enteritidis* in eggs. *Chapter 7 in Salmonella, CRC press.* Boca Raton, Florida.
- Mishu, B., Koehler, J., Lee, L.A., Rodrigue, D., Brenner, F., Blaka, P. and Tauxe, R.V. (1994). Outbreaks of *S.enteritidis* infections in the U.S, 1985-1991. *J. Infect. Dis.*, 169: 547-552.
- Benouda, A., Zouhdi, M. and Alaoui, M. (1997). Epidemiologie des Salmonella non typhoidiques. Revue Marocaine de Biologie-Infectiologie., 3: 59-62.
- Alial, F., Boushiha, A., Jouadi, Z., Adnane, F. and Abid, A. (2004). Forty-one pediatric cases of non-typhoidal Salmonellosis. *Medecine et maladies infectieuses.*, 34: 206-209.
- CDC. (2008). Salmonella Surviellance: annual summary, 2006. US. Department of Health and Human services, CDC, Atlanta, GA.
- Galanis, E., Lo Fo Wong, D.M., Patrick, M.E., Binsztein, N., Cieslik, A., Chalermchikit, T., Aidara-Kane, A., Ellis, A., Angulo, F.J., Weener, H.C. and World Health Organization Global Salm-Surv. (2006). Web-based surveillance and global *Salmonella* distribution, 2000-2002. *Emerg. Infect. Dis.* 12: 381-388.
- Public Health Agency of Canada. (2007). Salmonella Heidelberg—ceftiofur-related resistance in human and retail chicken isolates. Public Health Agency of Canada, Guelph, Ontario,Canada.http://www.phacaspc.gc.ca/ciparspicra/heidelberg/pdf/heidelberg\_e.pd f.
- European Commission. (2005). Trends and sources of zoonotic agents in animals, feeding stuffs, food and man in the European Union and Norway in 2003. Part 3: Salmonella. SANCO/339/2005. European Commission, Health and Consumer Protection Directorate-General, Directorate D-Food Safety. European Commission.
- 25. Saylers, A.A. and Whitt, D.D. (1994). Bacterial pathogenesis. *ASM Press, Washington, DC.* 229-243.

## International Journal of Pharmacy and Biological Sciences (e-ISSN: 2230-7605)

Ramya Putturu\* et al

Page 97.



- Rajashekara, G., Munir, Sh., Alexeyev, M.F., Halvorson, D.A., Wells, C.L. and Nagaraja, K.V. (2000). Pathogenic role of *Sef14* and *Sef17* and *sef21* fimbriae in *Salmonela enteric enteritidis* infection of chickens. *Appl. Environ. Microbiol.*, 66(4): 1759-1793.
- Kisiela, D.M., Kuczkowski, A., Wieliczko, I., Sambor, M., Mazurkiewicz, And Ugorski, M. (2003). Comparision of SefA, FimA, and AgfA fimbrial proteins of Salmonella enteritidis in their abilities to elicit humoral immune response in hens. Bull. Vet. Inst. Pulawy., 47: 95-105.
- Rychlik, I., Lovell, M.A. and Barrow, P.A. (1998). The presence of genes homologous to the k88 genes faeH on the virulence plasmid of *Salmonella* gallinarum. *Microbiol. Let.*, 159: 255-260.
- Dibb-Fuller, M.P., Allen-Vercoe, E., Thorns, C.J. and Woodward, M.J. (1999). Fimbriae- and flagellamediated association with and invasion of cultured epithelial cells by *Salmonella enteritidis*. *Microbiology.*, 145: 1023-1031.
- Gantois, I., Ducatelle, R., Pasmans, F., Haesebrouck, F., Gast, R., Humphrey, T.J. and Van Immerseel, F. (2009). Mechanisms of egg contamination by *Salmonella enteritidis*. FEMS *Microbiol. Rev.*, 718-738.
- Guard-Bouldin, J., Gast, R.K., Humphrey, T.J., Henzler, D.J., Morales, C. and Coles, K. (2004). Subpopulation characteristics of egg-contaminating *S.enteritidis* as defined by the lipopolysacharide O chain. *Appl. Environ. Microbiol.*, 70: 2756-2763.
- Humphrey, T. (2006). Public health aspects of Salmonella enterica in food production. In Mastroeni P. and Maskel, D., (Eds), Salmonella infections. Clinical immunological and molecular aspects. Cambridge university press, Newyork., 89-116.
- Hofer, E., Silva Filho, S.J. and Reis, E.M.F. (1998). Serovars de Salmonella isolates de materias primas e de racao para aves no brasil. *Pesquisa veterinaria Brasiliera.*, 18(1): 21-27.
- Gast, R.K., Holt, P.S., Nasir, M.S., Jolley, M.E. and Stone, H.D. (2003). Detection of *Salmonella enteritidis* in incubated pools of egg contents by flourescence polarization and lateral flow immunodiffusion. *Poultry Sci.*, 82: 687-690.
- 35. Letellier, A. (1999). Distribution of *Salmonellae* in swine herds in Quebec. *Vet. Microbiol.*, 67: 229-306.
- CDC, (2003). Outbreaks of *S.Enteritidis* infections associated with eating raw or under cooked shell eggs, United States, 1999- 2001. *MMWR.*, 51: 1149-1152.
- 37. OIE. (2008). Terrestrial manual: Salmonellosis., 1267-1283.
- Stephenson, P., Satchell, F.B., Allen, G. and Andrews, W.H. (1991). Recovery of *Salmonella* from shell eggs. *J. Assoc. Off. Anal. Chem.*, 74: 821-826.

## IJPBS |Volume 5| Issue 1 |JAN-MAR|2015|86-95

- Fox, J.G. and Gallus, C.B. (1977). Salmonella associated conjunctitis in a cat. J. American Vet. Med. Association., 171: 845-847.
- 40. Rabsch, W., Tschape, H. and Baumler, A.J. (2001). Non typhoidal Salmonellosis: Emerging problems. Microbes *Infect.*, 3: 237-247.
- Demczuk, W., Woodward, D., Ahmed, R., Clark, C., Tabor, h., Dore, K., Clampa, N. and Muckle, A. (2005). A laboratory surveillance data for enteric pathogens in Canada: annual summary 2002 and 2003. *Winnipeg, MB, Health* Canada.
- 42. Cox, J. (1995). *Salmonella enteritidis*: Virulence factors and invasive infection in poultry. *Trends Food Sci. Technol.*, 61: 407-410.
- Pelkonen, S., Romppanen, E.L., Siitonen, A. and Pelkonen, J. (1994). Differentiation of *Salmonella* serovar Infantis isolates from human and animal sources by fingerprinting IS 200 and 16S rrn Loci. *J. Clin. Microbiol.*, 32: 2128-2133.
- 44. Kapley, A., Lampel, K. and Purohit, H.J. (2000). Thermocycling steps and optimization of multiplex PCR. *Biotechnol Lett.*, 22: 1913-1918.
- Prakash, K. and Ray, B.G. (1969). Pyogenic meningitis caused by *Salmonella* cholera suis var. Kunzendorf. *Indian J. Med. Res.*, 58: 158-160.
- Mussini, C., Trenti, F., Manicardi, G., Mongiardo, N., Codeluppi, M., D'andrea, L., Guaraldi, G., Squadrini, T. and De Reinzo, B. (1995). Non-Typhoid Salmonella subdural empyema in a patient with AIDS. Scand. J. Infect. Dis., 27: 173-174.
- Saphra, I. and Winter, J.W. (1957). Clinical manifestations of Salmonellosis in man. *New Eng. J. med.*, 256: 1128-1134.
- 48. Kotto, R.M. and Bovelet, B.A. (1999). Infectious endocarditis due to *Salmonella typhimurium* in a HIV infected patient. *Cardiologie tropicale.*, 25: 79-80.
- Chao, H.C., Chin, C., Kong, M., Chang, L., Huang, Y., Lin, T. and Lou, C. (2000). Factors associated with intestinal perforation in children's non-typhi *Salmonella* toxic megacolon. *Pediatric Infect. Dis. J.*, 19: 1158-1162.
- Gurakan, F., Derman, O., Yasmaris, A. and Ece, A. (2000). Distinguishing features of *Salmonella* and Viral hepatitis. *Pediatric Infect. Dis. J.*, 19: 587.
- 51. Kalinowska, N.A. and Biesiada, G.G. (2000). Haemolytic uremic syndrome in a patient with bacterial food poisoning. *Przegl-lek.*, 57: 54-55.
- Chen, Y., Chen, T., Lu, P., Su, Y., Hwang, K., Tsai, J., Cheng, H. and Penh, C. (1999). *Salmonella cholera suis* bacteraemia in southern Taiwan. *Kaosiung J. Med. Sci.*, 15: 202-208.
- 53. Cohen, N.D., Martin, I.J., Simpson, B., Wallis, D.E. and Neibergs, H.E. (1996). Comparision of polymerase

## International Journal of Pharmacy and Biological Sciences (e-ISSN: 2230-7605)

Ramya Putturu\* et al



# Available Online through

## www.ijpbs.com (or) www.ijpbsonline.com

chain reaction and microbiological culture for detection of Salmonellae in equine faeces and environmental samples. Am. J. Vet. Res., 57: 780-786.

- 54. Pignato, S., Marino, A.M., Emanuelle, M.C., Lannotta, V., Carcappa, S. and Giammanco, G. (1995). Evaluation of new culture media for rapid detection and isolation of Salmonellae in foods. Appl. Environ. Microbiol., 61: 1996-1999.
- 55. Collighan, R.J. and Woodward, M.J. (2001). The SEF14 fimbrial antigen of S.enteritidis is encoded within a pathogenicity islet. Vet. Microbiol., 80: 235-245.
- 56. Clayton, D.J., Bowen, A.J., Hulme, A.W., Buckley, A.M. and Deacon, V.L. (2008). Analysis of the role of 13 major fimbrial subunits in colonization of the chicken intestines by S. enteritidis reveals a role for a novel locus. BMC Microbiol., 8: 228-228.
- 57. O'Mahony, M., Cowden, J., Smyth, B., Lynch, D., Hall, M., Rowe, B., Teare, E.I., Tettmar, R.E., Coles, A.M. and Gilbert, R.J. (1990). An outbreak of Salmonella saintpaul infection associated with bean sprouts. Epidemiol. Infect., 104: 229-235.
- 58. D'Aoust, J.Y. (1997). Salmonella spp, p. In Doyle MP, Beurhat, L.R. and Mantville, T.J. (Eds). Food microbiology. fundamentals and frontiers. ASM press, Washington D.C., 129-158.
- 59. Gast, R.K., and Beard, C.W. (1990). Production of Salmonella enteritidis – contaminated eggs by experimentally infected hens. Avian. Dis., 34: 438-446.
- 60. Hogue, A.T., Ebel, E.D., Thomas, L.A., Schlosser, W., Bufano, N. and Ferris, K. (1997). Surveys of Salmonella enteritidis in unpasteurized liquid egg and spent hens at slaughter. J. Food Prot., 60: 1194-200.
- 61. Hennessy, T.W., Hedberg, C.W., Slutssker, L., White, K.E., Besser-wiek, J., Moen, M.E., Feldman, J., Coleman, W.W., MacDonald, K.L. and Osterholm, M.T. (1996). A national outbreak of S.enteritidis infection from ice cream. N. Engl. J. Med., 334: 1281-1286.
- 62. Gast, R.K. and Holt, P.S. (2000). Deposition of phage type 4 and 13a Salmonella enteritidis strains in the yolk and albumen of eggs laid by experimentally infected hens. Avian Dis., 44: 706-710.
- 63. Gast, R.K., Petter, G. and Holt, P.S. (2002). Chracteristics of Salmonella enteritidis contamination in eggs after oral, aerosol, and intravenous inoculation of laying hens. Avian Dis., 46: 629-635.
- 64. Fajardo, T.A., Anantheswaran, R.C., Puri, V.M. and Knabel, S.J. (1994). Penetration of Salmonella enteritidis into eggs subjected to rapid cooling. J. Food. Prot., 58: 473-477.
- 65. Boonmar, S., Bangtrakulnonth, A., Pornrunangwong, S., Terajima, J., Watanabe, H., Kaneko, K.I. and Ogawa, M. (1998). Epidemiological analysis of Salmonella

#### IJPBS |Volume 5| Issue 1 |JAN-MAR|2015|86-95

enteritidis isolates from humans and broiler chickens in Thailand by phage typing and pulsed-field gel electrophoresis. J. Clinical Microbiol., 36: 971-974.

- 66. Hoop, R.K. and Pospischil, A. (1993). Bacteriological, serological, histological and immune-histochemical findings in laying hens with naturally acquired Salmonella enteritidis phage type 4 infection. Vet. Rec., 133: 391-393.
- 67. Gibson, D.L. (2000). Salmonella enteritidis thin aggregative fimbriae and the extracellular matrix. Ph.D Thesis, University of Victoria.
- 68. Wang, S.J., Yeh, D.B. and Wei, C. (2009). Specific PCR primers for the identification of Salmonella enteric serovar enteritidis in chicken related samples. J. Food and Drug Analysis., 17(3): 183-189.
- 69. Hassanien, R., Hassan Ali, S.F., El-Malek, A.M.A., Moemen., Mohamed, A. and Elsayh, K.I. (2011). Detection and identification of Salmonella species in minced beef and chicken meats by using multiplex PCR in assiut city. Vet. World., 4(1): 5-11.
- 70. El-Gazzar, F.E. and Marth, E.H. (1992). Salmonellae, Salmonellosis, and dairy foods: A review. J. Dairy Sci., 75: 2327-2343.
- 71. Ryan, C.A., Nickels, M.K., Hargeyy-bean, N.T., Potter, M.E., Endo, T., Mayer, L. Langkop, C.W., Gibson, C., McDonald, R.C., Kenny, R.T., Buhr, N.D., McDonnel, P.J., Martin, R.J., Cohen, M.L. and Blake, P.A. (1987). Massive outbreak of antimicrobial resistant salmonellosis traced to pasteurized milk. J. Am. Med. Assoc., 258: 3269-3274.
- 72. Bean, N.H., Griffin, P.M., Goulding, J.S. and Ivey, L.B. (1990). Food borne disease outbreaks, 5 years summery, 1983-1987. J. Food Prot., 53: 711-728.
- 73. Synnot, M.B., Brindley, M., Gray, J. and Dawson, J.K. (1998). An outbreak of Salmonella agona infection associated with precooked turkey meat. PHLS, Commun. Dis. Public Health., 1: 176-179.
- 74. Lecos, C. (1986). Of microbes and milk: America's worst Salmonella outbreak. Dairy and Food Sanitation., 6: 136-140.
- 75. Evans, M.R., Tromans, J.P., Dexter, E.L.S., Riebeiro, C.D. and Gardner, D. (1996). Consecutive Salmonella outbreaks traced to the same bakery. Epidemiol. Infect., 116: 161-167.
- 76. Rowe, B., Begg, N.T., Hutchison, D.N., Dawkin, H.C., Gilbert, R.J., Jacob, M., Hales, B.M., Rae, F.A. and Jepson, M. (1987). Salmonella eailing infections associated with consumption of infant dried milk. Lancet., ii: 900-903.
- 77. Zaidi, M.B., McDermott, P.F., Fedorka-Cray, P., Leon, V., Canche, C., Hubert, S.K., Abbott, J., Leo'n, D., Zhao, S., Headrick, M., and Tollefson, L. (2006). Nontyphoidal

#### International Journal of Pharmacy and Biological Sciences (e-ISSN: 2230-7605)

Ramya Putturu\* et al



Salmonella from Human Clinical Cases, Asymptomatic Children, and Raw Retail Meats in Yucatan, Mexico. *Clinical Infectious Diseases.*, 42: 21–8.

- Agarwal, R.K., Kapoor, K.N., Verma, J.C., Bachhil, V.N., Singh, B.R., Kumar, A., Sachan, N., Singh, D.K. and Malik, S.V.S. (1999). Isolation and characterization of Salmonella stockholm from beef. Indian J. Comp. Microbiol. Immunol. Infect. Dis., 20: 50-52.
- Das, M.S., Roy, D.K. and Das, S. (1990). Occurrence of Salmonellae in slaughtered pigs, goat meat, meat handlers and slaughter-house workers. J. Com. Dis., 22: 39-42.
- Singh, P., Gupta, R.S. and Misra, D.S. (1994). The enterobacteria from mutton and their enterotoxigenicity. *Indian J. Comp. Microbiol. Immunol. Infect. Dis.*, 15(1&2).
- Roels, T.H., Frazak, P.A. and Kazmierczak, J.J. (1997). Incomplete sanitation of a meat grinder and ingestion of raw ground beef: contributing factors to a large outbreak factors to a large outbreak of *S. typhimurium* infection. *Epidemiol. Infect.*, 119: 127-134.
- O'Mahony, M., Cowden, J., Smyth, B., Lynch, D., Hall, M., Rowe, B., Teare, E.I., Tettmar, R.E., Coles, A.M. and Gilbert, R.J. (1990). An outbreak of *Salmonella* saintpaul infection associated with bean sprouts. *Epidemiol. Infect.*, 104: 229-235.
- Evans, H.S., Madden, P., Douglas, C., Adak, G.K., O'Brien, S.J., Djuretic, T., Wall, P.G. and Stanwell-Smith, R. (1998). General outbreaks of infectious intestinal disease in England and Wales : 1995 and 1996. PHLS Commun. Dis. Public health, 1: 165-171.
- Arumugawsamy, R.K., Rasul, G., Hamid, S.N.A. and Cheah, C.T. (1995). Prevalence of *Salmonella* in raw and cooked foods in Malaysia. *Food Mocrobiol.*, 12: 3-8.
- Wilson, I.G. and Moore, J.E. (1996). Presence of Salmonella spp. and Campylobacter spp. in shell fish. Epidemiol. Infect., 116: 147.



## IJPBS |Volume 5| Issue 1 |JAN-MAR|2015|86-95

- Prescott, J.F. and Gellner, O.S. (1984). Intestinal carriage of *Campylobacter jejuni* and *Salmonella* by chicken flocks at slaughter. *Can. J. Comp. Med.*, 48(3): 329-331.
- Barnhart, H.M., Dreesen, D.W., Bastein, R. and Pancorbo, O.C. (1991). Prevalence of *Salmonella enteritidis* and other serovars in ovaries of layer hens at time of slaughter. *J. Food. Prot.*, 54: 488-491.
- Turkyilmaz, S., Hazimoglu, S. and Bozdogan, B. (2009). Antimicrobial susceptibility and resistance genes in Salmonella enteric serovar enteritidis isolated from turkeys. Israel J. Vety. Med., 64(3).
- He, G.Z., Tian, W.Y. Qian, N., Cheng, A.C. and Deng, S.X. (2010). Quantitative studies of the distribution pattern for *Salmonella enteritidis* in the internal organs of chicken after oral challenge by a real-time PCR. *Vet. Res. Commun.*, 34: 669-676.
- *90.* Taylor, J. and McCoy, J.H. (1969). *Salmonella* and Arizona infections. *Food borne Infections and Intoxications*. 3-72.
- Jay, J.M. (2000). Food borne gastroenteritis caused by Salmonella and shigella. Modern Food Microbiol. 6: 511-528.
- Vanhooser, S.L. and Welsh, R.D. (1995). Isolation of Salmonella species from ratites. J. Vet. Diagn. Invest., 7: 268-269.
- Prusak-Sochaczewski, E. and Luong, J.H.T. (1989). An improved ELISA method for the detection of *Salmonella* typhimurium. *J. Appl. Bacteriol.*, 66: 127-135.
- Olsen, J.E., Aabo, S., Nielsen, E.O. and Nielsen, B.B. (1991). Isolation of a *Salmonella* specific DNA hybridization probe. *APMIS.*, 99: 114-120.
- Josefsen, M.H., Krause, M., Hansen, F. and Hoorfar, J. (2007). Optimization of a 12-hour TaqMan PCR-based method for detection of *Salmonella* bacteria in meat. *Appl. Environ. Microbiol.*, 73: 3040-3048.



 $_{Page}95$ 

International Journal of Pharmacy and Biological Sciences (e-ISSN: 2230-7605)

Ramya Putturu\* et al