

International Journal of Pharmacy and Biological Sciences ISSN: 2321-3272 (Print), ISSN: 2230-7605 (Online) IJPBS™ | Volume 8 | Issue 4 | OCT-DEC | 2018 | 612-622



Research Article | Biological Sciences | Open Access | MCI Approved | ज्ञान-विज्ञान विमुक्तये |UGC Approved Journal |

## CHARACTERIZATION AND POTENTIAL APPLICATIONS OF HALOVIBRIO VARIABILIS MLA3 (A HETEROTROPHIC BACTERIUM) ISOLATED FROM A SALINE-ALKALINE LAKE OF RAJASTHAN, INDIA

Deepesh Kumar Neelam, Akhil Agrawal, Anuj Kumar Tomer and Pawan K Dadheech\*

Department of Microbiology, School of Life Sciences, Central University of Rajasthan, Bandarsindri-305817 (Ajmer), Rajasthan, India

\*Corresponding Author Email: pdadheech@curaj.ac.in

#### ABSTRACT

Halophiles are salt loving microorganisms. These have ability to produce industrially important enzymes showing activity at high salt concentration and alkaline condition. This study describes the isolation, physiological and biochemical characterization, phylogenetic analysis of a moderately halophilic bacterial strain MLA3 and screening of the strain for industrially important enzymes production. The strain MLA3 was isolated from Sambhar Lake, Rajasthan, India. It is a gram-negative, motile, rod shaped with an optimal growth observed at 10% (w/v) NaCl concentration, pH 8.0 and 30 °C. On the basis of 16S rRNA gene sequence analysis, the strain MLA3 was phylogenetically identified as members of the genus Halovibrio and clustered in clade with type species Halovibrio variabilis DSM 3050. The occurrence of the studied strains is an addition of the genus Halovibrio in microbial diversity of Sambhar Lake. The MLA3 was found to be capable to hydrolyse skim milk, Tween80 and starch. It showed maximum hydrolysis zone for starch and skim milk. The amylase and protease enzyme production were estimated to be 7.3  $\pm$  0.05 U/ml and 1.7  $\pm$  0.03 U/ml, respectively. The present study indicates that Sambhar Lake is a treasure of still many unexplored microorganisms which are capable of producing industrially significant enzymes.

#### **KEY WORDS**

Sambhar Lake, Halophile, 16S rRNA gene, Phylogenetic study, Halovibrio, Halophilic enzymes

#### 1. INTRODUCTION

Hypersaline environments such as salt lakes, salt pan and marine water are habitats for survival of microbial community of halophiles in the entire world [1-4]. Soda lakes such as Howz Soltan Lake, Iran [5]; Wadi Natrum, Egypt [6, 7]; Great salt lake, Utah [8]; Mono Lake, California [9] and Lonar lake, India [10] were more emphasized for exploration of culture dependent diversity of halophilic bacteria. In this context, Sambhar Lake attracted researchers more due to its dual nature i.e., salinity and alkalinity [11]. On the basis of salinity and alkalinity, many earlier studies focused on the phylogenetic diversity of halophiles and haloalkaliphiles of the lake and characterized different genera such as Natronobacterium [12, 13], Halobacterium [14], Gracilibacillus [15], Geomicrobium [16], Alkalibacillus Halobiforma [17], Hallobacillus [18], [19], Oceanobacillus [20] and Lentibacillus [21]. Due to unique characteristics of halophiles, many studies focused not only on phylogenetic diversity but also their industrial and biotechnological applicability [15, 22-25]. It was reported that subgroups members of genus Halovibrio and Halomonas were phylogenetically distinct [26]. A number of strains belonging to Halomonas have been reported from Sambhar lake and also examined for industrial applications [27, 28].



Considering the novelty and significance of halophilic heterotophic bacteria and paucity of work on microbial diversity of Sambhar lake, the present study was undertaken to characterize and phylogenetic analysis of a strain MLA3 isolated from this soda lake. The strain MLA3 was also evaluated for the qualitative production of extracellular hydrolytic enzymes such as protease, lipase and amylase as well as quantitative production of amylase and protease.

#### 2. MATERIALS AND METHODS

#### Chemicals

All chemicals used in the present study were procured from HiMedia Laboratories, India

#### Sample collection

The water sample was collected in summer season from opposite site of sludge gate, Guda Jhapog dam area (26°58'N 75°5'E) of Sambhar lake (a saline-alkaline lake), Rajasthan, India. Sample was stored in sterile bottles and transported to laboratory, at Central University of Rajasthan, Ajmer for further analyses.

#### Isolation and purification of strain MLA3

Water sample was initially inoculated for enrichment in the modified MGM broth medium [29] containing (g L<sup>-1</sup>) NaCl, 220; KCl, 5.4; MgSO<sub>4</sub>.7H<sub>2</sub>O, 26.9; MgCl<sub>2</sub>.6H<sub>2</sub>O, 23.1; CaCl<sub>2</sub>.2H<sub>2</sub>O, 0.8; peptone, 10.0, 0.8; sodium lactate, 2.5; and yeast extract, 2. The pH 8.0 of the medium was adjusted with NaOH and incubated at 37 °C on an orbital shaker incubator (Waiometra, India) at 120 rpm for 7 days. Further, the colony of the strain MLA3 was purified by successive streaking on MGM medium agar plates. Isolated strain was stored at 4 °C in refrigerator for further investigations.

# Determination of optimum temperature, salinity and pH for growth

The growth studies were carried out in MGM modified broth medium. The different ranges of temperature (25-55 °C), pH (6-12) and salinity (0-30%) were employed to find out optimal growth of the isolated strain [30]. The turbidity was measured at  $OD_{600}$  nm by using spectrophotometer (HALO DB-30 UV-Visible double beam, Dynamica, Singapore). All experiments were performed in triplicates.

#### Morphological studies of colony and cell

The appearance of colony such as shape, colour, opacity, margin and elevation was observed by magnifying lens (10X) on agar plate containing MGM modified agar medium. Gram staining was performed using earlier protocol [31]. The other morphological features of bacterial strain were observed at 1000X using Axio Lab. A1 compound microscope (Carl Zeiss, Germany).

#### **Biochemical characterization**

A series of biochemical tests such as indole production, methyl red, Voges-Proskauer, catalase activity, nitrate reduction [32], citrate utilization and oxidase activity [33] were performed in triplicates by placing the culture at 30 °C and pH 8.0 for 7 days.

#### Extracellular lipolytic activity

Lipolytic activity of the strain was determined using 1% Tween 80 in MGM modified broth medium containing (g L<sup>-1</sup>) NaCl, 100; KCl, 5.4; MgSO<sub>4</sub>.7H<sub>2</sub>O, 26.9; MgCl<sub>2</sub>.6H<sub>2</sub>O, 23.1; CaCl<sub>2</sub>.2H<sub>2</sub>O, 0.8; peptone, 10.0, 0.8; sodium lactate, 2.5 yeast extract, 2 and pH 8.0. The strain MLA3 was incubated on solided plates at 30 °C for 7 days. The formation of conspicuous halos/zone around the growing isolate indicated the lipolytic activity [34].

#### Extracellular protease activity

The proteolytic activity of the strain was examined in MGM modified broth medium containing (g  $L^{-1}$ ) NaCl, 100; KCl, 5.4; MgSO<sub>4</sub>.7H<sub>2</sub>O, 26.9; MgCl<sub>2</sub>.6H<sub>2</sub>O, 23.1; CaCl<sub>2</sub>.2H<sub>2</sub>O, 0.8; peptone, 10.0, 0.8; sodium lactate, 2.5; skim milk, 10; yeast extract, 2 and pH 8.0. For protease activity, the strain MLA3 was inoculated on skim milk agar plates and incubated at 30 °C for 7 days. The hydrolysis around the culture gave indication of proteolytic activity [35].

#### Quantitative assay of protease enzyme production

Quantitative protease production was performed in 50 ml of MGM modified broth medium containing (g L<sup>-1</sup>) gelatin, 10; casein enzymatic hydrolysate, 10; NaCl, 100; KCl, 5.4; MgSO<sub>4</sub>.7H<sub>2</sub>O, 26.9; MgCl<sub>2</sub>.6H<sub>2</sub>O, 23.1; CaCl<sub>2</sub>.2H<sub>2</sub>O, 0.8; peptone, 10.0, 0.8; sodium lactate, 2.5 yeast extract, 2 and pH 8.0 in 250 ml Erlenmeyer flasks. The MLA3 culture was incubated on orbital incubator shaker (Waiometra, India) at 30 °C with 150 rpm for 7-10 days. Samples were withdrawn at 1-day interval and centrifuged at 10,000 rpm for 10 min at 4 °C. The cell free extract was used as crude enzyme to measure protease activity [36]. The enzyme substrate reaction mixture containing 0.5 ml enzyme was added with 3.0 ml casein (0.6% in 20mM Tris-HCl buffer, pH 8.0) and reaction mixture incubated at 37 °C for 10 min. The reaction was terminated by adding 3.2 ml of TCA mixture (0.11 M trichloro acetic acid, 0.22 M sodium



acetate, 0.33 M acetic acid) and incubated for 30 min at room temperature. The reaction mixture was centrifuged at 14000 g for 10 min at 4 °C and supernatant absorbance was observed at 280 nm by spectrophotometer (HALO DB-30 UV-Visible double beam, Dynamica). One unit of alkaline protease activity was defined as the amount of enzyme liberating 1  $\mu$ g of tyrosine/min under assay conditions. Enzyme units were measured using tyrosine (0–100  $\mu$ g) as standard [37].

#### Extracellular amylase activity

The amylase activity was determined in MGM modified broth medium containing (g L<sup>-1</sup>) starch, 10; NaCl, 100; KCl, 5.4; MgSO<sub>4</sub>.7H<sub>2</sub>O, 26.9; MgCl<sub>2</sub>.6H<sub>2</sub>O, 23.1; CaCl<sub>2</sub>.2H<sub>2</sub>O, 0.8; peptone, 10.0, 0.8; sodium lactate, 2.5 yeast extract, 2 and pH 8.0. Isolate MLA3 was inoculated on solidified plates and incubated at 30 °C for 7 days. After incubation, plates were flooded with Gram's iodine solution (3% (w/v), KI and 0.3% (w/v) iodine). A clear zone around the growth indicated hydrolysis of starch [38].

#### Quantitative assay of amylase enzyme production

Quantitative amylase production was performed in 50 ml of MGM modified broth medium containing (g L<sup>-1</sup>) starch, 10; NaCl, 100; KCl, 5.4; MgSO<sub>4</sub>.7H<sub>2</sub>O, 26.9; MgCl<sub>2</sub>.6H<sub>2</sub>O, 23.1; CaCl<sub>2</sub>.2H<sub>2</sub>O, 0.8; peptone, 10.0, 0.8; sodium lactate, 2.5 yeast extract, 2 and pH 8.0 in 250 ml Erlenmeyer flasks. The strain MLA3 was inoculated on orbital shaker (Waiometra, India) at 30 °C with 150 rpm for 7-10 days. Samples were withdrawn at 1-day intervals and centrifuged at 12000 g for 20 min at 4 °C. The cell-free supernatants were used for the amylase assay. The amylase activity was measured by using DNS method and maltose as the standard. One unit of amylase is defined as the amount of enzyme that liberates one l µmol reducing sugar (maltose) ml<sup>-1</sup>, min<sup>-1</sup> at 30 °C [39-41].

#### **Genomic DNA extraction**

Genomic DNA of halophilic bacterial strain MLA3 was extracted using protocol [42] with some modifications. Briefly, bacterial culture was centrifuged at 9000 g for 5 minutes followed by addition of lysis solution (500  $\mu$ l) and incubated at 65 °C for 5 minutes in water bath. Cell suspension was extracted three times with organic solvent using equal amount of Chloroform: Isoamyl alcohol (24:1) and gently emulsified by inversion 3-5 times and centrifuged at 9000 g for 5 minutes. Aqueous phase was separated into sterile centrifuge tube and extracted with equal volume of phenol: chloroform: isoamyl alcohol (25:24:1) at 9000 g for 5 minutes. Obtained aqueous phase were transferred into sterile centrifuge tube and extracted with equal amount of chloroform: isoamyl alcohol (24:1) at 9000 g for 5 minutes. Upper aqueous phase containing DNA was transferred to a new tube and equal amount of Propan-2-OL (iso-propyl alcohol) was added and mixed gently by inversion at room temperature (37°C) followed by centrifugation at 9000 g for 3 minutes. supernatant was removed completely, and DNA pellet was dissolved in 70  $\mu$ l of NaCl solution. Thereafter, 300  $\mu$ l of cold absolute ethanol was added, the DNA precipitated out in 7 minutes at -20 °C which was centrifuged at 9000 g for 5 minutes. Ethanol was removed, and the was pellet washed once with 70% cold ethanol at 9000 g for 5 minutes. Later, the pellet was dissolved in 50 µl deionzed water by gentle vortexing and kept at -20 for further use.

#### 16S rRNA gene sequencing

The 16S rRNA gene was amplified using PCR master mix (Genei<sup>™</sup> Company) using universal primers 27F and 1495R [43]. Amplified genomic DNA was purified with GeneJET PCR purification kit (Thermo Fisher, India) as per the manufacturer's protocol. Purified product was sequenced commercially (Eurofins Genomics India, Bangalore) to get partial sequences of 16S rRNA gene.

#### **Phylogenetic analysis**

A clean 1466 bp sequence was generated by commercial sequencing of amplified product. The similarity search was done using the 'Identify' option of the EzTaxon database http://www.eztaxon.org [44] to find out the closest strain for molecular identification. Other appropriate 16S rRNA gene sequences of related strains particularly from strains originated from Sambhar Lake were also retrieved for the phylogenetic analysis. Multiple sequence alignment was performed using MUSCLE program [45] inbuilt available MEGA 7 [46]. The alignment was manually edited to obtain unambiguous data. A total of 1262 positions in the final dataset was used for phylogenetic analysis. All the phylogenetic analyses were carried out using the software package MEGA version 7. The suitable model of phylogeny with the lowest Bayesian Information Criterion (BIC) score was selected using the MEGA 7 program which led to the selection of the TN93+G model having the lowest BIC score. The phylogenetic tree was constructed using the maximum likelihood (ML) method. Bootstrapping of



1000 replications was calculated to find out confidence values for the edges of the maximum likelihood tree. *Planococcus citreus* (X62172) was chosen as the outgroup.

#### Nucleotide sequence accession numbers

The nucleotide sequence has been deposited in the GenBank (NCBI database) under accession number MK156419.

#### 3. RESULTS AND DISCUSSION

#### Colony and cell morphology

The present study revealed that colonies of the strain MLA3 were circular with entire margin, smooth surface, sticky nature, orange colored, raised and opaque on MGM modified medium agar plate, as shown in Table 1. The morphological characters rather than colony color are similar to *Halovibrio variabilis* and *Pseudomonas halophila* [47]. Cell morphology result showed that cells were gram negative and rod shaped (fig.1b and Table 1). These results correspond to earlier reported strains *Halovibrio variabilis, Pseudomonas halophila* and *Halovibrio denitrificans* strain HGD 3 [47, 48].




Fig. 1. (a) Colonies of the *Halovibrio* sp. strain MLA3 on MGM modified medium. (b) Bright field microscopy showing the negative rod cell

Table 1: The colony and cell morphology similarity features of *Halovibrio* sp. strain MLA3 with type strain *Halovibrio variabilis* and *Pseudomonas halophila* 

Characters	<i>Halovibrio</i> sp. MLA3	Halovibrio variabilis DSM 3050	Pseudomonas halophila DSM 3051
Isolation source	Sambhar lake, India	Great Salt Lake, Utah, USA	Great Salt Lake, Utah, USA
Shape	Circular	Circular	Circular
Colour	Orange	Light brown	Reddish-brown
Margin	Entire	Entire	Entire
Surface	Smooth	Smooth	Smooth
Elevation	Raised	Raised	Raised
Gram reaction	Negative	Negative	Negative
Cell shape	Curved rod	Curved rod	Rod
Cell size (µm) (width and length)	0.3-0.5 x 1.0- 3.8	0.5-0.8 x 1.0-3.0	0.8-1.0 X 1.5-5.0
Motility	Motile	Motile	Motile

#### Physiological and biochemical characterization

The results of physiological parameters for the growth showed that the strain MLA3 was able to grow optimally at 30 °C, 10% (w/v) NaCl concentration and at pH 8.0

(Fig. 2a, 2b, 2c and Table 2). Whereas, previously reported halophilic strain of *Halovibrio variabilis* DSM 3050 from Great Salt Lake, Utah, USA showed optimal growth at 33  $^{\circ}$ C, 9.35  $^{\circ}$  (w/v) NaCl concentration and at



pH 7.5. Another strain of *Pseudomonas halophila* was able to grow optimally at 28°C, 4.56% (w/v) NaCl concentration and at pH 7.0 [47]. Likewise, *Halovibrio denitrificans* HGD 3 from hypersaline lake sediment from Central Asia showed optimal growth at 28 °C, 11.6

% (w/v) NaCl concentration and at pH range 6.7-8.5 [48]. Moreover, *Halomonas* sp. strain SL121 from Sambhar lake could grow optimally at 11% (w/v) NaCl concentration and pH 8.9 [28].




Fig. 2. Growth pattern of the *Halovibrio* sp. strain MLA3 (a) Temperature 25-55 °C. (b) 0-30 % NaCl concentration in MGM modified medium. (c) pH ranges 5 -12. Each value is a mean of three replicates data with ± standard deviation.

The strain MLA3 showed negative result for indole formation, methyl red and voges-proskauer, citrate utilization, and urease activity (Table 2). It showed positive result for nitrate reduction, oxidase and catalase activity. Similarly, *Halovibrio variabilis* DSM 3050 showed positive result for oxidase and catalase activity [47].

Characters	Halovibrio sp.	Halovibrio	Pseudomonas
Characters	MLA3	variabilis DSM 3050	halophila DSM 3051
NaCl optimum range (% w/v)	10	9.35	4.56
pH optimum	8.0	7.5	7.0
Temperature optimum (°C)	30	33	28
Indole	-	-	-
Methyl Red	-	-	-
Voges-Proskauer	-	-	-
Nitrate	-	-	-
Citrate	-	ND	ND
Catalase	+	+	+
Oxidase	+	+	+


Table 2: Differential features of physiological and biochemical characterization of *Halovibrio* sp. strain MLA3 with type strain *Halovibrio variabilis* and *Pseudomonas halophila* 




#### **Phylogenetic analysis**

The strain MLA3 showed 99.97% identity with coverage of 100% to Pseudomonas halophila DSM 3050(T) in search 1466 bp of 16S rRNA gene using EzTaxon database. During the course of studies on halophilic, Gram-negative bacteria, a comparison of two strains of halophilic bacteria, Pseudomonas halophila DSM 3050 and Halomonas variabilis DSM 3051 (formerly Halovibrio variabilis) demonstrated that the characteristics of strain DSM 3050 corresponded to the original description of Halovibrio variabilis and those of DSM 3051 to P. halophila, both of which had been isolated from the Great Salt Lake in Utah [47, 49]. It was concluded that type species Halovibrio variabilis should be associated with type strain DSM 3050 [49]. Thus,

MLA3 is identified as the strain of *Halovibrio variabilis* on the basis of percentage identity of 16S rRNA gene. The phylogenetic tree constructed to determine the affiliation of the strain MLA3 are shown in Fig.3. The investigated strain MLA3 falls in the cluster in which type strain of *Halovibrio variabilis and H. denitrificans* HGD3 is included with support of significant bootstraps. The occurrence of the studied strains is an addition of the genus *Halovibrio* in microbial diversity of Sambhar Lake. This cluster also comprised strains related to *Pseudomonas halophila*, those may be affiliated to genus *Halovibrio*. Moreover, phylogenetically close species of the genus *Halomonas* have been reported from Sambhar Lake [28].





#### Production of extracellular hydrolytic enzymes

Halovibrio variabilis MLA3 was able to produce the protease, amylase and lipase (fig. 4, fig. 5 and fig. 6). Whereas, type strain Halovibrio variabilis DSM 3050 showed production of protease alone confirmed by hydrolysis of casein [47, 50]. In the present study, quantitative assay was further performed for amylase and protease enzyme production on the basis of maximum hydrolyse zone. The maximum protease production  $1.7 \pm 0.02$  U/ml was observed after 7 days of incubation (fig.7). Our result corresponds to earlier findings showing that halophilic bacteria are well known for the protease enzyme production [19, 22, 51]. However, Halomonas variabilis SL121 isolated from Sambhar Lake showed only amylase enzyme production

and protease production was not reported by this strain. The present study describes the production of protease enzyme by the strain MLA3 at qualitative as well quantitative level. The amylase production started after 1 day and maximum production was recorded on  $7^{th}$  day (7.3 ± 0.05 U/ml) and later on production of enzyme decreased (fig. 8). Some other studies also showed amylase production by halophilic bacteria such as *Halomonas meridian* [52], *Bacillus* sp. Strain TSCVKK [53], *Bacillus cereus* [54]. Further, study will be focused on the optimization of carbon sources, nitrogen sources (organic and inorganic), effect of different factors such as agitation, inoculum size, inoculum age, metal ions, surfactant, additives, inhibitors and chelating agents on amylase production and its activity.






Fig 4. Skim milk hydrolysis (a) control (b) Halovibrio sp. strain MLA3




Fig 5. Starch hydrolysis (a) control (b) Halovibrio sp. strain MLA3




Fig. 6. Tween 80 hydrolysis (a) control (b) Halovibrio sp. strain MLA3




Fig. 7. Protease production during growth of *Halovibrio* sp. strain MLA3. Each value is a mean of three replicates data with ± standard deviation.




Fig. 8. Amylase production during growth of *Halovibrio* sp. strain MLA3. Each value is a mean of three replicates data with ± standard deviation.

#### 4. CONCLUSIONS

The present study exhibited the occurrence of bacteria of genus *Halovibrio* in Sambhar Lake, India. Moreover, the strain MLA3 found ability to produce industrially important enzymes such as amylase, protease and lipase which could meet the industrial needs. Thus, the present study shows that Sambhar lake is a treasure of

many unexplored microorganisms. It should be explored extensively for evaluation of microorganisms for phylogenetic studies and also for their biotechnological applications.



#### 5. ACKNOWLEDGEMENTS

The authors would like to thank Department of Microbiology, Central University of Rajasthan for providing the basic infrastructural facilities required for the present work. DKN is thankful to University Grant Commission, New Delhi and AKT to Central University of Rajasthan for providing financial support in the form of fellowships.

#### 7. REFERENCES

- 1. DasSarma S., DasSarma P., Halophiles. eLS: (2006)
- 2. Oren A., Life at high salt concentrations. In the Prokaryotes. Springer; 421-440, (2013)
- Ma Y., Galinski E.A., Grant W.D., Oren A., Ventosa A., Halophiles 2010: life in saline environments. Appl Environ Microbiol, 76 (21): 6971-6981, (2010)
- Ventosa A., Nieto J.J., Oren A., Biology of moderately halophilic aerobic bacteria. Microbiol Mol Biol Rev, 62 (2): 504-544, (1998)
- Rohban R., Amoozegar M.A., Ventosa A., Screening and isolation of halophilic bacteria producing extracellular hydrolyses from Howz Soltan Lake, Iran. J Ind Microbiol Biotechnol, 36 (3): 333-340, (2009)
- Oremland R.S., A random biogeochemical walk into three soda lakes of the western USA: With an introduction to a few of their microbial denizens. InPolyextremophiles. Springer; 179-199, (2013)
- Oren A., Two centuries of microbiological research in the Wadi Natrun, Egypt: a model system for the study of the ecology, physiology, and taxonomy of haloalkaliphilic microorganisms. In Polyextremophiles. Springer; 101-119, (2013)
- Kjeldsen K.U., Loy A., Jakobsen T.F., Thomsen T.R., Wagner M., Ingvorsen K., Diversity of sulfate-reducing bacteria from an extreme hypersaline sediment, Great Salt Lake (Utah). FEMS Microbiol Ecol, 60 (2): 287-298, (2007)
- Humayoun S.B., Bano N., Hollibaugh J.T., Depth distribution of microbial diversity in Mono Lake, a meromictic soda lake in California. Appl Environ Microbiol, 69 (2): 1030-1042, (2003)
- Joshi A.A., Kanekar P.P., Kelkar A.S., Shouche Y.S., Vani A.A., Borgave S.B., Sarnaik S.S., Cultivable bacterial diversity of alkaline Lonar Lake, India. Microb Ecol, 55 (2): 163-172, (2008)
- Upasani V.N., Microbiological Studies on Sambhar Lake (Salt of Earth) Rajasthan, India. In Proceedings of Taal2007: The 12th World Lake Conference. 2008: 450.
- 12. Upasani V., Desai S., Sambhar Salt Lake. Arch Microbiol, 154 (6): 589-593, (1990)
- 13. Upasani V.N., Desai S.G., Moldoveanu N., Kates M., Lipids of extremely halophilic archaeobacteria from

saline environments in India: a novel glycolipid in *Natronobacterium* strains. Microbiology, 140 (8): 1959-1966, (1994)

- Sundaresan S., Ponnuchamy K., Rahaman A.A., Biological management of Sambhar lake saltworks (Rajasthan, India). In Proceedings of the 1st International Conference on the Ecological Importance of Solar Saltworks, 20-22 October 2006. Santorini; 2006: 199-208.
- Giridhar P.V., Chandra T., Production of novel haloalkali-thermo-stable xylanase by a newly isolated moderately halophilic and alkali-tolerant *Gracilibacillus* sp. TSCPVG. Process Biochem, 45 (10): 1730-1737, (2010)
- Karan R., Khare S., Purification and characterization of a solvent-stable protease from *Geomicrobium* sp. EMB2. Environ Technol, 31 (10): 1061-1072, (2010)
- Rawal C., Raval V., Bhimani H., Bhensdadia D., Kothari C., Patel A., Bhatt V., Parmar N., Sajnani M., Koringa P., Whole-genome shotgun sequencing of the extremophile *Alkalibacillus haloalkaliphilus* C-5, of Indian origin. J Bacteriol, 194 (17): 4775-4775, (2012)
- Ramadoss D., Lakkineni V.K., Bose P., Ali S., Annapurna K., Mitigation of salt stress in wheat seedlings by halotolerant bacteria isolated from saline habitats. SpringerPlus, 2 (1): 6, (2013)
- Gupta M., Aggarwal S., Navani N.K., Choudhury B., Isolation and characterization of a protease-producing novel haloalkaliphilic bacterium *Halobiforma* sp. strain BNMIITR from Sambhar lake in Rajasthan, India. Ann Microbiol, 65 (2): 677-686, (2015)
- Gaur A., Prasad A., Parveen A., Oceanobacillus Sp. Moderate Halophilic Bacterium Isolated from Sambhar Salt Lake. Int J Pharm Biol Sci, 11 (2): 13, (2017)
- 21. Singh N., Choudhury B., Potential of *Lentibacillus* sp. NS12IITR for production of lipids with enriched branched-chain fatty acids for improving biodiesel properties along with hydrocarbon co-production. Extremophiles: 1-11, (2018)
- Karan R., Singh S., Kapoor S., Khare S., A novel organic solvent tolerant protease from a newly isolated *Geomicrobium* sp. EMB2 (MTCC 10310): production optimization by response surface methodology. N Biotechnol, 28 (2): 136-145, (2011)
- Nigam V., Singhal P., Vidyarthi A., Mohan M., Ghosh P., Studies on keratinolytic activity of alkaline proteases from halophilic bacteria. Int J Pharm Biol Sci, 4: 389-399, (2013)
- Gaur A., Sharma G., Mohan D., Isolation and characterization of moderate halophilic salinicoccus sp. Producing extracellular hydrolytic enzymes. J Glo Bio, 4 (9): 3379-3383, (2015)
- 25. Vijay A., Arora S., Gupta S., Chhabra M., Halophilic starch degrading bacteria isolated from Sambhar Lake, India, as



potential anode catalyst in microbial fuel cell: A promising process for saline water treatment. Bioresour Technol, 256: 391-398, (2018)

- Dobson S., McMeekin T., Franzmann P., Phylogenetic relationships between some members of the genera *Deleya*, *Halomonas*, and *Halovibrio*. Int J Syst Evol Microbiol, 43 (4): 665-673, (1993)
- Tiwari S., Singh P., Tiwari R., Meena K.K., Yandigeri M., Singh D.P., Arora D.K., Salt-tolerant rhizobacteriamediated induced tolerance in wheat (*Triticum aestivum*) and chemical diversity in rhizosphere enhance plant growth. Biol Fertility Soils, 47 (8): 907, (2011)
- Sahay H., Mahfooz S., Singh A.K., Singh S., Kaushik R., Saxena A.K., Arora D.K., Exploration and characterization of agriculturally and industrially important haloalkaliphilic bacteria from environmental samples of hypersaline Sambhar lake, India. World J Microbiol Biotechnol, 28 (11): 3207-3217, (2012)
- Makhdoumi-Kakhki A., Amoozegar M.A., Kazemi B., PaiC L., Ventosa A., Prokaryotic diversity in Aran-Bidgol Salt Lake, the largest hypersaline playa in Iran. Microbes Environ, 27 (1): 87-93, (2012)
- Amoozegar M.A., Fatemi A.Z., Karbalaei-Heidari H.R., Razavi M.R., Production of an extracellular alkaline metalloprotease from a newly isolated, moderately halophile, *Salinivibrio* sp. strain AF-2004. Microbiol Res, 162 (4): 369-377, (2007)
- Gram C., The differential staining of Schizomycetes in tissue sections and in dried preparations. Fortschitte der Medicin, 2: 185-189, (1884)
- Ventosa A., Quesada E., Rodriguez-Valera F., Ruiz-Berraquero F., Ramos-Cormenzana A., Numerical taxonomy of moderately halophilic Gram-negative rods. Microbiology, 128 (9): 1959-1968, (1982)
- León M.J., Fernández A.B., Ghai R., Sánchez-Porro C., Rodriguez-Valera F., Ventosa A., From metagenomics to pure culture: isolation and characterization of the moderately halophilic bacterium *Spiribacter salinus* gen. nov., sp. nov. Appl Environ Microbiol: AEM. 00430-00414, (2014)
- Gutiérrez C., González C., Method for simultaneous detection of proteinase and esterase activities in extremely halophilic bacteria. Appl Microbiol, 24 (3): 516, (1972)
- 35. brahim A.S., Al-Salamah A.A., Elbadawi Y.B., El-Tayeb M.A., Ibrahim S.S.S., Production of extracellular alkaline protease by new halotolerant alkaliphilic *Bacillus* sp. NPST-AK15 isolated from hyper saline soda lakes. Electron J Biotechnol, 18 (3): 236-243, (2015)
- Patel R., Dodia M., Singh S.P., Extracellular alkaline protease from a newly isolated haloalkaliphilic *Bacillus* sp.: Production and optimization. Process Biochem, 40 (11): 3569-3575, (2005)

- Karbalaei-Heidari H.R., Amoozegar M.A., Hajighasemi M., Ziaee A.-A., Ventosa A., Production, optimization and purification of a novel extracellular protease from the moderately halophilic bacterium *Halobacillus karajensis*. J Ind Microbiol Biotechnol, 36 (1): 21-27, (2009)
- Sahoo S., Roy S., Maiti S., A High Salt Stable α-Amylase by *Bacillus* Sp. MRS6 Isolated from Municipal Solid Waste; Purification, Characterization and Solid-State Fermentation. Enz Eng, 5 (3): 1-8, (2016)
- Mesbah N.M., Wiegel J., Halophilic alkali-and thermostable amylase from a novel polyextremophilic *Amphibacillus* sp. NM-Ra2. Int J Biol Macromol, 70: 222-229, (2014)
- Miller G.L., Use of dinitrosalicylic acid reagent for determination of reducing sugar. Anal Chem, 31 (3): 426-428, (1959)
- Sharma A., Satyanarayana T., High maltose-forming, Ca<sup>2+</sup>-independent and acid stable α-amylase from a novel acidophilic bacterium, *Bacillus acidicola*. Biotechnol Lett, 32 (10): 1503-1507, (2010)
- Barbier G., Godfroy A., Meunier J.-R., Quérellou J., Cambon M.-A., Lesongeur F., Grimont P.A., Raguénès G., *Pyrococcus glycovorans* sp. nov., a hyperthermophilic archaeon isolated from the East Pacific Rise. Int J Syst Evol Microbiol, 49 (4): 1829-1837, (1999)
- Chen X., Du X., Wang W., Zhang J., Sun Z., Liu W., Li L., Sun T., Zhang H., Isolation and identification of cultivable lactic acid bacteria in traditional fermented milk of Tibet in China. Int J Dairy Technol, 63 (3): 437-444, (2010)
- 44. Kim O.-S., Cho Y.-J., Lee K., Yoon S.-H., Kim M., Na H., Park S.-C., Jeon Y.S., Lee J.-H., Yi H., Introducing EzTaxone: a prokaryotic 16S rRNA gene sequence database with phylotypes that represent uncultured species. Int J Syst Evol Microbiol, 62 (3): 716-721, (2012)
- Edgar R.C., MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res, 32 (5): 1792-1797, (2004)
- Kumar S., Stecher G., Tamura K., MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. Mol Biol Evol, 33: 1870-1874, (2016)
- Fendrich C., Halovibrio variabilis gen. nov. sp. nov., *Pseudomonas halophila* sp. nov. and a new halophilic aerobic coccoid Eubacterium from Great Salt Lake, Utah, USA. Syst Appl Microbiol, 11 (1): 36-43, (1988)
- 48. Sorokin D.Y., Tourova T., Galinski E., Belloch C., Tindall B., Extremely halophilic denitrifying bacteria from hypersaline inland lakes, *Halovibrio denitrificans* sp. nov. and *Halospina denitrificans* gen. nov., sp. nov., and evidence that the genus name *Halovibrio* Fendrich 1989 with the type species *Halovibrio variabilis* should be associated with DSM 3050. Int J Syst Evol Microbiol, 56 (2): 379-388, (2006)



- Sorokin D.Y., Tindall B., The status of the genus name Halovibrio Fendrich 1989 and the identity of the strains Pseudomonas halophila DSM 3050 and Halomonas variabilis DSM 3051. Request for an opinion. Int J Syst Evol Microbiol, 56 (2): 487-489, (2006)
- Edbeib M.F., Wahab R.A., Kaya Y., Huyop F., In silico characterization of a novel dehalogenase (DehHX) from the halophile *Pseudomonas halophila* HX isolated from Tuz Gölü Lake, Turkey: insights into a hypersalineadapted dehalogenase. Ann Microbiol, 67 (5): 371-382, (2017)
- Sa A., Singh A., Garg S., Kumar A., Kumar H., Screening, Isolation and Characterization of Protease Producing Moderately Halophilic Microorganisms. Asian Jr. of Microbiol. Biotech. Env. Sc., 14 (4): 603-612, (2012)

Received:05.08.18, Accepted: 08.09.18, Published:01.10.2018

- 52. Coronado M.-J., Vargas C., Hofemeister J., Ventosa A., Nieto J.J., Production and biochemical characterization of an α-amylase from the moderate halophile *Halomonas meridiana*. FEMS Microbiol Lett, 183 (1): 67-71, (2000)
- Kiran K.K., Chandra T., Production of surfactant and detergent-stable, halophilic, and alkalitolerant alphaamylase by a moderately halophilic *Bacillus* sp. strain TSCVKK. Appl Microbiol Biotechnol, 77 (5): 1023-1031, (2008)
- 54. Vijayabaskar P., Jayalakshmi D., Shankar T., Amylase production by moderately halophilic *Bacillus cereus* in solid state fermentation. Afr J Microbiol Res, 6 (23): 4918-4926, (2012)

### \*Corresponding Author:

Pawan K Dadheech\* Email: pdadheech@curaj.ac.in