



Study on Biofilm Forming Ability of *Staphylococcus* Spp. Isolated from Bovine Mastitis Milk

Khushal Patel¹, Palak Rahevar¹ and Anju Kunjadiya^{2*}

¹Ashok & Rita Patel Institute of Integrated Study & Research in Biotechnology and Allied Sciences, New Vallabh Vidyanagar-388121, Gujarat, India

²Centre for Interdisciplinary Studies in Science and Technology (CISST), Sardar Patel University, V. V. Nagar-388120, Anand, Gujarat, India

Received: 20 Oct 2018 / Accepted: 12 Nov 2018 / Published online: 1 Jan 2019

Corresponding Author Email: anjukunjadiya@gmail.com

Abstract

Mastitis is an inflammation of mammary gland and udder tissue. One of the vital etiological agent *Staphylococcus* sp., mainly *Staphylococcus aureus* is responsible. Biofilm forming ability of such microorganism is considered as one of the vital virulence factors. This study was designed to screen the biofilm forming ability of mastitis-causing pathogens. A total of 48 *Staphylococcus* spp. was investigated which were isolated from mastitis infected milk samples using Baird parker and Mannitol salt differential media having drug resistivity observed from previous study. Tube assay and Microtitre plate assay were conducted for the detection of biofilm for these antibiotic-resistant strains. In tube assay detection method, 19 strong biofilm formers containing all *S. aureus*; 19 moderate formers containing 12 *S. aureus* & 7 *Staphylococcus* spp. and 10 weak biofilm formers containing all *Staphylococcus* spp. were observed at 24 hr. and after additional incubation of 12 hr leads all the isolates to strong biofilm forming category. Similarly, in microtitre plate assay out of 31 Coagulase positive and methicillin-resistant *Staphylococcus aureus*, 9 were moderate and 22 were strong biofilm formers, while out of studied 17 *Staphylococcus* spp., 9 were observed weak and 8 were moderate biofilm formers. No significance difference was observed after more incubation period i.e. 12-24 hr. in microtitre plate assay. Tube assay can be considered as the best method for preliminary detection of biofilm as it can be used to study the growth pattern of microorganism and may help in diagnosis and treatment of such infections caused by pathogens.

Keywords

Biofilm, ELISA, Microtitre, *Staphylococcus* spp.

INTRODUCTION

Mastitis in bovines known as an inflammation of mammary glands [20]. Many etiological agents are responsible to cause this clinical condition having an

ability to form a biofilm. Biofilms are well-defined as the group of bacteria enclosed in a self-synthesized extracellular polymeric matrix (EPM), which attaches to an abiotic or a biotic surface [22].

Biofilm forming ability of a microorganism is considered to be one of the notable virulent factors. *Staphylococcus aureus* and *Staphylococcus epidermidis* are recognised to be vital bacteria for causing the clinical infection [5].

Such infections involving biofilm-forming bacteria are extremely difficult to eliminate because biofilms avenge antibiotic dissemination and it also prevents typical immune responses against infections [3]. The presence of biofilm formation is considered to be an important feature for survival and virulence. It has been identified that resistant nature of *S. aureus* against antibiotics mainly methicillin are due to biofilms. Some studies have shown that the difficulty to treat staphylococcal infection using conventional antibiotics is very difficult [17].

Different pathways and time ranges are observed in the formation of biofilms and sometimes its detection may vary according to the time taken for formation. Time is a dependable parameter for biofilm formation. The time required for biofilm formation is depending upon bacterial strains and it should be taken into consideration for the development of suitable antibiotic rehabilitation measures [7, 14].

Detection of biofilm forming ability of a bacteria at primary level is necessary. Few studies addressed that microtiter plate assay can be used as a rapid and simple method to screen biofilm formation of strains [2, 16].

Here, we studied the biofilm-forming ability of coagulase positive *Staphylococcus aureus* (CoPSA) and coagulase negative *Staphylococcus* spp. (CoNS) of clinical mastitic infection using two different approaches, one is simple tube assay for qualitative and microtitre plate assay to quantitative detection.

MATERIAL AND METHOD

Bacterial strains

We studied biofilm formation assay on 48 strains of *Staphylococcus* sp. Out of which 31 were coagulase positive *Staphylococcus aureus* which was resistant to methicillin and remained 17 *Staphylococcus* spp. were multidrug resistant including methicillin from the previous study. Isolates were confirmed and antibiotic sensitivity assay was performed for all the isolates in a previous study [15].

Tube assay

Identification of biofilm formation qualitatively was performed by tube assay method as described. Briefly, 1.9 ml Luria-Bertani (HiMedia, M1245) broth was inoculated with 100 μ l of overnight grown culture broth and incubated for 24 hr. 48 hr. and 72

hr. at 37°C with control without culture [11]. The tubes were decanted and washed with Phosphate Buffer Saline (PBS) (Sigma Aldrich, MFC00131855) at pH 7.3 and tubes were subjected to dry. Staining of dried tubes was done with 0.1 % crystal violet stain (Sigma-Aldrich, C0775). Excess stain was removed by washing the tubes with deionized water (Sigma-Aldrich, 99053). Later, tubes were dried in an inverted position and observed for biofilm formation.

Microtitre plate assay

The isolates were screened for their capability to form biofilm by microtitre plate method with modification with culture dilutions and incubation time [7, 13, 21]. Isolates from fresh nutrient agar plates were inoculated in LB broth supplemented with 1% glucose and incubated for 18 hours at 37°C in a stagnant condition. Sterile flat bottom 96 wells plate (HiMedia, DM3) was filled with 0.2 ml aliquots of the diluted overnight cultures (Dilution Factor 10^{-1}). The control well contained only LB broth without inoculation. The plates were incubated for 24 hr and 48 hr at 37°C to observe biofilm.

After the respective incubation period, the content of each well was blandly removed by lightly tapping / instant inverting the plates. The wells were washed twice with phosphate buffer saline pH 7.3 to remove free-floating planktonic bacteria. The plates were then stained with 0.1% crystal violet solution and washed. Excess stain was washed off completely with 70% ethanol and plates were kept for drying. In order to quantify the adhered cells, 220 μ l of decolouring solution (ethanol/acetone, 80:20%) was added to each well for 15 min. The absorption of the eluted stain was measured with a micro ELISA auto reader at a wavelength of 630 nm. These values were considered as an evidence of attachment to surface and forming biofilms. The experiment was performed in triplicates with proper hygiene conditions and the mean optical density values were considered.

Statistical Analysis

Biofilm forming ability was evaluated for coagulase-positive *S. aureus*(CoPSA) and coagulase negative *Staphylococcus* sp. (CoNS) using Friedman's test (SPSS 20) including 24 hr. and 48 hr. data of microtitre plate assay [23].

RESULTS AND DISCUSSION

Formation of biofilm was confirmed with the presence of attachment on the wall and bottom of the tube as shown in figure 1 [7].



Figure 1. Biofilm attached to the bottom of the tube

As a result of tube assay, positive biofilm formation was considered when a visible stained film was observed on the wall and bottom of the tube. Ring formation at the liquid interface was not considered an indication of biofilm formation. Tubes were examined and the amount of biofilm formation was scored as negative (-), weak (+), moderate (+++) and strong (++++).

Tube assay was performed on 48 isolates, out of which 19 *S. aureus* showed attachment of biofilm at the bottom and wall of the tube when stained with crystal violet after 24 hr. and remaining 29 isolates showed attachment after 24-36 hr (Table 1).

Table 1. Biofilm formation by Tube assay after 24 hr. & 24-36 hr. incubation

Biofilm Formation	Isolates showing biofilm after 24 hr. incubation	Isolates showing biofilm after 24-36 hr. incubation
Negative (-)	Control	Control
Weak formation (+)	10 (<i>Staphylococcus</i> spp.)	-
Moderate formation (+++)	19 (12 <i>S. aureus</i> ; 07 <i>Staphylococcus</i> spp.)	-
Maximum formation (++++)	19 (<i>S. aureus</i>)	29 (12 <i>S. aureus</i> , 17 <i>Staphylococcus</i> spp.)

Microtitre plate assay gave promising and comparatively similar results to a preliminary evaluation of tube assay. Microtitre plate assay is an updated method of tube assay in a smaller scale with the larger number of samples. In this method categorization of biofilm was done based on absorbance value. On the basis of microtitre plate assay, biofilm formation ability was weak for 9

Staphylococcus spp. isolate (OD <0.120), moderate for 17 isolates (9 MRSA, 8 *Staphylococcus* spp. where OD range was between 0.120-0.240 and high for 22 (*S. aureus*) isolates having absorbance above 0.240. The results at the point justifies tube staining outcomes (Table 2; Figure 1). After 12 hr. to 24 hr. of absorbance measurement; negligible change in OD value was observed.

Table 2. Biofilm Categorization on the basis of microtitre plate assay

Absorbance at 630 nm	Biofilm Formation
<0.120	Non/Weak
0.120 – 0.240	Moderate
>0.240	High




Figure 2: Numbers of isolates showed biofilms in category

According to the Friedman's test, no significant difference was observed between groups (CoPSA and CoNS for time intervals) in terms of biofilm-forming ability at each time point (24 hr: $p = 0.027$; 48 hr: $p = 0.040$) and verified no difference in outputs after 24 hr supplementary incubation.

When free whirling microbial cells encounter any surface, normally they switch themselves from a planktonic life type to the slimy communities which are known as biofilms that are enclosed by a dense layer of the self-produced slippery matrix material [12].

The ability of a pathogenic organism to form biofilm is a trait that is diligently associated with bacteria's virulence and persistence. Thus, many persistent and chronic bacterial infections like mastitis are nowadays believed to be linked with the formation of biofilm by pathogens [4, 9].

In this study, we analysed the ability of multi drug resistant *Staphylococcus* spp. to form a biofilm. *S. aureus* is a pliable opportunistic pathogen whose ability to persist and multiply in a variety of environments causes a wide spectrum of diseases in animals. This may lead to an increase in the risk of animal health. Biofilm forming ability of *Staphylococcus* sp. from mastitis cases is nowadays a major concern pertaining to animal health and milk quality.

A study from microtitre plate assay showed all the 31 *S. aureus* strains were found biofilm producers, classified as strong (70.97% = 22/31), moderate (29.03% = 09/31), and in *Staphylococcus* spp. 52.94% (09/17) were weak and 47.04% (08/17) were moderate biofilm producers. These data are in

accordance with the tube assay outputs where all the isolates were found biofilm producers.

It is very notable observation that all the 48 (100%) *Staphylococcus* spp. were biofilm producer as well as antibiotics resistant. This virulent factor assures the pathogenicity of *Staphylococcus* spp. and its connexion and conservation in the mammary gland tissue of udder. The results of this study are in accordance with a study from China, USA, Taiwan and India [4, 10, 12, 13, 21].

For detection of biofilm Congo Red Agar method is also used for primary detection but according to a study by T. Mathur *et al.*, it suggests that tube assay method for detection of biofilm forming ability of an organism is accurate and reproducible for screening [13].

Prevalence of biofilm producing ability of *Staphylococcus* spp. in different colonizing conditions may defer. It is highly depending upon the in-vitro conditions. In many cases, simple bacteria are unable to produce such biofilm until and unless it is provided with the susceptible environment to enhance toxins within and that is directly affecting the strain health. It has been previously demonstrated that phenotypic expression of biofilm formation is highly susceptible to in vitro conditions and hence can be detected variably by different methods [6, 7, 23].

This study was designed on single species biofilms and since beginning single-species biofilms have been extensively investigated. However, the biofilms in nature comprises mostly multiple species whether from same genera or different, where interspecies interactions help in shaping development, structure

and molecular functions from biofilm populations *S. aureus* [1, 8, 18, 19].

CONCLUSION

Detection of biofilm by the mainstream producers *Staphylococcus aureus* and *Staphylococcus* spp. can be detected in 24-36 hr. cultures. Tube assay and microtitre plate assays are the best approaches for preliminary screening qualitatively and quantitatively. The understanding of time progression of biofilm expression may subsidize the microbiologist to improve the treatment strategies with the use of antimicrobials for Staphylococcal infected mastitis therapy because it's an important virulent factor of microorganism.

ACKNOWLEDGEMENT

Authors are gratefully thanking the substantial support of CISST, Sardar Patel University; acknowledge SICART, Charutar Vidya Mandal for providing support and platform to conduct experiments.

ETHICAL APPROVAL

No need for ethical approval.

FUNDING

The authors received no financial support for the research.

CONFLICT OF INTEREST

The authors declare that they have no competing interests.

REFERENCE

- Al-rubaye SMH, Al-jumaily EF, Abdul-ratha HA (2016) Biofilm Production by *Staphylococcus aureus* isolated from Bovine Mastitis Related with Resistance to the Antibiotics. International Journal of Current Microbiology and Applied Sciences 5:33–44
- Barakat GI, Nabil YM (2016) Correlation of mupirocin resistance with biofilm production in methicillin-resistant *Staphylococcus aureus* from surgical site infections in a tertiary centre, Egypt. Journal of Global Antimicrobial Resistance 4:16–20. doi: 10.1016/j.jgar.2015.11.010
- Brackman G, Breyne K, De Rycke R, Vermote A, Van Nieuwerburgh F, Meyer E, Van Calenbergh S, Coenye T (2016) The Quorum Sensing Inhibitor Hamamelitannin Increases Antibiotic Susceptibility of *Staphylococcus aureus* Biofilms by Affecting Peptidoglycan Biosynthesis and eDNA Release. Scientific reports 6:20321. doi: 10.1038/srep20321
- Cha JO, Yoo JI, Yoo JS, Chung HS, Park SH, Kim HS, Lee YS, Chung GT (2013) Investigation of Biofilm Formation and its Association with the Molecular and Clinical Characteristics of Methicillin-resistant *Staphylococcus aureus*. Osong Public Health Res Perspect 4:225–232. doi: 10.1016/j.phrp.2013.09.001
- Chavhan SK, Kalorey DR, Nagdive AA, Purohit HJ, Barbudhe SB, Kurkure NV (2012) Molecular characterization of intercellular adhesion gene in *Staphylococcus aureus* isolated from bovine mastitic milk. Tropical Animal Health and Production 44:247–252. doi: 10.1007/s11250-011-0009-9
- Da Cunha MDLRDS, Peresi E, Oliveira Calsolari RA, Araújo JP (2006) Detection of enterotoxins genes in coagulase-negative staphylococci isolated from foods. Brazilian Journal of Microbiology 37:70–74. doi: 10.1590/S1517-83822006000100013
- Dhanawade NB, Kalorey DR, Srinivasan R, Barbudhe SB, Kurkure N V. (2010) Detection of intercellular adhesion genes and biofilm production in *Staphylococcus aureus* isolated from bovine subclinical mastitis. Veterinary Research Communications 34:81–89. doi: 10.1007/s11259-009-9326-0
- Fey PD, Olson ME (2010) Current concepts in biofilm formation of *Staphylococcus epidermidis*. FutureMicrobiol 5:917–933. doi: 10.2217/fmb.10.56.Current
- Fox LK, Zadoks RN, Gaskins CT (2005) Biofilm production by *Staphylococcus aureus* associated with intramammary infection. Veterinary Microbiology 107:295–299. doi: 10.1016/j.vetmic.2005.02.005
- Jain A, Agarwal A (2009) Biofilm production, a marker of pathogenic potential of colonizing and commensal staphylococci. Journal of Microbiological Methods 76:88–92. doi: 10.1016/j.mimet.2008.09.017
- Lotfi G, Hafida H, Nihel K, Abdelmonaim K, Fatima NAS, Walter Z (2014) Detection of biofilm formation of a collection of fifty strains of *Staphylococcus aureus* isolated in Algeria at the University Hospital of Tlemcen. Journal of Bacteriology Research 6:1–6. doi: 10.5897/JBR2013.0122
- Marques VF, Motta CC da, Soares B da S, Melo DA de, Coelho S de M de O, Coelho I da S, Barbosa HS, Souza MMS de (2017) Biofilm production and beta-lactamic resistance in Brazilian *Staphylococcus aureus* isolates from bovine mastitis. Brazilian Journal of Microbiology 48:118–124. doi: 10.1016/j.bjm.2016.10.001
- Mathur T, Singhal S, Khan S, Upadhyay D, Fatma T, Rattan A (2006) Detection of biofilm formation among the clinical isolates of Staphylococci: An evaluation of three different screening methods. Indian Journal of Medical Microbiology 24:25. doi: 10.4103/0255-0857.19890
- Mirani ZA, Aziz M, Khan MN, Lal I, Hassan N ul, Khan SI (2013) Biofilm formation and dispersal of *Staphylococcus aureus* under the influence of oxacillin. Microbial Pathogenesis 61–62:66–72. doi: 10.1016/j.micpath.2013.05.002
- Patel K, Joshi C, Kunjadiya A (2017) Isolation and Molecular Identification of *Staphylococcus* spp. from

- Bovine Mastitis Milk Samples. International Journal of Advanced Research 5:2320–5407. doi: 10.21474/IJAR01/4818
16. Purkrtova S, Babulikova J, Karpiskova R, Demnerova K, Pazlarova J (2011) Antimicrobial Factors Effects on Biofilm Formation in *Staphylococcus aureus*. Czech Journal of Food Sciences 29:S1–S10
 17. Qayyum A, Khan JA, Hussain R, Khan A, Avais M, Ahmad N, Hassan MF (2016) Molecular Characterization Of *Staphylococcus aureus* Isolates Recovered From Natural Cases Of Subclinical Mastitis In Cholistani Cattle And Their Antibacterial Susceptibility. 53:971–976
 18. Raza A, Muhammad G, Sharif S, Atta A (2013) Biofilm Producing *Staphylococcus aureus* and Bovine Mastitis: A Review. Molecular Microbiology Research. doi: 10.5376/mmr.2013.03.0001
 19. Silva NCC, Guimarães FF, Manzi MDP, Gómez-Sanz E, Gómez P, Araújo JP, Langoni H, Rall VLM, Torres C (2014) Characterization of methicillin-resistant coagulase-negative staphylococci in milk from cows with mastitis in Brazil. Antonie van Leeuwenhoek, International Journal of General and Molecular Microbiology 106:227–233. doi: 10.1007/s10482-014-0185-5
 20. Taverna F, Negri A, Piccinini R, Zecconi A, Nonnis S, Ronchi S, Tedeschi G (2007) Characterization of cell wall associated proteins of a *Staphylococcus aureus* isolated from bovine mastitis case by a proteomic approach. 119:240–247. doi: 10.1016/j.vetmic.2006.09.007
 21. Torlak E, Korkut E, Uncu AT, Şener Y (2017) Biofilm formation by *Staphylococcus aureus* isolates from a dental clinic in Konya, Turkey. Journal of Infection and Public Health 1–5. doi: 10.1016/j.jiph.2017.01.004
 22. Yarets Y, Rubanov L, Novikova I, Shevchenko N (2013) The Biofilm-forming capacity of *Staphylococcus aureus* from chronic wounds can be useful for determining Wound-Bed Preparation methods. EWMA Journal 13:7–13
 23. Zmantar T, Ben Slama R, Fdhila K, Kouidhi B, Bakhrouf A, Chaieb K (2017) Modulation of drug resistance and biofilm formation of *Staphylococcus aureus* isolated from the oral cavity of Tunisian children. The Brazilian Journal of Infectious Diseases 21:27–34. doi: 10.1016/j.bjid.2016.10.009