



SYNTHESIS, CHARACTERIZATION AND *IN VITRO* BIOLOGICAL STUDIES: Cu (II) COMPLEXES OF N-(2-HYDROXYBENZYL)-AMINO ACID LIGANDS AND 4, 4'-BIPYRIDINE

Santha Lakshmi S^{*1}, Saranya J² and Chithra J³

^{*1}Assistant professor, PG and Research Department of Chemistry, D.K.M College for Women (Autonomous), Vellore-632001.

^{2&3} Research Scholar, D.K.M College for Women (Autonomous), Vellore-632001.

*Corresponding Author Email: santhalakshmi76@gmail.com

ABSTRACT

Currently, coordination chemistry occupies a larger area of inorganic research. Copper (II) complexes derived from Schiff bases have attracted attention of the researchers for their preparative, accessibilities and structural varieties. Copper (II) complexes containing Schiff bases and heterocyclic compounds were reported to show very good biological activities. In the present work, a series of five Schiff base Cu (II) complexes derived from N-(salicylidene)-L-aminoacids [L-valine, L-tryptophan, L-leucine, L-alanine and L-cysteine] and 4,4'-bipyridine were designed, synthesized and tested for biological activities. All the complexes were characterized by physicochemical and spectral studies such as UV-Visible, FTIR, EPR, TGA and CV. In vitro antimicrobial studies were carried out against bacterial strains such as *S. aureus*, *E. coli* and *P. aeruginosa* and fungal strains such as *A. flavus*, *A. fumigates* and *Rhizopus*. In addition, in vitro larvicidal and antioxidant studies were also performed. The FTIR spectra confirmed that the Cu (II) ion is coordinated to imine nitrogen, phenolic oxygen and one of the oxygen anion of carboxylate group present in the Schiff base ligand and confirms that the Schiff base acted as a tridentate ligand. The cyclic voltammogram of all the complexes suggested two one electron transfer with quasi-reversible redox reactions. The biological activity data of the Cu (II) complex showed the better antimicrobial, antioxidant and larvicidal activity.

KEY WORDS

Schiff base Cu (II) complexes, spectral studies, in vitro antibacterial, antifungal, larvicidal and antioxidant studies.

INTRODUCTION:

Today metal complexes play an essential role in the drug industries and in agronomy. ^[1] In the case of metalloproteins, copper (II) complexes act as an active site and plays an important role in biological systems. ^[2] Recently, many researchers have been focusing their work on synthesis and characterisation of copper complexes due to their ease of preparation, structural varieties and biological applications. The Schiff base copper complexes have wide applications in many fields

such as food industry, dye industry, analytical chemistry, catalysis, agrochemical, anti-inflammable activity, antiradical activities and biological activities. ^[3-5] Further, Schiff base copper complexes are promising agents in pharmaceuticals. ^[6, 7] Schiff bases are considered to be privileged ligand in the field of Coordination chemistry ^[8] which was first reported by Hugo Schiff. ^[9-10] The nucleophilic attack of amines on electrophilic carbon of aldehydes or ketones end with the result of forming a compound in which C=O

double bond is replaced by a C=N double bond. This type of compound is known as imine, azomethine or a Schiff base. ^[11, 12] Schiff base ligand derived from salicylaldehyde is a significant component in enlightening the ideal coordination geometries of copper complexes and its derivatives, which are found to possess attractive biological activities. ^[13] Today, a number of diseases in the world such as malaria, dengue and filariasis are generated by Mosquitoes. To prevent mosquito borne diseases and to improve public health it is necessary to control the larvae of mosquitoes. Copper complexes have been shown to possess better larvicidal activity. ^[14,15] Till now, there is a lack in the preparations of Schiff base copper complexes as pharmaceuticals agents. Based on these backdrops, an attempt was made to synthesise Schiff base Cu (II) complexes and to investigate their biological properties by *in vitro* method.

In the present work, a series of Schiff base Cu (II) complexes were synthesised using N-(salicylidene)-L-aminoacids [L-valine, L-tryptophan, L-leucine, L-alanine and L-cysteine] and 4,4'-bipyridine. The synthesized Cu (II) complexes were characterised by physicochemical, spectral and *in vitro* antimicrobial, antioxidant and larvicidal studies.

MATERIALS AND METHODS:

All common laboratory chemicals and reagents used were of analytical grade. The electronic absorption spectra of the complexes were recorded on a SYSTRONICS 2201 spectrometer at room temperature with the sample concentration of (10^{-3} M) in DMSO. The FITR spectra were recorded on SHIMADZU spectrometer between 4000-400 cm^{-1} range, using KBr pellet. The EPR spectra were recorded in solid state at room temperature using Bruker EMX -10/2.7 spectrometer. The electrochemical activity of the Cu (II) complexes were studied in DMSO (10^{-3} M) on a cyclic voltammetry model CHI617C electrochemical analyser with the scan rate of 0.1 V/s in the potential range +2 to -2 V using TBAP (Tetrabutylammonium perchlorate) as supporting electrolyte. Thermal stability of complex **1** was carried out on the instrument SDT Q600 V20.9 Build 20 model from room temperature to 800 °C under a nitrogen atmosphere at the heating rate of 20°C/min.

Template synthesis of Schiff base metal complexes:

Schiff base Cu (II) complexes were synthesised as per the literature. ^[16] L-valine (0.351 g, 3 mmol) and KOH

(0.336 g, 6 mmol) were dissolved in water. To this an ethanolic solution (25 mL) of salicylaldehyde (0.3 mL, 3 mmol) was added. The reaction mixture was stirred for an hour at 60 °C. The solution turned yellow. Then, copper (II) acetate monohydrate (0.61 g, 3 mmol) was added and the reaction mixture was stirred for 2 h, followed by addition of 4, 4'- bipyridine (0.468 g, 3 mmol). The mixture was stirred for another 2 h at the same temperature. The resultant green coloured product was filtered, washed with ethanol and dried. Similar procedure was adopted for the preparation of the other Cu (II) complexes using the amino acids L-tryptophan (0.6126 g, 3 mmol), L-leucine (0.3935 g, 3 mmol), L-alanine (0.2672g, 3 mmol) and L-cysteine (0.36343 g, 3 mmol).

Antimicrobial studies (*in vitro*):

The synthesised Schiff base Cu (II) complexes were screened for their biological activities to test their bio-efficacy. ^[17] The copper complexes have been screened for their antimicrobial activities against the bacterial strains of *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *E. coli* and fungal strains of *Aspergillus flavus*, *Aspergillus fumigates* and *Rhizopus* using agar well diffusion method. 10^{-3} M test solution in DMSO was used for the study. Ampicillin and polymycin B sulphate were used as a standard for the study of antibacterial and antifungal activities respectively.

Antioxidant activity:

Hydrogen peroxide scavenging activity:

Hydrogen peroxide scavenging activity was carried out as per the literature. ^[18, 19] A solution of hydrogen peroxide (40 mM) was prepared in phosphate buffer (50 mM, pH 7.4). The concentration of hydrogen peroxide is determined by adsorption at 230 nm using a spectrophotometer. Synthesized complexes with the concentration of 2 mg /1 mL, 4 mg /1 mL, 6 mg /1 mL in DMF are added to hydrogen peroxide and the absorption at 230 nm was observed after 10 min against blank solution containing phosphate buffer without hydrogen peroxide. The percentage of hydrogen peroxide scavenging activity was calculated using the following equation. ^[19]

$$\% \text{ scavenged (H}_2\text{O}_2) = \frac{A_c - A_s}{A_c} \times 100$$

where, A_c is the absorbance of the control solution; A_s is the absorbance of the sample solution.

Larvicidal activity:

The eggs and egg rafts of *Culex quinque fasciatus* were procured from Zonal Entomological Unit, Vellore, Tamil

Nadu. *Culex quinque fasciatus* larvae were maintained in the laboratory as per the literature.^[20] The larvicidal activity was assessed by the procedure of WHO guidelines with some modification^[21] and number of dead larvae, percentage of mortality in each batch were counted for every 24 h exposure period by using various concentration (6 mg, 4 mg, 2 mg and 1mg) of synthesized Cu (II) complexes. The treated larvae were

mounted on a slide and examined under a microscope for image capture.

RESULTS:

The analytical data of the synthesized Cu (II) complexes such as molecular formula, molar conductance and colour are shown in Table 1.

Table 1. Analytical data of the Schiff base Cu (II) complexes

| Complex Formula | Complex code | Molecular formula | Molecular weight | Melting /Decomposition point °C | Colour | Molar Conductance Ohm ⁻¹ cm ² mol ⁻¹ |
|---|--------------|---|------------------|---------------------------------|-------------|---|
| [Cu ₂ L ₂ ¹ L ² (H ₂ O) ₄] | 1 | C ₃₄ H ₄₂ O ₁₀ N ₄ Cu ₂ | 794 | 204 | Dark green | 8.05 |
| [Cu ₂ L ₂ ³ L ² (H ₂ O) ₄] | 2 | C ₄₆ H ₄₄ O ₁₀ N ₆ Cu ₂ | 966 | 210 | Light green | 6.9 |
| [Cu ₂ L ₂ ⁴ L ² (H ₂ O) ₄] | 3 | C ₃₆ H ₄₆ O ₁₀ N ₄ Cu ₂ | 820 | 215 | Dark green | 5.75 |
| [Cu ₂ L ₂ ⁵ L ² (H ₂ O) ₄] | 4 | C ₃₀ H ₃₄ O ₁₀ N ₄ Cu ₂ | 736 | 150 | Green | 9.2 |
| [Cu ₂ L ₂ ⁶ L ² (H ₂ O) ₄] | 5 | C ₃₀ H ₃₄ O ₁₀ N ₄ S ₂ Cu ₂ | 800 | 219 | Green | 4.6 |

where, L₂¹-N- salicylidene-L-valine, L₂³-N- salicylidene-L-tryptophan, L₂⁴-N- salicylidene-L-leucine, L₂⁵-N- salicylidene-L-alanine, L₂⁶-N- salicylidene-L-cysteine, L²-4, 4'-bipyridine.

UV-Vis. spectra:

The UV-Visible spectral data are represented in the Table 2.

Table 2. The UV-Vis. spectral data of Schiff base Cu (II) complexes

| Complex code | Absorption (λ _{max} nm) | | |
|--------------|----------------------------------|------|-----|
| | π→π* | n→π* | d→d |
| 1 | 251 | 389 | 685 |
| 2 | 240 | 374 | 648 |
| 3 | 265 | 370 | 703 |
| 4 | 267 | 375 | 672 |
| 5 | 269 | 371 | 645 |

FTIR spectra:

The assignments of important infrared spectral data are listed in Table 3.

Table 3. Table 3. FTIR spectral data of the Schiff base Cu (II) complexes

| Complex code | C=N | COO ⁻ | | Δν = [ν _{as} - ν _s] | M-O | M-N |
|--------------|------|------------------|----------------|--|-----|-----|
| | | ν _{as} | ν _s | | | |
| 1 | 1584 | 1533 | 1317 | 216 | 463 | 546 |
| 2 | 1628 | 1593 | 1348 | 246 | 418 | 549 |
| 3 | 1632 | 1566 | 1340 | 226 | 480 | 544 |
| 4 | 1562 | 1470 | 1332 | 138 | 473 | 569 |
| 5 | 1593 | 1535 | 1389 | 146 | 472 | 538 |

EPR spectra:

The EPR spectrum is mainly used to analyse the paramagnetic nature and geometry of the metal complexes. The g-value of the Schiff base Cu (II) complexes are given in Table 4.

Table 4. EPR data for Schiff base Cu (II) complexes.

| Complex code | g value |
|--------------|---------|
| 1 | 2.03 |
| 2 | 2.01 |
| 3 | 2.09 |
| 4 | 2.04 |
| 5 | 2.06 |

Cyclic voltammetry:

The redox potentials of all the complexes of present series are summarised in Table 5.


Table 5. Redox potentials for the Schiff base Cu (II) complexes

| Complex code | Epc (V) | Epa (V) | ΔE_p (V) | $E_{1/2}$ (V) | Ipc μA | Ipa μA | Ipa/Ipc |
|--------------|---------|---------|------------------|---------------|-------------|-------------|---------|
| 1 | -1.650 | 0.180 | 1.830 | -0.735 | -0.510 | 0.595 | -1.166 |
| | -1.319 | 2.870 | 4.189 | 0.775 | -0.938 | 0.967 | -1.030 |
| 2 | -1.630 | 0.395 | 2.025 | -0.617 | -0.392 | 0.323 | -0.823 |
| | -1.296 | 0.847 | 2.143 | -0.224 | -0.843 | 0.919 | -1.090 |
| 3 | -0.741 | 0.348 | 1.089 | -0.196 | -0.750 | 0.739 | -0.985 |
| | -0.567 | 0.761 | 1.328 | 0.097 | -0.466 | 0.678 | -1.454 |
| 4 | -0.704 | 0.312 | -0.392 | -0.196 | -0.659 | 0.348 | -0.528 |
| | -0.475 | 0.733 | 1.208 | 0.129 | -0.503 | 0.770 | -1.530 |
| 5 | -0.768 | 0.352 | 1.120 | -0.208 | -0.757 | 0.396 | -0.523 |
| | -0.483 | 0.858 | 1.341 | 0.187 | -0.537 | 0.860 | -1.601 |

Epa = anodic peak potential; Epc = cathodic peak potential; Ipa = anodic peak current; Ipc = cathodic current peak; Ipa/Ipc = number of electrons; ΔE_p = Epa - Epc; scan rate = 0.1 V/s.

Thermogravimetric analysis:

TGA provides useful information about the thermal stability of the synthesized complexes. The TGA curve of the complex **1** is represented in Fig1.


Figure 1. TGA curve of complex 1
Antibacterial and antifungal activity:

The zone of inhibition values of the synthesized metal complexes against the growth of the bacteria and fungi under investigation are measured in mm. The



corresponding data are summarized in Tables 6 and 7 and the graphical representation is given in the Fig 2 & 3.

Table 6. Antibacterial activity of the Schiff base Cu (II) complexes

| Bacteria | Zone of Inhibition (mm) | | | | | Ampicillin |
|----------------------|-------------------------|-----------|-----------|-----------|-----------|------------|
| | Complex 1 | Complex 2 | Complex 3 | Complex 4 | Complex 5 | |
| <i>S. aureus</i> | 16 | 18 | 15 | 6 | 11 | 10 |
| <i>E. coli</i> | 15 | 13 | 19 | 19 | 12 | 11 |
| <i>P. aeruginosa</i> | 15 | 18 | 10 | 19 | 12 | 10 |

Table 7. Antifungal activity data of the Schiff base Cu (II) complexes

| Fungi | Zone of Inhibition (mm) | | | | | Gentamicin |
|---------------------|-------------------------|-----------|-----------|-----------|-----------|------------|
| | Complex 1 | Complex 2 | Complex 3 | Complex 4 | Complex 5 | |
| <i>A. flavus</i> | 15 | 12 | 23 | 16 | 20 | 20 |
| <i>A. fumigatus</i> | 21 | 19 | 25 | 15 | 24 | 15 |
| <i>Rhizopus</i> | 12 | 15 | 21 | 20 | 22 | 15 |


Figure 2. Graphical representation of antibacterial activity of the Schiff base Cu (II) complexes

Figure 3. Graphical representation of antifungal activity of the Schiff base Cu (II) complexes

Antioxidant activity:

The antioxidant scavenging activity of the synthesized Schiff base Cu (II) complexes in hydrogen peroxide (H₂O₂) was monitored and the values are listed in Table 8.

Table 8. Antioxidant scavenging activity values of Cu (II) complexes

| Complex code | % antioxidant scavenging activity | | |
|--------------|-----------------------------------|-------|-------|
| | 2 mg | 4 mg | 6 mg |
| 1 | 90.29 | 92.23 | 93.20 |
| 2 | 91.26 | 92.23 | 93.20 |
| 3 | 86.40 | 89.32 | 90.29 |
| 4 | 85.43 | 92.23 | 94.17 |
| 5 | 88.34 | 89.32 | 91.26 |

Larvicidal activity:

The larvicidal activity of synthesized Cu(II) complexes was studied against *C. quinquefasciatus* and the values are depicted in Tables 9 & 10.

Table 9. Larvicidal activity of Schiff base Cu (II) complexes

| Complex code | Concentration /Mortality | | | |
|--------------|--------------------------|-----------|-----------|-----------|
| | 6mg/200mL | 4mg/200mL | 2mg/200mL | 1mg/200mL |
| 1 | 16 | 11 | 7 | 3 |
| 2 | 17 | 12 | 6 | 3 |
| 3 | 16 | 10 | 6 | 2 |
| 4 | 17 | 12 | 7 | 3 |
| 5 | 17 | 11 | 7 | 4 |

Table 10. Statistical analysis of larvicidal activity of the Schiff base Cu(II) complexes

| Complex code | Concentration /Mortality \pm SD | | | | χ^2 | Df |
|--------------|-----------------------------------|---------------|---------------|---------------|----------|----|
| | 6mg/200mL | 4mg/200mL | 2mg/200mL | 1mg/200mL | | |
| 1 | 80 \pm 6.81 | 55 \pm 6.47 | 35 \pm 7.66 | 15 \pm 6.80 | 23.75 | 3 |
| 2 | 85 \pm 6.91 | 60 \pm 6.62 | 30 \pm 7.81 | 15 \pm 3.26 | 23.98 | |
| 3 | 80 \pm 6.50 | 50 \pm 6.80 | 30 \pm 6.69 | 10 \pm 6.00 | 21.86 | |
| 4 | 85 \pm 5.93 | 60 \pm 6.29 | 35 \pm 7.22 | 15 \pm 8.21 | 22.55 | |
| 5 | 85 \pm 5.82 | 55 \pm 6.66 | 35 \pm 6.71 | 20 \pm 8.24 | 18.90 | |

Mean value of triplicates; Control-Nil mortality; df- significant at $p < 7.81$

DISCUSSION:

The synthesized Schiff base Cu (II) complexes are freely soluble in DMSO, DMF and ethanol. The lower molar conductivity value of the complexes indicates their non-electrolytic nature. [22] The electronic spectrum of Cu (II) complexes were shifted to longer wave length with increasing the intensity which may be attributed to the donation of lone pair of electron present in the nitrogen atom of the Schiff base to the metal ions. The absorption band observed around 270 nm and 300 nm corresponds to $\pi \rightarrow \pi^*$ and $n \rightarrow \pi^*$ transitions respectively. [23] The green colour observed for the Cu (II) complexes are due to the presence of a broad $d \rightarrow d$ transitions in the visible region of the UV-Visible spectrum.

In the FTIR spectra an intense band appeared around 1590 cm^{-1} for the complexes corresponds to the coordination of imine nitrogen with Cu (II) ions. [24] The difference between asymmetric and symmetric stretching frequencies ($\Delta\nu = [\nu_{as}\text{ COO}^- - \nu_s\text{ COO}^-]$) were found to be higher than the corresponding free carboxylate anion. This confirmed the monodentate coordination of the carboxylate anion present in the Schiff base ligand with the Cu (II) ions. [25] The bands appeared between $418\text{--}480\text{ cm}^{-1}$ and $538\text{--}550\text{ cm}^{-1}$ are assigned to the formation of M-O and M-N coordination respectively. [26] FTIR spectral data of all the Schiff base

Cu (II) complexes revealed that the Cu (II) ion is coordinated through phenolic oxygen, imine nitrogen and oxygen atom present in the carboxylate group of the Schiff base ligand and act as a tridentate ligand. ESR spectra of the complexes displayed an isotropic spectrum with the g_{iso} values between 2.01-2.09, indicating an axial symmetry with all the principal axes aligned parallel symmetry i.e., $g_{xx} = g_{yy} = g_{zz}$. Such a spectrum is expected in complexes with octahedral geometry. [27]

All the binuclear Cu (II) complexes undergo two one-electron reduction and oxidation at different potential. Cyclic voltagrams for all the complexes are similar and the cathodic (Ipc) and anodic (Ipa) peak currents are not equal. This indicates the quasi-reversible nature of the electron transfer process. [28] Two reduction waves are obtained in the cathodic region corresponding to stepwise one-electron reductions through a $\text{Cu}^{\text{II}}\text{Cu}^{\text{I}}$ intermediate to give a binuclear $\text{Cu}^{\text{I}}\text{Cu}^{\text{I}}$ species. All the binuclear copper (II) complexes showed two quasi-reversible reduction waves. [29] The reason for the observation of two reduction waves may be due to electronic change between the copper ions. After the first one-electron reduction some of the electron density is transferred from the reduced copper ion to the other copper ion.

The thermal decomposition of the complex **1** takes place in four stages. The 1st stage takes place in a single step that is between 40 to 120°C with mass loss of 13.65 % it may due to elimination of water molecule. In the 2nd stage the decomposition occurs in the temperature range of 120 to 300° C. This range corresponds to the decomposition of 4, 4'-bipyridine ligand ^[29] and the maximum range of mass loss was recorded as 46.86 %. The 3rd stage corresponds to the decomposition of Schiff base ligand in the temperature range of 300 to 530°C. The final residue with 22.9 % was probably due to CuO. ^[29]

Schiff base Cu (II) complexes structure enhances the coordination and chelating which was tends to make more powerful and potent bacteriostatic agents, thus inhibiting the growth of the bacteria also destroying them more forcefully. ^[30] In the present study, the lower activity of some metal complexes may be due to the lower penetrating tendency of them through lipid membrane. Hence this could neither block nor inhibit the growth of the microorganism. ^[30]

The Schiff base Cu (II) complexes **1** and **2** showed very good antioxidant scavenging activity with 90.29 % and 91.26 % respectively with minimum concentration of 2 mg. The highest mortality of 85 % was obtained for the complexes **2**, **4** and **5**. The complexes **1** and **3** exhibited 80 % of mortality values. The average larval mortality data were subjected to statistical analysis for calculating standard deviation and chi-square values for synthesized Schiff base Cu (II) complexes. The calculated values were lesser than table value hence the results with $p < 7.81$ were considered to be statistically significant.

CONCLUSION:

In the present study, we have designed and synthesized a series of Schiff base Cu (II) complexes using N-(salicylidene)-L-aminoacids [L-valine, L-tryptophan, L-leucine, L-alanine and L-cysteine] and 4,4'-bipyridine. The synthesized Schiff base Cu (II) complexes were characterized by UV-Visible, FTIR, EPR, TGA and CV. The biological activities such as *in vitro* antibacterial, *in vitro* antifungal, antioxidant and larvicidal activity was assayed. All the complexes exhibited significant *in vitro* biological activities. However, further *in vivo* studies are necessary to prove their biological activities.

REFERENCES:

1. Nancy Mathew M., Sithambaresan M.R.P., Kurup., Spectral studies of copper (II) complexes of tridentate hydrazone ligands with heterocyclic compounds as coligands: X ray crystal structure of one acyl hydrazone copper (II) complex. Spectrochim. Acta A, 79, 1154-1161, (2011)
2. Chakraborty A., Kumar P., Ghosh K., Roy P., Evaluation of a Schiff base copper complex compound as potent anticancer molecule with multiple targets of action. Eur J Pharmacol, 647(1-3), 1-12, (2010)
3. Minkel D.T., Chanstier C.H., Petering D.H., Reactions of 3-Ethoxy-2-oxobutylaldehyde Bis(N4-dimethylthiosemicarbazono)-Zinc (II) with tumor cells and mitochondria. Mol Pharmacol, 12(6), 1036-1044, (1976)
4. Liberta A.E., West D.X., Antifungal and antitumor activity of heterocyclic thiosemicarbazones and their metal complexes: current status. Biometals, 5(2), 121-126, (1992)
5. Marzano C., Pellei M., Tisato F., Santini C., Copper complexes as anticancer agents. Anti-Cancer Agents Med. Chem, 9(2), 185-211, (2009)
6. Pandey A., Dewangan D., Verma S., Mishra A., Dubey R.D., Synthesis of schiff bases of 2-amino-5-aryl-1, 3, 4-thiadiazole and its analgesic, anti-inflammatory, antibacterial and antitubercular activity. Int J Chem Tech Res, 3, 178-184, (2011)
7. Chandramouli C., Shivanand M.R., Nayanbhai T.B., Bheemachari B., Udupi R.H., Synthesis and biological screening of certain new triazoleschiff bases and their derivatives bearing substituted benzothiazole moiety. J Chem Pharm Res, 4, 1151-1159, (2012)
8. Schiff H., Justus Leibigs., Eineneue Reshe Organischer Diamine (in German), Ann Chem, 140, 92-137, (1864)
9. Schiff H., Justus Liebig's., Mittheilungen aus dem universitätslaboratorium in Pisa: Eine neue reihe organischer basen, Ann Chem, 131(1), 118-119, (1864)
10. Dhar D.N., Taploo C.L., Schiff bases and their applications, J Sci Ind Res, 41(8), 501-506, (1982)
11. Kplodziej B., Grech E., Schiff W., Kamieński B., Pazio A., Woźniak K., The NMR and X-ray study of L-arginine derived Schiff bases and its cadmium complexes. J. Mol. Str, 1063,145-152, (2014)
12. Geetha K., Santha Lakshmi S., Synthesis, Characterisation and Antimicrobial studies of Dinuclear copper (II) complexes derived from pentadentate Schiff base ligand. Res. J. Chem. Sci, 4(3),68-75, (2014)
13. Verma G., Marella A., Shaquiquzzaman M., Akhtar M., Ali M.R., A review exploring biological activities of hydrazones. J Pharm Bioallied Sci, 6, 69-80, (2014)
14. Young I.S., Woodside J.V. Antioxidants in health and disease. J Clin Pathol, 54, 176-86, (2001)

15. Rice-Evans C.A., Gopinathan V., Oxygen toxicity, free radicals and antioxidants in human disease: Biochemical implications in atherosclerosis and the problems of premature neonates. *Essays Biochem*, 29, 39–63, (1995)
16. Saranya J., Sundaramurthy Santha Lakshmi., In vitro antioxidant, antimicrobial and larvicidal studies of schiff base transition metal complexes. *J.Chem. Pharm. Res*, 7(4), 180-186, (2015)
17. Ananthanarayan A., Jayaram Paniker C.K., Bisopsies of tissue. *Text book of microbiology*. 8 th Ed, University press, India, (2009)
18. Vinusha H.M., Shiva Prasad K., Chandan S., Muneera Begum., Imino-4-Methoxyphenol Thiazole Derived Schiff Base Ligands: Synthesis, Spectral Characterization and Antimicrobial Activity. *Chem. Sci. J*, 6(3), (2015)
19. Lobo V., Patil A., Phatak A., Chandra N., Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacogn Rev*, 4(8), 118-126, (2010)
20. Rahuman A.A., Gopalakrishnan G., Ghouse B.S., Arumugam S., Himalayan B., Effect of Feronialimonia on mosquito larvae, *Fitoterapia*, 7, 553-555, (2000)
21. Prasad R.N., Agrawal M., Sharma M., Synthesis, characterization, theoretical studies and biological activities of mixed ligand complexes of Cr (III), Mn(II), Fe(III), Co(II), Ni(II) and Cu(II) with 2-hydroxybenzaldehyde and oxalic acid. *J. Serb. Chem. Soc*, 67, 229-234, (2002)
22. Chinnaswamy., Jayabalakrishnan., Karuannan., Natrajan., Ruthenium (II) carbonyl complexes with triennials shift buses and their antimicrobial activity. *Transmet. Chem*, 27, 75 – 79, (2002)
23. Robert Boggess K., David Zatko A., The use of conductivity data for the structure determination of metal complexes. *J. Chem. Edn*, 52(10), 649-665, (1975)
24. Angel Garcia-Raso J., Juan Fiol., Adela Lopez-Zafra., A Jose Castro., Araceli Cabrero., Ignasi Mata., Elies Molins, Crystal structure of the copper (II) ternary complex of *N*-salicylidene-l-serinato with 2, 6-diaminopyridine.: Toxicity studies against *Drosophila melanogaster*. *Polyhedron*, 22, 403-409, (2003)
25. Nakamoto K., *Infrared and Raman Spectra of Inorganic and Coordination Compounds*. 5th Edition, Wiley-Interscience Publication, New York, (1978)
26. Santha lakshmi S., Geetha K., Mahadevi P., Tridentate Schiff base (ONO) transition metal complexes: Synthesis, crystal structure, spectroscopic and larvicidal studies. *J. Chem. Sci*, 128, 1113-1118, (2016)
27. Kannappan R., Mahalakshmy R., Rajendiran T.M., Venkatesan R., Sambasiva Rao P., Magnetic, catalytic, EPR and electrochemical studies on binuclear copper (II) complexes derived from 3,4-disubstituted phenol. *Indian Academy of Sciences*, 115(1), 1-14, (2003)
28. Prabu R., Vijayaraj A., Jagadish L., Kaviyaranan V., Narayanan V., New Unsymmetric Dinuclear Copper (II) Complexes of Trans-disubstituted Cyclam Derivatives: Spectral, Electrochemical, Magnetic, Catalytic, Antimicrobial, DNA Binding and Cleavage Studies. *Bull. Korean Chem. Soc*, 32(5), 1669-1678, (2011)
29. Samik Gupta., Sachindranath Pal., Anil Kumar barik., Arijit Hazra., Somnath Roy., Tarak Nath Mandal., Shie-Peng., Gene-Hsiang Lee., Salah El Fallah., Javier Tercero., Susanta Kumar Kar., Synthesis, characterization and magnetostructural correlation studies on three binuclear copper complexes of pyrimidine derived Schiff base ligands. *Polyhed*, 27, 2519-2528, (2009)
30. Da Silva C.M., da Silva D.L., Modolo L.V., Alves R.B., de Resende M.A., Martins C. V.B., de Fátima Â., Schiff bases: A short review of their antimicrobial activities, *J of Advan. Res*, 2(1), 1–8, (2011)