

RESTORATION OF SAND DUNE VEGETATION FROM CUDDALORE AREA, SOUTHEAST COAST OF INDIA

M.P.Arulmoorthy*, M.Srinivasan¹

^{1,2} Centre of Advanced Study in Marine Biology, Faculty of Marine Sciences, Annamalai University,
Parangipettai – 608 502 Tamil Nadu, India

*Corresponding Author Email: mparulguru@gmail.com

ABSTRACT

Coastal sand dunes (CSD) florae were under constant anthropogenic and natural pressure due to rapid elimination of sand dunes and its associated vegetation; as a result, its associated indigenous knowledge with them is also gradually disappearing. Such biodiversity rich and useful ecosystems need immediate restoration and conservation actions. Cuddalore coastal area is prone to both anthropogenic and natural disaster. Cyclone Thane hit Cuddalore coast on 29th and 30th of December 2011 with wind speeds of up to 135 kmph (83 mph) and tidal surges reaching 1.5 metres (5 feet), is worth mentioning apart from the tsunami hit during December 2004. Industrialization has occupied nearly 500 acres of Coastal land which causes pollution and destruction of sand dune vegetation. Restoration of degraded area by propagation of plants (*Ipomoea pescaprae* and *Spinifex littoralis* (which are natural sand binders) by plucking a portion of the creeper from the denser area and planting it in pits dug at a depth of 30 to 40 cm. The planting of creepers was made at an interval of 2 m distance each in 6 pits. The restoration work was started from October 2012, 90 % survival was found during the restoration study. The best season for this program in this area was between October to January. After three months, 30cm of growth was observed in the plants. As that of mangroves restoration programmes this sand dune vegetation flora should also be encouraged by all the countries in the world.

KEY WORDS

Coastal Sand dunes, cuddalore, pollution, southeast coast.

INTRODUCTION

Coastal sand dunes are natural structures which protect the coastal environment by absorbing energy from wind, tide and wave action. Despite geographical differences, sand dunes have been considered as a specific ecosystem due to several common environmental features. CSD constitute a variety of microenvironments due to substrate mobility and physical processes. Plants establishing on coastal sand dunes are subjected to several environmental fluctuations which affect their growth, survival and community structure. CSDs dynamic but fragile buffer zones of sand and vegetation where the

following three characteristics can be found: large quantities of sand; persistent wind capable of moving the sand; suitable locations for sand to accumulate.

CSD formulations ultimately depend on embayment size and prevailing wind energy (Kumar et al.1993). Their heights differ in response to adequate sand supply, climate and local topographic features (Barbour et al. 1985). Plants on coastal dunes are specially adapted to withstand various environmental stresses which allow them to grow, establish and to trap sand in such harsh conditions of coastal zones, so they are mostly represented by herbs, shrubs, creepers or runners (Sridhar et al. 2007)

The role of vegetation in dune formation is critical and is that of a wind trap, sand binder and dune stabilizer (Wagner 1964; Dahm et al. 2005). The foliage of dune plants breaks wind activity leading to less erosive activity on the lee side (Chapman 1976). Pioneer zone, intermediate zone and back/ forests zone were recognized earlier in coastal dunes and later several workers found shore, foredune, main dune width wind ward and lee ward slopes, wet dune slacks and back dunes with plateaus, holes that supporting grasslands scrub forests, thus portraying complex ecosystem diversity (Wood house 1978; Hesp 2004). Temperate coastal dunes are well studied and documented (Koske and gemma 1997; Sridhar and Bhagya 2007) as compared to studies on tropical coastal dunes (Kulkarni et al. 1997; Sridhar and Bhagya 2007).

The Ecological roles and functions of coastal dunes include: essential store of sediments, protecting the land behind them from storm erosion and potential sea level rise; filter for rainwater and groundwater and in some situations, provided aquatic habitats such as dune lakes; protection of islands from storm surges, hurricanes and erosion; trapping of the windblown sand and prevention of sand being blown further inland by the vegetation; habitats for specially adapted plants, birds, and animals - several of which are now rare or endangered; a range of unique landforms and processes which have intrinsic value and are of scientific interest; and nesting sites for sea turtles and birds.


This paper aims to generate a restoration of *Ipomoea pescaprae* and *Spinifex littoralis* on coastal sand dune in the coramandal coast, with special reference to Cuddalore coastline. Apparently, this is the first sand dune restoration work carried out in Cuddalore coast.

MATERIALS AND METHODS

Study area

Cuddalore is located on the Coramandal coast in 11°44'39" N and 79°47'00" E, lies 25km (16mi) south of Pondicherry and coast of the bay of bengal . *Devanampattinam, Thazhanguda, Singarathoppu, Pudupettai, Annangkovil* are the coastal village with sand dune coverage of about 30 km² extent, present at a distance of about 48 km towards south on the way to Parangipettai from the Cuddalore main town (**Map:1**).

The coastal border has a length of 30 km and a breadth ranging from four to six hundred meters. Superficially, the coast is flat and sandy. The coastal zone of Cuddalore comprises newer and older dunes including saline areas of clayey texture. The study area experiences mean annual temperature of 30.0 °C and mean annual rainfall about 1,311-1,172 mm. The mean number of annual rainy days is 55, the mean monthly temperature ranges from 21.3°- 30.2°C. The climate is tropical dissymmetric with the bulk of the rainfall during northeast monsoon October-December (Indian Meteorological Department - Chennai).



Map 1: Showing Cuddalore coastal area

Preliminary survey

A preliminary survey was carried out in *Devanampattinam, thazhanguda, Singarathoppu,*

Pudupettai, Annangkovil coastal villages in Cuddalore. Due to high industrial development, construction of thermal power plant in Puthupettai and harbor

construction in annankovil affects the coastal sand dunes and destroys the sand dune plants (*Ipomea pescaprae* and *Spinifex littoralis*) in high rate. In my

survey 30-40% of sand dunes in Cuddalore coast are disturbed by the development activities carried out in the coastal areas and industrialation.

Devanampattinam



Fig:1 Sand dune with *ipomea pescaprae* affected by the artificial shield build by humans



Fig:2 *spinifex littoralis* destruction due to human impact

Thazhanguda



Fig:3 Sand dune destroyed by the cyclone "thane"



Fig:4 sand dune destructured by human impact

Singarathoppu



Fig:5 Sand dune with *ipomea pescaprae*



Fig:6 sand dune with temple

Pudupettai



Fig:7 Grazing of *ipomea pes-caprae* in sand dune



Fig:8 sand dune with *cashewrina* and *ipomea*

Annangkovil



Fig:9 Grazing of *ipomea pes-caprae* by the cattles in sand dunes



Fig:10 Destruction of sand dunes by harbor construction

Restoration

Restoration work was carried out in Devanampattinam, Thazhanguda, Singarathoppu, Pudupettai and Annangkovil coastal villages in Cuddalore coast from the month of October 2012. In each villages two to three sand dunes were selected for *ipomea pes-caprae* and *spinifex* restoration. Restoration of degraded area by propagation of plants by plucking a portion of plant from the denser

area i.e., two to three meter length of *ipomea pes-caprae* were plucked and cut into small plants by giving two to three internodes and planting it in pits dug at a depth of 30 to 40 cm. The planting of creepers was made at an interval of 2 m distance each in 6 pits. For one site an area of 16 m X 16 m was covered with fence. Watering the plants and fencing were followed after the planting.



Fig:11 Dugging of pits



Fig:12 Planting ipomea pescaprae



Fig:13 Twenty days growth



Fig:14 Watering to ipomea and spinifex



Fig: 15, 16 Fencing of ipomea vegetated sand dunes in pudupettai and singarathoppu villages

RESULTS AND CONCLUSION

Field investigation and the preliminary assessment in the Cuddalore coastal villages indicate the sand dunes were in the immense pressure. In *Devanampattinam* (11°44'39" N and 79°47'00" E) totally 12 sand dunes were surveyed, in this 6 to 7 dunes are without any sand dune plants and are in the stage of extinct (Fig :1,2). In *thazhanguda* (11°46'02" N and 79°47'27" E) totally 5 dunes were surveyed, in this all Sand dunes

are without any vegetation (Fig:3, 4). In *singarathoppu* (11°43'26" N and 79°46'57" E) and *pudupettai* (11°30'59" N and 79°46'09" E) 12 sand dunes were surveyed in this mostly all sand dunes in *singarathoppu* with high vegetation of ipomea and spinifex (Fig: 5, 6) and in pudupettai due to the thermal power plant construction all sand dunes are destroyed and polluted by the humans (Fig: 7, 8). *Annangkovil* (11°30'08" N and 79°46'19" E) nearly all

sand dunes are disturbed by harbor construction with poor vegetation (**Fig: 9, 10**).

Restoration works were carried out in five coastal villages in Cuddalore from the month October 2012 (**Fig: 11, 12, 13, and 14**). *Ipomea pescaprae* and *Spinifex littoralis* are the plant species restored in these villages by plucking a portion of the creeper from the denser area and planting it in pits dug at a depth of 30 to 40 cm. The planting of creepers was made at an interval of 2 m distance each in 6 pits. 90% of growth was recorded in my study from the month of October 2012 to January 2013. *Ipomea pescaprae* growth was reached nearly 30 cm from three months but *Spinifex littoralis* was not showing any level of growth. After two months, fencing of the vegetated sand dunes were made particularly in

pudupettai and singarathoppu villages (**Fig: 15, 16**). The table (**Table: 1, 2**) shows the growth rate of *Ipomea pescaprae* and *Spinifex littoralis* for the three months in the five villages.

In my study, sand dunes in Cuddalore coastal area is under pressure due to the development activities carried out in this area. Sand dune vegetation is the good way for protecting sand dunes in Cuddalore coastal areas. *Ipomea pescaprae* is a dune plant has great ability to grow in the Cuddalore area. Turtle nesting activity is carried out in high level in Cuddalore coastal area. For to conserve the coastal living organisms and protecting coastal villages from natural hazards, the sand dunes should be conserve and manage by routine monitoring and restoration activity.

Table: 1: *Ipomea pescaprae* plant growth in five villages from October 2012 to January 2013

PLANT SPECIES	VILLAGES	MONTH	GROWTH RATE(cm)
<i>Ipomea pescaprae</i>	Devanampattinum	October 2012	5-10
		November 2012	10-17
		December 2012	17-23
		January 2013	23-30
	Thazhanguda	October 2012	5-9
		November 2012	10-19
		December 2012	19-23
		January 2013	23-30
	Singarathoppu	October 2012	5-10
		November 2012	10-15
		December 2012	17-21
		January 2013	21-28
	Pudupettai	October 2012	5-8
		November 2012	10-16
		December 2012	17-24
		January 2013	24-27
	Annangkovil	October 2012	5-7
		November 2012	8-17
		December 2012	17-24
		January 2013	24-27

Table: 2: Spinifex littoralis plant growth in five villages from October 2012 to January 2013

PLANT SPECIES	VILLAGES	MONTH	GROWTH RATE(cm)
<i>Spinifex littoralis</i>	Devanampattinam	October 2102	-
		November2012	-
		December2012	-
		January 2013	0.5-1.5
	Thazhanguda	October 2102	1.5-2.0
		November2012	-
		December 2012	-
		January 2013	-
	Singarathoppu	October 2102	-
		November2012	-
		December2012	-
		January 2013	-
	Pudupettai	October 2102	-
		November2012	-
		December2012	-
		January 2013	1.0-2.3
	Annangkovil	October 2102	-
		November2012	-
		December2012	-
		January 2013	-

ACKNOWLEDGEMENT

The Authors are thankful to University Grants Commission (UGC) for their financial support. The authors also thank the authorities of Faculty of Marine Sciences, CAS in Marine Biology, Annamalai University for their support and encouragement.

REFERENCES

- Arun, A.B., K.R. Beena, N.S. Raviraja and K.R. Sridhar. 1999. Coastal sand dunes - A neglected ecosystem. *Current Science* 77: 19-21.
- Barbour, M.G., T.M. De Jong and B.M. Palvik. 1985. Marine beach and dune plant communities. Physiological ecology of North Americal communities. *Restoration Ecology* 6: 59-68.
- Donatiu, P. 2000. First Line of Defence, in the Growing Idea. Spring Edition. Greening Australia Queensland, Brisbane.
- Hesp, 2004. Coastal dunes in the Tropics & temperate regions: Location, formation, morphology and vegetation process; p.29-65 In: M.L. Martínez and N.P. Psuty (ed.). *Coastal dunes: Ecology and Conservation*. Berlin: Springer-Verlag.
- Koske R.E, and J.N Gemma. 1997. Mycorrhizae and succession in plantings of beachgrass in sand dunes. *American Journal of Botany* 84: 118-130.
- Kulkarni, S.S., N.S. Raviraja and K.R. Sridhar. 1997. Arbuscular mycorrhizal fungi of tropical sand dunes of west coast of India. *Journal of Coastal Research* 13: 931-936.
- Kumar M, E. Goossens and R.Goossens. 1993. Assessment of sand dune change detection in Rajasthan (Thar) Desert. *International Journal of Remote Sensing* 14(9): 1689-1703
- Sridhar K.R, and B. Bhagya. 2007. *Coastal sand dune vegetation: a potential source of food, fodder and pharmaceuticals*. Electronic database available at <http://www.lrrd.org/lrrd19/6/srid19084.htm>. Captured on 19 June 2009.
- Wagner, R.H. 1964. The Ecology of dunes – strand habitat of North Carolina. *Ecological Monographs* 34: 79-96.
- Wood house, W.W. 1978. Dune building and stabilization with vegetation. *U.S Army crop of engineers* 3: 9-104.



***Corresponding Author:**

Arulmoorthy*

Centre of Advanced Study in Marine Biology,
Faculty of Marine Sciences, Annamalai University,
Parangipettai – 608 502 Tamil Nadu, India